

GRAPHS WITHOUT INDUCED P_5 AND C_5 *

GABOR BACSÓ AND ZSOLT TUZA[†]

Computer and Automation Institute
Hungarian Academy of Sciences
1111 Budapest, Kende u. 13–17, Hungary

Abstract

Zverovich [Discuss. Math. Graph Theory **23** (2003), 159–162.] has proved that the domination number and connected domination number are equal on all connected graphs without induced P_5 and C_5 . Here we show (with an independent proof) that the following stronger result is also valid: Every P_5 -free and C_5 -free connected graph contains a minimum-size dominating set that induces a complete subgraph.

Keywords: graph domination, connected domination, complete subgraph, forbidden induced subgraph, characterization.

2000 Mathematics Subject Classification: 05C69.

1. The Results

For a (simple, undirected) graph G with vertex set $V(G)$ and edge set $E(G)$, a subgraph $H \subseteq G$ is called a *dominating subgraph* if every $v \in V(G) \setminus V(H)$ is adjacent to some $w \in V(H)$. The *domination number* of G is the smallest number of vertices in a dominating subgraph H of G ; and the *connected-domination number* is the minimum under the further requirement that the dominating subgraph $H \subseteq G$ be *connected*.

More than a decade ago, the present authors [1] and independently Cozzens and Kelleher [4] proved that a connected graph without induced paths and cycles on five vertices contains a dominating *complete subgraph*.

*Research supported in part by the grant OTKA T-032969.

[†]Also affiliated with the Department of Computer Science, University of Veszprém, Hungary.

Recently, Zverovich [6] showed that if such a graph has domination number k , then it also contains a *connected* dominating subgraph on k vertices. In this note we prove the following common generalization of these two results:

Theorem 1. *If a connected graph without induced P_5 and C_5 subgraphs has domination number k , then it is dominated by a complete subgraph on k vertices.*

This theorem will be proved in Section 2. Let us note that it has the following equivalent formulation, which is in fact a *necessary and sufficient condition*:

In each connected, induced subgraph of a graph G , the minimum number of vertices in a complete subgraph dominating G exists and is equal to the domination number, if and only if no induced subgraphs of G are isomorphic to P_5 and C_5 .

The reason is that P_5 and C_5 themselves do not have dominating cliques; i.e., if the property is required hereditarily, then P_5 and C_5 have to be forbidden induced subgraphs. Recalling from [6] that both P_5 and C_5 have domination number 2 and connected-domination number 3, also the following characterization theorem of [6] follows:

In each connected, induced subgraph of a graph G , the connected-domination number is equal to the domination number, if and only if G contains no induced subgraphs isomorphic to P_5 and C_5 .

In a more general setting, a theory has been developed for solving graph-class equations of the form $Dom(\mathcal{D}) = Forb(\mathcal{F})$ where $Dom(\mathcal{D})$ consists of the graphs in which every connected, induced subgraph is dominated by some induced subgraph $D \in \mathcal{D}$, and $Forb(\mathcal{F})$ means that no induced subgraph $F \in \mathcal{F}$ may occur. For recent results and more references of this kind, please see [2].

We also consider here the other extreme, i.e., domination with *largest* complete subgraphs. In this context, a subclass of P_5 -free graphs has been studied in [3]. It is proved there that if a connected graph contains at least one triangle but it is $2K_2$ -free (i.e., it does not contain any pair of disjoint edges as an induced subgraph), then it is dominated by some of its complete subgraphs of *maximum size*. As noted in [1], this result does not remain valid for P_5 -free (and C_5 -free) graphs. A simple counterexample is the graph, that we denote by P_5^+ , obtained from P_5 by joining the middle vertex with one of the endpoints. In Section 3 we prove:

Theorem 2. *If a connected graph does not contain P_5 , C_5 , and P_5^+ as an induced subgraph, then it is dominated by a complete subgraph of maximum size.*

Similarly to Theorem 1, also this result can be transformed to a necessary and sufficient condition:

Every connected, induced subgraph H of a graph G is dominated by a maximum-size complete subgraph of H , if and only if G contains no induced subgraph isomorphic to P_5 , C_5 , and P_5^+ .

2. Small Complete Subgraphs

Before proving the main result of the paper, we state an auxiliary observation. In the proof we shall use the following terminology. If H is a dominating subgraph, a *private neighbor* of a vertex $w \in V(H)$ is a vertex z in $V(G) \setminus V(H)$ such that w is the unique neighbor of z inside $V(H)$.

Lemma 1. *Suppose that G contains no induced P_5 and C_5 , and let H be a dominating induced subgraph of G . If H has a connected component H' which is not complete, then there exists a vertex $x \in V(H')$ such that*

1. $H' - x$ is connected, and
2. $H - x$ is dominating in G .

Proof. Let x and x' be two vertices at maximum distance apart in the induced subgraph H' . (It will be irrelevant whether or not their distance in G is smaller than in H' .) If H' is not complete, then x and x' are non-adjacent. Let us select a shortest x - x' path P in H' .

By the choice of x and x' , both $H' - x$ and $H' - x'$ are connected. If some of them dominates G , then the lemma is proved. Otherwise, each of x and x' has at least one private neighbor, say y and y' , respectively. In this case, however, the subgraph induced by $V(P) \cup \{y, y'\}$ either is a 5-cycle or contains an induced P_5 , a contradiction. ■

Proof of Theorem 1. Given the graph G without induced P_5 and C_5 , we select a dominating subgraph H such that

1. $|V(H)|$ is minimum, and

2. H has as few connected components as possible.

By Lemma 1, each component of H is a complete subgraph of G . We need to prove that H has just one component. Suppose, for a contradiction, that H is disconnected.

Case 1. Some vertex has neighbors in more than one component of H . Let w be such a vertex. Then the subgraph induced by $V(H) \cup \{w\}$ in G is dominating, and has fewer components than H . Since the component containing w is not complete, Lemma 1 allows us to remove a non-cutting vertex from this component, hence obtaining another dominating induced subgraph that has the same number of vertices as H , but with fewer components. This contradicts the choice of H .

Case 2. Each vertex has all of its neighbors in the same component of H . Since H is dominating and disconnected, but G is connected, there exists an edge $w'w''$ inside $G - H$, and two distinct components K', K'' of H , such that w' has neighbors in K' and w'' has neighbors in K'' . Then the subgraph H^+ induced by $V(H) \cup \{w', w''\}$ is dominating, and one of its components, say H' , is induced by $V(K') \cup V(K'') \cup \{w', w''\}$. This H' is not complete, and in order to keep it connected and make it complete, one needs to remove at least two vertices (one from $V(K') \cup \{w''\}$ and one from $V(K'') \cup \{w'\}$). On the other hand, H^+ contains just two more vertices than one of the *smallest* dominating subgraphs, H , therefore applying Lemma 1 twice we obtain a minimum-size dominating subgraph H^- . This H^- has fewer components than H , again a contradiction. ■

3. Large Complete Subgraphs

In the proof below, the notation $N(u)$ means the “open neighborhood” of vertex u (i.e., the set of vertices adjacent to u).

Proof of Theorem 2. We prove the assertion by contradiction. Let G be a graph without induced P_5 , C_5 , and P_5^+ , which is a minimal counterexample in the sense that, for every vertex v , if the subgraph $G - v$ is connected, then it is dominated by some of its maximum-size complete subgraphs. We denote by ω the maximum clique-size in G .

Choose any K_ω in G . Let v be a vertex at maximum distance from this K_ω . Since G is assumed to be a counterexample, this distance is at least 2.

Clearly, $G - v$ is connected and still has maximum clique size ω . Therefore, by the minimality of G , it is dominated by a subgraph $K \simeq K_\omega$. This K does not dominate v , but v has some neighbor, say w , dominated by K . Since K is of maximum size, $V(K) \setminus N(w) \neq \emptyset$.

Let us observe next that $V(K) \setminus N(w)$ consists of just one vertex. Indeed, otherwise we could select two vertices in $V(K) \setminus N(w)$ and one in $V(K) \cap N(w)$, hence obtaining an induced subgraph (the selected triangle with v and w) isomorphic to P_5^+ , a contradiction.

We denote by y the single vertex of $V(K) \setminus N(w)$. It follows that the set $X = V(K - y) \cup \{w\}$ also induces K_ω in G . If it is not dominating, then some vertex z is adjacent to y and nonadjacent to the entire X . We finally choose a vertex $x \in V(K) \cap N(w)$. If vz is not an edge, then v, w, x, y, z induce P_5 ; and if vz is an edge, then they induce a C_5 . In either case, we obtain a contradiction that proves the theorem. ■

Note added in February 2004. One of the referees has informed us that Theorem 1 has been discovered independently by Goddard and Henning [5]; and another referee has observed that it can also be derived from the result of Zverovich by an argument avoiding Lemma 1.

References

- [1] G. Bacsó and Zs. Tuza, *Dominating cliques in P_5 -free graphs*, Periodica Math. Hungar. **21** (1990) 303–308.
- [2] G. Bacsó and Zs. Tuza, *Structural domination of graphs*, Ars Combinatoria **63** (2002) 235–256.
- [3] F.R.K. Chung, A. Gyárfás, W.T. Trotter and Zs. Tuza, *The maximum number of edges in $2K_2$ -free graphs of bounded degree*, Discrete Math. **81** (1990) 129–135.
- [4] M.B. Cozzens and L.L. Kelleher, *Dominating cliques in graphs*, Discrete Math. **86** (1990) 101–116.
- [5] W. Goddard and M.A. Henning, *Total domination perfect graphs*, to appear in Bull. ICA.
- [6] I.E. Zverovich, *Perfect connected-dominant graphs*, Discuss. Math. Graph Theory **23** (2003) 159–162.

Received 21 July 2003
Revised 5 February 2004