

ESTIMATION OF CUT-VERTICES IN  
EDGE-COLOURED COMPLETE GRAPHS

ADAM IDZIK

*Akademia Świętokrzyska*  
*15 Świętokrzyska street, 25-406 Kielce, Poland*  
and

*Institute of Computer Science*  
*Polish Academy of Sciences*  
*21 Ordonia street, 01-237 Warsaw, Poland*

**e-mail:** adidzik@ipipan.waw.pl

All graphs considered here are finite simple graphs, i.e., graphs without loops, multiple edges or directed edges. For a graph  $G = (V, E)$ , where  $V$  is a vertex set and  $E$  is an edge set, we write sometimes  $V(G)$  for  $V$  and  $E(G)$  for  $E$  to avoid ambiguity. We shall write  $G \setminus v$  instead of  $G_{V \setminus \{v\}} = (V \setminus \{v\}, E \cap 2^{V \setminus \{v\}})$ , the subgraph induced by  $V \setminus \{v\}$ . A vertex  $v \in V(G)$  is called a *cut-vertex* of  $G$  if  $G$  is connected and  $G \setminus v$  is not. By a *k-edge-colouring* of a graph we mean any finite partition of the set of its edges into  $k$  subsets. A graph  $(V, E)$  with a given  $k$ -edge-colouring  $(E^1, \dots, E^k)$  ( $E^i \cap E^j = \emptyset$  for  $i \neq j$ ;  $i, j \in \{1, \dots, k\}$  and  $\bigcup_{i \in \{1, \dots, k\}} E^i = E$ ) is denoted by  $(V, E^1, \dots, E^k)$ . The graphs  $(V, E^i)$  are called monochromatic subgraphs of  $(V, E^1, \dots, E^k)$ ,  $i \in \{1, \dots, k\}$ . As usual, by  $K_m$  we denote the complete graph with  $m$  vertices.

Let  $c(G^i)$  denote the number of cut-vertices of  $G^i$  in a monochromatic subgraph  $G^i = (V, E^i)$  of a  $k$ -edge-coloured complete graph  $K_m = (V, E^1, \dots, E^k)$  ( $i \in \{1, \dots, k\}$ ).

Given a  $k$ -edge-coloured graph  $G = (V, E^1, \dots, E^k)$ , we define  $F^i = E \setminus E^i$ ,  $G^i = (V, E^i)$ ,  $\bar{G}^i = (V, F^i)$ , where  $E = \bigcup_{i \in \{1, \dots, k\}} E^i$  and  $i \in \{1, \dots, k\}$ . Here  $G^i$  is a monochromatic subgraph of  $G$  and  $\bar{G}^i$  its complement in  $G$ .

**Theorem** (Idzik, Tuza, Zhu). *Let  $(E^1, \dots, E^k)$  be a  $k$ -edge-colouring of  $K_m$  ( $k \geq 2$ ,  $m \geq 4$ ), such that all the graphs  $\bar{G}^1, \dots, \bar{G}^k$  are connected.*

- (i) *If one of the subgraphs  $G^1, \dots, G^k$  is 2-connected, say  $G^i$ , then  $c(\bar{G}^i) \leq m - 2$  and  $c(\bar{G}^j) = 0$  for  $j \neq i$  ( $i, j \in \{1, \dots, k\}$ ).*
- (ii) *If none of the graphs  $G^1, \dots, G^k$  is 2-connected, and one of them is connected, say  $G^i$ , then  $c(\bar{G}^i) \leq 2$  ( $i \in \{1, \dots, k\}$ ).*
- (iii) *If none of the graphs  $G^1, \dots, G^k$  is 2-connected, and one of them is disconnected, say  $G^i$ , then  $c(\bar{G}^i) \leq 1$  ( $i \in \{1, \dots, k\}$ ).*

**Problem.** Let  $(E^1, \dots, E^k)$  be a  $k$ -edge-colouring of  $K_m$  ( $k \geq 2$ ,  $m \geq 4$ ). What is the cardinality of the set of the sum of cut-vertices of  $\bar{G}^i$  in the case none of  $G^i$  is 2-connected and (a) two of  $G^i$  are connected or (b) two of  $G^i$  are disconnected and  $c(\bar{G}^i) = 1$  ( $i \in \{1, \dots, k\}$ ) ?

Observe that in both cases (a) and (b) all the graphs  $\bar{G}^1, \dots, \bar{G}^k$  are connected.

This problem is related to some theorems presented in [1] and [2].

## References

- [1] J. Bosák, A. Rosa and Š. Znám, *On decompositions of complete graphs into factors with given diameters*, in: P. Erdős and G. Katona, eds., *Theory of Graphs, Proceedings of the Colloquium Held at Tihany, Hungary* (Academic Press, New York, 1968) 37–56.
- [2] A. Idzik and Z. Tuza, *Heredity properties of connectedness in edge-coloured complete graphs*, *Discrete Math.* **235** (2001) 301–306.

Received 21 November 2003