

## ON A SPHERE OF INFLUENCE GRAPH IN A ONE-DIMENSIONAL SPACE

ZBIGNIEW PALKA AND MONIKA SPERLING

*Department of Algorithmics and Programming*  
*Adam Mickiewicz University*  
*Umultowska 87, 61–614 Poznań, Poland*

**e-mail:** palka@amu.edu.pl

**e-mail:** dwight@amu.edu.pl

### Abstract

A sphere of influence graph generated by a finite population of generated points on the real line by a Poisson process is considered. We determine the expected number and variance of societies formed by population of  $n$  points in a one-dimensional space.

**Keywords:** cluster, sphere of influence graph.

**2000 Mathematics Subject Classification:** Primary 60D05;  
Secondary 60C05, 05C80.

## 1. Introduction

Let  $X = \{X_1, X_2, \dots, X_n\}$  be the set of  $n$  points of  $R^d$  chosen randomly and independently with the same probability. Let

$$r(X_i) = \min_{X_j \in X \setminus \{X_i\}} d(X_i, X_j)$$

denote the minimum distance between  $X_i$  and any other point in  $X$ . The open ball

$$B_i = \left\{ X \in R^d : d(X_i, X) < r_i \right\}$$

with center  $X_i$  and radius  $r_i$  is the *sphere of influence graph* at  $X_i$  ( $i = 1, \dots, n$ ). The random sphere of influence graph  $SIG(X)$  has vertex set  $X$  with edges corresponding to pairs of intersecting spheres of influence.

In other words two vertices, say  $X_i$  and  $X_j$ , are connected by an edge if and only if

$$r(X_i) + r(X_j) > d(X_i, X_j).$$

The definition of the sphere of influence graph was introduced in [10] by Touissant. These graphs have been widely investigated recently. It is known that on the Euclidean plane the sphere of influence graph always has a vertex of degree at most 18 (see [5], for related results see [1, 7]). Füredi [4] showed that the expected number of edges  $E(n, \mathcal{N})$  of the random sphere of influence graph on  $n$  vertices in normed space  $\mathcal{N}$  is equal to

$$E(n, \mathcal{N}) = C(d)n + o(n),$$

where  $C(d)$  is a constant depending only on the dimension of the space and

$$\frac{\pi}{8}2^d < C(d) < \left(1 + \frac{1}{2d}\right)\frac{\pi}{8}2^d.$$

This result was also proved independently by Chalker et al in [2]. In [6] Hitczenko, Janson and Yukich proved analogue result for variance. They showed

$$c(d)n \leq \text{Var}(n, \mathcal{N}) \leq C(d)n,$$

where constants  $c(d)$  and  $C(d)$  depend only of the space dimension.

Consider a population of  $n$  points generated by some random process in  $R^d$  and its resulting sphere of influence graph. We thereby generate clusters of points that are connected by edges. We call these clusters *societies*. The following questions arise:

- Let  $M$  denote the number of societies formed. What is the distribution of  $M$ ?
- Let  $N$  denote the size of society, i.e., the number of individuals (points) in a society. What is the distribution of  $N$ ?
- Form the convex hull of each society. What is
  - the content (area, volume) covered by a society?
  - the fraction of  $R^d$  that is contained in some society, as  $n \rightarrow \infty$ ?

In this paper our main concern is with the random variable  $M$ .

## 2. One-Dimensional Societies

Let the population consist of  $n$  points,  $X_i, 1 \leq i \leq n$ , generated on the real line by a Poisson process. Let  $X_{(i)}$  denote the corresponding order statistics and let

$$A_i = X_{(i+1)} - X_{(i)}, \quad 1 \leq i \leq n - 1,$$

denote the lengths of the spacings between adjacent points. Societies are determined by the relative magnitudes of the spacings. The  $A_i$  are identically distributed. Moreover, the distribution of the vector of ranks of the  $A_i$  is discrete uniform.

Consider now the number of societies  $M$  formed by a population of  $n$  points. Clearly,  $M$  satisfies  $1 \leq M \leq \lfloor \frac{n}{2} \rfloor$ . For fixed  $n$ , let

$$P_n(M = m) = P_n(m)$$

denote the distribution of  $M$ . Obviously,  $P_2(1) = P_3(1) = 1$ .

The following technical lemma will be helpful in the proof of the main theorem.

**Lemma 1.** *Let  $2 \leq m \leq \lfloor \frac{n}{2} \rfloor$ . If for  $n \geq 4$*

$$P_n(m) = \sum_{i=2}^{n-2} \frac{1}{4} P_i(1) P_{n-i}(m-1)$$

and for  $n \geq 2$

$$P_n(1) = (n-1)2^{2-n}$$

then

$$P_n(m) = 2^{2-n} \binom{n-1}{2m-1}$$

for  $n \geq 2$ .

**Proof.** Let  $m = 2$ . Then

$$\begin{aligned} P_n(2) &= \sum_{i=2}^{n-2} \frac{1}{4} P_i(1) P_{n-i}(1) = \sum_{i=2}^{n-2} \frac{1}{4} 2^{2-i} (i-1) 2^{2-n+i} (n-i-1) \\ &= 2^{2-n} \sum_{i=1}^{n-3} i(n-i-2) = 2^{2-n} \binom{n-1}{3}. \end{aligned}$$

Assume that lemma is true for  $m \leq j$  and let  $m = j + 1$ . Then by induction

$$\begin{aligned} P_n(j+1) &= \sum_{i=2}^{n-2} \frac{1}{4} P_i(1) P_{n-i}(j) = \sum_{i=2}^{n-2} \frac{1}{4} 2^{2-i}(i-1) 2^{2-n+i} \binom{n-i-1}{2j-1} \\ &= 2^{2-n} \sum_{i=1}^{n-3} i \binom{n-i-2}{2j-1} = 2^{2-n} \binom{n-1}{2j+1} \end{aligned}$$

which completes the proof.  $\blacksquare$

**Theorem 2.** Let  $E_n(M)$  and  $Var_n(M)$  denote the mean and the variance of the number of societies formed in a population of  $n$  individuals. Then

$$E_n(M) = \begin{cases} 2 & \text{for } n = 2, \\ \frac{n+1}{4} & \text{for } n \geq 3, \end{cases}$$

and

$$Var_n(M) = \begin{cases} 0 & \text{for } n = 2, 3, \\ \frac{n-1}{16} & \text{for } n \geq 4. \end{cases}$$

**Proof.** Let us assume that  $A_{n-1} \geq A_{n-2}$ . Then independently from the value of  $A_{n-3}$ , vertices  $X_{(n-1)}$  and  $X_{(n-2)}$  are connected by an edge. So by the above assumption the number of societies formed by population of  $n$  points is equal to one with probability

$$\frac{1}{2} P_{n-1}(1).$$

Now, let  $A_{n-1} < A_{n-2}$ . In this case the existence of only one society formed by  $n$  points, under condition that first  $n-2$  points formed one society, depends on lengths  $A_{n-3}, A_{n-2}, A_{n-1}$ . Notice that two vertices  $X_{(n-1)}$  and  $X_{(n-2)}$  are not connected by an edge if the following inequality holds

$$A_{n-1} + A_{n-3} < A_{n-2}.$$

Assume that  $A_{n-3} + A_{n-2} + A_{n-1} = l$ . Then the probability of the event

$$A_{n-2} > \frac{1}{2}l,$$

i.e., probability that vertices  $X_{(n-1)}$  and  $X_{(n-2)}$  are not connected by an edge, is equal to

$$P\left(A_{n-2} > \frac{1}{2}l\right) = \frac{\frac{1}{8}l^2}{\frac{1}{2}l^2} = \frac{1}{4}.$$

Thus we obtain that if  $A_{n-1} < A_{n-2}$ , the number of societies formed by population of  $n$  points is equal to one with probability

$$\frac{1}{2}P_{n-1}(1) - \frac{1}{4}P_{n-2}(1).$$

Consequently population of  $n$  individuals forms one society with the probability

$$P_n(1) = P_{n-1}(1) - \frac{1}{4}P_{n-2}(1).$$

Solving this recurrence equation and considering boundary conditions we obtain

$$(*) \quad P_n(1) = (n - 1)2^{2-n}, \quad n \geq 2.$$

Let  $B_i$  denote the event that two vertices, say  $X_{(i)}$  and  $X_{(i+1)}$ , are the first ones that are not connected by an edge in the sphere of influence graph. It means that the number of societies formed by population of first  $i$  vertices is equal to one, while population of first  $i + 2$  vertices form two societies and the number of societies formed by population of last  $n - i$  points is equal to  $m - 1$ , assuming that  $M = m$ . Then

$$P_n(M = m|B_i) = P_{n-i}(M = m - 1).$$

Therefore for  $n \geq 4$

$$\begin{aligned} P_n(M = m) &= \sum_{i=2}^{n-2} P_n(m|B_i)P(B_i) \\ &= \sum_{i=2}^{n-2} \frac{1}{4}P_i(1)P_{n-i}(m - 1). \end{aligned}$$

This and (\*) imply (see Lemma 1) that

$$P_n(m) = 2^{2-n} \binom{n - 1}{2m - 1}$$

for  $1 \leq m \leq \lfloor \frac{n}{2} \rfloor$ .

Now we can calculate the expected value of number of societies formed by  $n$  points. For  $n \geq 3$  we have

$$\begin{aligned} E_n(M) &= \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} iP_n(i) = \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} 2^{2-n} \binom{n-1}{2i-1} i \\ &= 2^{1-n} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \binom{n-1}{2i-1} 2i = 2^{1-n} (2^{n-2} + (n-1)2^{n-3}) \\ &= \frac{n+1}{4}. \end{aligned}$$

Consequently

$$E_n(M) = \begin{cases} 2 & \text{for } n = 2, \\ \frac{n+1}{4} & \text{for } n \geq 3. \end{cases}$$

The second moment (for  $n \geq 4$ ) is equal to

$$\begin{aligned} E_n(M^2) &= \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} i^2 P_n(i) = \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} 2^{2-n} \binom{n-1}{2i-1} i^2 \\ &= 2^{-n} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \binom{n-1}{2i-1} (2i)^2 \\ &= 2^{-n} (n2^{n-2} + (n-1)2^{n-3} + (n-1)(n-2)2^{n-4}) \\ &= \frac{n(n+3)}{16}. \end{aligned}$$

And thus we obtain

$$\text{Var}_n(M) = \begin{cases} 0 & \text{for } n = 2, 3, \\ \frac{n-1}{16} & \text{for } n \geq 4. \end{cases} \quad \blacksquare$$

Although we formulated the problem for  $R^d$ , we provided results only for the one-dimensional case. Even for simpler model of nearest neighbour graph (see [11] and [3]), higher-dimensional situations become complex enough to require simulation.

## References

- [1] P. Avis and J. Horton, *Remarks on the sphere of influence graph*, in: ed. J.E. Goodman, *et al.* Discrete Geometry and Convexity (New York Academy of Science, New York) 323–327.
- [2] T. Chalker, A. Godbole, P. Hitczenko, J. Radcliff and O. Ruehr, *On the size of a random sphere of influence graph*, Adv. in Appl. Probab. **31** (1999) 596–609.
- [3] E.G. Enns, P.F. Ehlers and T. Misi, *A cluster problem as defined by nearest neighbours*, The Canadian Journal of Statistics **27** (1999) 843–851.
- [4] Z. Füredi, *The expected size of a random sphere of influence graph*, Intuitive Geometry, Bolyai Math. Soc. **6** (1995) 319–326.
- [5] Z. Füredi and P.A. Loeb, *On the best constant on the Besicovitch covering theorem*, in: Proc. Coll. Math. Soc. J. Bolyai **63** (1994) 1063–1073.
- [6] P. Hitczenko, S. Janson and J.E. Yukich, *On the variance of the random sphere of influence graph*, Random Struct. Alg. **14** (1999) 139–152.
- [7] L. Guibas, J. Pach and M. Sharir, *Sphere of influence graphs in higher dimensions*, in: Proc. Coll. Math. Soc. J. Bolyai **63** (1994) 131–137.
- [8] T.S. Michael and T. Quint, *Sphere of influence graphs: a survey*, Congr. Numer. **105** (1994) 153–160.
- [9] T.S. Michael and T. Quint, *Sphere of influence graphs and the  $L_\infty$ -metric*, Discrete Appl. Math. **127** (2003) 447–460.
- [10] Toussaint, *Pattern recognition of geometric complexity*, in: Proceedings of the 5th Int. Conference on Pattern Recognition, (1980) 1324–1347.
- [11] D. Warren and E. Seneta, *Peaks and eulerian numbers in a random sequence*, J. Appl. Prob. **33** (1996) 101–114.

Received 9 September 2004

Revised 4 May 2005