

NOTE

**A RESULT RELATED TO THE  
LARGEST EIGENVALUE OF A TREE**

GURUSAMY RENGASAMY VIJAYAKUMAR

*School of Mathematics*  
*Tata Institute of Fundamental Research*  
*Homi Bhabha Road, Colaba, Mumbai 400 005, India*  
**e-mail:** vijay@math.tifr.res.in

**Abstract**

In this note we prove that  $\{0, 1, \sqrt{2}, \sqrt{3}, 2\}$  is the set of all real numbers  $\ell$  such that the following holds: every tree having an eigenvalue which is larger than  $\ell$  has a subtree whose largest eigenvalue is  $\ell$ .

**Keywords:** eigenvalues of a graph, characteristic polynomial.

**2000 Mathematics Subject Classification:** 05C50, 15A18.

For terminology and notation, we follow [8]. The path with  $n$  vertices and the star with  $n$  edges are denoted by  $P_n$  and  $K_{1,n}$ , respectively. The largest eigenvalue and the least one of a graph  $G$  are denoted by  $\Lambda(G)$  and  $\lambda(G)$ , respectively. Let  $A$  be the adjacency matrix of  $G$ . Then  $|xI - A|$ , the characteristic polynomial of  $G$ , is denoted by  $\phi(G; x)$ . In [1], it has been found that  $\{-2, -\sqrt{2}, -1, 0\}$  is the set of all real numbers  $\ell$  such that if the least eigenvalue of a graph is less than  $\ell$ , then the least eigenvalue of one of its induced subgraphs is equal to  $\ell$ . A result similar to this one is proved in this note: we determine  $\mathcal{L}$  which is defined to be the set of all real numbers  $\ell$  such that the following holds: if  $T$  is a tree with  $\Lambda(T) > \ell$ , then for some subtree  $U$  of  $T$ ,  $\Lambda(U) = \ell$ . To prove our result, we need the following facts:

- (1) If  $F$  is a forest and  $u$  is a vertex of  $F$ , then

$$\phi(F; x) = x\phi(F-u; x) - \sum_{v \in N(u)} \phi(F-u-v; x). \quad (\text{See [8, Page 468].})$$

- (2)  $\Lambda(P_5) = \sqrt{3}$ . (This fact can be easily derived by using the above formula; for more information in this connection, see [5] and [4, Problems 1.29 and 11.5].)
- (3) For each  $n \in \mathbb{N}$ ,  $\Lambda(K_{1,n}) = \sqrt{n}$ . (By using (1), it can be easily verified that  $\phi(K_{1,n}; x) = x^{n-1}(x^2 - n)$ ; see [8, Pages 453–454] for an alternative method.)
- (4) If  $H$  is a proper subgraph of a connected graph  $G$ , then  $\Lambda(H) < \Lambda(G)$ . (See [2, Page 178].)

Obviously  $0 \in \mathcal{L}$ . Let  $T$  be any tree. If  $\Lambda(T) > 1$ , then  $K_2$  is a subtree of  $T$ . Therefore  $1 \in \mathcal{L}$ . If  $\Lambda(T) > \sqrt{2}$ , then  $K_{1,2}$  is a subtree of  $T$ . Therefore by (3),  $\sqrt{2} \in \mathcal{L}$ .

Let  $T$  be a tree with  $\Lambda(T) > \sqrt{3}$ . By (2) and (4),  $T$  cannot be a subtree of  $P_4$ . Therefore it contains  $P_5$  or  $K_{1,3}$ ; now (2) and (3) imply that  $T$  has a subtree whose largest eigenvalue is  $\sqrt{3}$ . Therefore  $\sqrt{3} \in \mathcal{L}$ .

In [7], the family of all graphs  $G$  with  $\Lambda(G) = 2$  has been determined. By using this family, the following result can be derived.

- (5) Every graph  $G$  with  $\Lambda(G) > 2$  has a (connected) subgraph  $H$  with  $\Lambda(H) = 2$ .

A shorter method of classifying the above mentioned family has been found in [3]; in its process of classification, (5) has been observed; but it has not been stated explicitly. Note that (5) is an easy consequence of the main result of [6]: every signed graph  $S$  with  $\lambda(S) < -2$  has an induced subgraph  $R$  with  $\lambda(R) = -2$ . Confining (5) to trees we find that  $2 \in \mathcal{L}$ .

Summary of what we have observed so far:


- (6)  $0, 1, \sqrt{2}, \sqrt{3}, 2 \in \mathcal{L}$ .

Now we proceed to show that  $\mathcal{L}$  does not have elements other than those listed above. As a prelude to this end, we have the following observation.

- (7) A real number  $\ell$  does not belong to  $\mathcal{L}$  when  $\ell^2 \notin \mathbb{Z}$ . (Reason: for any integer  $m > \ell^2$ , by (3),  $\Lambda(K_{1,m}) > \ell$  but for each subtree  $U$  of  $K_{1,m}$ ,  $\Lambda(U) \neq \ell$ .)

The main work of this note is concerned with constructing for each  $k \in \mathbb{N}$ , a tree  $T$  such that (i)  $\Lambda(T) > \sqrt{k+4}$  and (ii) for each proper subtree  $U$  of  $T$ ,  $\Lambda(U) < \sqrt{k+4}$ . If  $p, q, r$  are three nonnegative integers, then the tree

$T(p, q, r)$  is formed from  $K_{1,p}$ ,  $K_{1,q}$  and  $r$  copies of  $K_2$ , by joining the vertex of degree  $p$  in  $K_{1,p}$  with the vertex of degree  $q$  in  $K_{1,q}$  and joining the latter with one vertex of each  $K_2$ . Thus, the degree of the center of  $K_{1,q}$  in the new tree is  $q + r + 1$ .



The tree  $T(2, 1, 6)$

In the recursive formula given by (1), taking  $F$  to be  $T(p, q, r)$  and  $u$  to be the vertex of degree  $q + r + 1$  mentioned above, we get

$$\begin{aligned} \phi(T(p, q, r); x) &= xx^{p-1}(x^2 - p)x^q(x^2 - 1)^r - x^p x^q (x^2 - 1)^r \\ &\quad - qx^{p-1}(x^2 - p)x^{q-1}(x^2 - 1)^r - rx^{p-1}(x^2 - p)x^q x(x^2 - 1)^{r-1}. \end{aligned}$$

Simplifying we get

$$\begin{aligned} \phi(T(p, q, r); x) &= x^{p+q-2}(x^2 - 1)^{r-1} [(x^2 - 1)(x^2 - p)(x^2 - q) - (r + 1)x^4 + (pr + 1)x^2]. \end{aligned}$$

**Theorem.** *If  $k$  is an integer which exceeds 1, then  $\sqrt{k + 3} \notin \mathcal{L}$ .*

**Proof.** The characteristic polynomials of the trees  $T(2, 1, k)$ ,  $T(2, 0, k)$ ,  $T(1, 1, k)$  and  $T(2, 2, k - 1)$  given by the above formula can be expressed as follows

$$\begin{aligned} \phi(T(2, 1, k); x) &= x(x^2 - 1)^{k-1} \{(x^2 - k - 3)x^2(x^2 - 2) - 2\}; \\ \phi(T(2, 0, k); x) &= (x^2 - 1)^{k-1} \{(x^2 - k - 3)[x^2(x^2 - 1) + k] + k(k + 3)\}; \\ \phi(T(1, 1, k); x) &= (x^2 - 1)^{k-1} \{(x^2 - k - 3)[x^2(x^2 - 1) + 1] + k + 2\}; \\ \phi(T(2, 2, k - 1); x) &= x^2(x^2 - 1)^{k-2} \{(x^2 - k - 3)(x^2 - 1)^2 + (k - 1)\}. \end{aligned}$$

Since  $\phi(T(2, 1, k); \sqrt{k+3}) < 0$  and  $\phi(T(2, 1, k); \infty) = \infty$ , it follows that the largest root of  $\phi(T(2, 1, k); x)$  exceeds  $\sqrt{k+3}$ ; i.e.,  $\Lambda(T(2, 1, k)) > \sqrt{k+3}$ . Let  $U$  be a proper subtree of  $T(2, 1, k)$ ; note that  $U$  is a subgraph of either  $T(2, 0, k)$  or  $T(1, 1, k)$  or  $T(2, 2, k-1)$ . Since the largest eigenvalue of each of the latter trees is less than  $\sqrt{k+3}$  because this eigenvalue is a root of one of the above polynomials which are positive on the interval  $[\sqrt{k+3}, \infty)$ , by (4) it follows that  $\Lambda(U) < \sqrt{k+3}$ . Therefore  $\sqrt{k+3} \notin \mathcal{L}$ . ■

Now combining (6), (7) and the above theorem, we get our result. Since the spectrum of any tree is symmetric about the origin (see [2, Page 178]), the dual of this result, obtained from its statement in the abstract by replacing the words ‘larger’, ‘largest’, and the numbers  $1, \sqrt{2}, \sqrt{3}, 2$  by ‘less’, ‘least’ and  $-1, -\sqrt{2}, -\sqrt{3}, -2$  respectively also holds; i.e., for a real number  $\ell$ , each tree  $T$  with  $\lambda(T) < \ell$  has a subtree  $U$  with  $\lambda(U) = \ell$  if and only if  $\ell \in \{0, -1, -\sqrt{2}, -\sqrt{3}, -2\}$ .

### Acknowledgement

The author expresses his gratitude to the referee for pointing out some mistakes and for suggesting some modifications.

### REFERENCES

- [1] M. Doob, *A surprising property of the least eigenvalue of a graph*, Linear Algebra and Its Applications **46** (1982) 1–7.
- [2] C. Godsil and G. Royle, *Algebraic Graph Theory* (Springer, New York, 2001).
- [3] P.W.H. Lemmens and J.J. Seidel, *Equiangular lines*, Journal of Algebra **24** (1973) 494–512.
- [4] L. Lovász, *Combinatorial Problems and Exercises* (North-Holland Publishing Company, Amsterdam, 1979).
- [5] A.J. Schwenk, *Computing the characteristic polynomial of a graph*, in: Graphs and Combinatorics, eds. R.A. Bari and F. Harary, Springer-Verlag, Lecture Notes in Math. **406** (1974) 153–172.
- [6] N.M. Singhi and G.R. Vijayakumar, *Signed graphs with least eigenvalue  $< -2$* , European J. Combin. **13** (1992) 219–220.
- [7] J.H. Smith, *Some properties of the spectrum of a graph*, in: Combinatorial Structures and their Applications, eds. R. Guy, H. Hanani, N. Sauer and J. Schönheim, Gordon and Breach, New York (1970), 403–406.

- [8] D.B. West, Introduction to Graph Theory, Second edition (Printice Hall, New Jersey, USA, 2001).

Received 3 October 2007

Revised 10 June 2008

Accepted 10 June 2008