

THE COMPETITION NUMBERS OF JOHNSON GRAPHS

SUH-RYUNG KIM*, BORAM PARK*^{†‡}

Department of Mathematics Education
Seoul National University, Seoul 151-742, Korea

YOSHIO SANOS[§]

Research Institute for Mathematical Sciences
Kyoto University, Kyoto 606-8502, Japan

Abstract

The competition graph of a digraph D is a graph which has the same vertex set as D and has an edge between two distinct vertices x and y if and only if there exists a vertex v in D such that (x, v) and (y, v) are arcs of D . For any graph G , G together with sufficiently many isolated vertices is the competition graph of some acyclic digraph. The competition number $k(G)$ of a graph G is defined to be the smallest number of such isolated vertices. In general, it is hard to compute the competition number $k(G)$ for a graph G and to characterize all graphs with given competition number k has been one of the important research problems in the study of competition graphs.

The Johnson graph $J(n, d)$ has the vertex set $\{v_X \mid X \in \binom{[n]}{d}\}$, where $\binom{[n]}{d}$ denotes the set of all d -subsets of an n -set $[n] = \{1, \dots, n\}$, and two vertices v_{X_1} and v_{X_2} are adjacent if and only if $|X_1 \cap X_2| = d - 1$. In this paper, we study the edge clique number and the competition number of $J(n, d)$. Especially we give the exact competition numbers of $J(n, 2)$ and $J(n, 3)$.

*This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) (KRF-2008-531-C00004).

[†]The author was supported by Seoul Fellowship.

[‡]Corresponding author. *E-mail address*: kawa22@snu.ac.kr

[§]The author was supported by JSPS Research Fellowships for Young Scientists. The author was also supported partly by Global COE program “Fostering Top Leaders in Mathematics”.

Keywords: competition graph, competition number, edge clique cover, Johnson graph.

2010 Mathematics Subject Classification: 05C69, 05C75.

1. INTRODUCTION

The *competition graph* $C(D)$ of a digraph D is a simple undirected graph which has the same vertex set as D and has an edge between two distinct vertices x and y if and only if there is a vertex v in D such that (x, v) and (y, v) are arcs of D . The notion of a competition graph was introduced by Cohen [3] as a means of determining the smallest dimension of ecological phase space (see also [4]). Since then, various variations have been defined and studied by many authors (see [11, 15] for surveys and [1, 2, 7, 8, 9, 10, 12, 14, 19, 20] for some recent results). Besides an application to ecology, the concept of competition graph can be applied to a variety of fields, as summarized in [17].

Roberts [18] observed that, for a graph G , G together with sufficiently many isolated vertices is the competition graph of an acyclic digraph. Then he defined the *competition number* $k(G)$ of a graph G to be the smallest number k such that G together with k isolated vertices is the competition graph of an acyclic digraph.

A subset S of the vertex set of a graph G is called a *clique* of G if the subgraph of G induced by S is a complete graph. For a clique S of a graph G and an edge e of G , we say e is *covered by* S if both of the endpoints of e are contained in S . An *edge clique cover* of a graph G is a family of cliques such that each edge of G is covered by some clique in the family. The *edge clique cover number* $\theta_E(G)$ of a graph G is the minimum size of an edge clique cover of G . We call an edge clique cover of G with the minimum size $\theta_E(G)$ a *minimum edge clique cover* of G . A *vertex clique cover* of a graph G is a family of cliques such that each vertex of G is contained in some clique in the family. The *vertex clique cover number* $\theta_V(G)$ of a graph G is the minimum size of a vertex clique cover of G . Dutton and Brigham [5] characterized the competition graphs of acyclic digraphs using edge clique covers of graphs.

Roberts [18] observed that the characterization of competition graphs is equivalent to the computation of competition numbers. It does not seem to be easy in general to compute $k(G)$ for a graph G , as Opsut [16] showed

that the computation of the competition number of a graph is an NP-hard problem (see [11, 13] for graphs whose competition numbers are known). For some special graph families, we have explicit formulae for computing competition numbers. For example, if G is a chordal graph without isolated vertices then $k(G) = 1$, and if G is a nontrivial triangle-free connected graph then $k(G) = |E(G)| - |V(G)| + 2$ (see [18]).

In this paper, we study the competition numbers of Johnson graphs. We denote an n -set $\{1, \dots, n\}$ by $[n]$ and the set of all d -subsets of an n -set by $\binom{[n]}{d}$. The *Johnson graph* $J(n, d)$ has the vertex set $\{v_X \mid X \in \binom{[n]}{d}\}$, and two vertices v_{X_1} and v_{X_2} are adjacent if and only if $|X_1 \cap X_2| = d - 1$ (for reference, see [6]). For example, the Johnson graph $J(5, 2)$ is given in Figure 1.




Figure 1. The Johnson graph $J(5, 2)$.

As it is known that $J(n, d) \cong J(n, n - d)$, we assume that $n \geq 2d$. Our main results are the following.

Theorem 1. For $n \geq 4$, we have $k(J(n, 2)) = 2$.

Theorem 2. For $n \geq 6$, we have $k(J(n, 3)) = 4$.

We use the following notation and terminology in this paper. For a digraph D , an ordering v_1, v_2, \dots, v_n of the vertices of D is called an *acyclic ordering* of D if $(v_i, v_j) \in A(D)$ implies $i < j$. It is well-known that a digraph D is acyclic if and only if there exists an acyclic ordering of D . For a digraph D and a vertex v of D , the *out-neighborhood* of v in D is the set $\{w \in V(D) \mid (v, w) \in A(D)\}$. A vertex in the out-neighborhood of a vertex v in a digraph D is called a *prey* of v in D . For simplicity, we denote the

out-neighborhood of a vertex v in a digraph D by $P_D(v)$ instead of usual notation $N_D^+(v)$. For a graph G and a vertex v of G , we define the (*open*) neighborhood $N_G(v)$ of v in G to be the set $\{u \in V(G) \mid uv \in E(G)\}$. We sometimes also use $N_G(v)$ to stand for the subgraph induced by its vertices.

2. A LOWER BOUND FOR THE COMPETITION NUMBER OF $J(n, d)$

In this section, we give lower bounds for the competition number of the Johnson graph $J(n, d)$.

Lemma 3. *Let n and d be positive integers with $n \geq 2d$. For any vertex x of the Johnson graph $J(n, d)$, we have $\theta_V(N_{J(n,d)}(x)) = d$.*

Proof. If $d = 1$, then $J(n, d)$ is a complete graph and the lemma is trivially true. Assume that $d \geq 2$. Take any vertex x in $J(n, d)$. Then $x = v_A$ for some $A \in \binom{[n]}{d}$. For any vertex v_A in $J(n, d)$, the set

$$S_i(v_A) := \{v_B \mid B = (A \setminus \{i\}) \cup \{j\} \text{ for some } j \in [n] \setminus A\}$$

forms a clique of $J(n, d)$ for each $i \in A$. To see why, take two distinct vertices v_B and v_C in $S_i(v_A)$. Then $B = (A \setminus \{i\}) \cup \{j\}$ and $C = (A \setminus \{i\}) \cup \{k\}$ for some distinct $j, k \in [n] \setminus A$. Clearly $|B \cap C| = d - 1$, and so v_B and v_C are adjacent in $J(n, d)$.

Take a vertex v_B in $N_{J(n,d)}(v_A)$. Then $B = (A \setminus \{i\}) \cup \{j\}$ for some $i \in A$ and $j \in [n] \setminus A$ and so $v_B \in S_i(v_A)$. Thus $\{S_i(v_A) \mid i \in A\}$ is a vertex clique cover of $N_{J(n,d)}(v_A)$. Thus $\theta_V(N_{J(n,d)}(v_A)) \leq d$. On the other hand,

$$|((A \setminus \{i\}) \cup \{j\}) \cap ((A \setminus \{i'\}) \cup \{j'\})| = d - 2$$

if $i, i' \in A$ and $j, j' \in [n] \setminus A$ satisfy $i \neq i'$ and $j \neq j'$ (such i, i', j, j' exist since $n \geq 2d \geq 4$). This implies that $\theta_V(N_{J(n,d)}(v_A)) \geq d$. Hence $\theta_V(N_{J(n,d)}(v_A)) = d$. ■

Opsut [16] gave a lower bound for the competition number of a graph G as follows:

$$k(G) \geq \min\{\theta_V(N_G(v)) \mid v \in V(G)\}.$$

Together with Lemma 3, we have $k(J(n, d)) \geq d$ for positive integers n and d satisfying $n \geq 2d$. The following theorem gives a better lower bound for $k(J(n, d))$ if $d \geq 2$.

Theorem 4. For $n \geq 2d \geq 4$, we have $k(J(n, d)) \geq 2d - 2$.

Proof. Put $k := k(J(n, d))$. Then there exists an acyclic digraph D such that $C(D) = J(n, d) \cup I_k$, where $I_k = \{z_1, z_2, \dots, z_k\}$ is a set of isolated vertices. Let $x_1, x_2, \dots, x_{\binom{n}{d}}, z_1, z_2, \dots, z_k$ be an acyclic ordering of D . Let $v_1 := x_{\binom{n}{d}}$ and $v_2 := x_{\binom{n}{d}-1}$. By Lemma 3, we have $\theta_V(N_{J(n,d)}(x_i)) = d$ for $i = 1, \dots, \binom{n}{d}$. Thus v_i has at least d distinct prey in D , that is,

$$(2.1) \quad |P_D(v_i)| \geq d.$$

Since $x_1, x_2, \dots, x_{\binom{n}{d}}, z_1, z_2, \dots, z_k$ is an acyclic ordering of D , we have

$$(2.2) \quad P_D(v_1) \cup P_D(v_2) \subset I_k \cup \{v_1\}.$$

Moreover, we may claim the following:

Claim. For any two adjacent vertices v_{X_1} and v_{X_2} of $J(n, d)$, we have $|P_D(v_{X_1}) \setminus P_D(v_{X_2})| \geq d - 1$.

Proof of Claim. Suppose that v_{X_1} and v_{X_2} are adjacent in $J(n, d)$. Then $|X_1 \cap X_2| = d - 1$ and

$$|[n] \setminus (X_1 \cup X_2)| \geq 2d - |X_1| - |X_2| + |X_1 \cap X_2| = d - 1.$$

We take $d - 1$ elements from $[n] \setminus (X_1 \cup X_2)$, say z_1, z_2, \dots, z_{d-1} , and put $X_1 \cap X_2 := \{y_1, y_2, \dots, y_{d-1}\}$.

For each $1 \leq j \leq d - 1$, we put $Z_j := X_1 \cup \{z_j\} \setminus \{y_j\}$. Then $|Z_j| = d$ and so v_{Z_j} is a vertex in $J(n, d)$. Note that $|Z_j \cap X_1| = d - 1$ and $|Z_j \cap X_2| = d - 2$. Thus v_{Z_j} is adjacent to v_{X_1} while it is not adjacent to v_{X_2} . Therefore

$$P_D(v_{X_1}) \cap P_D(v_{Z_j}) \neq \emptyset \quad \text{and} \quad P_D(v_{X_2}) \cap P_D(v_{Z_j}) = \emptyset.$$

This implies

$$(2.3) \quad P_D(v_{X_1}) \setminus P_D(v_{X_2}) \supseteq \bigcup_{j=1}^{d-1} (P_D(v_{X_1}) \cap P_D(v_{Z_j})),$$

and, trivially, for each $j \in \{1, \dots, d - 1\}$,

$$(2.4) \quad |P_D(v_{X_1}) \cap P_D(v_{Z_j})| \geq 1.$$

Note that $|Z_j \cap Z_i| = d - 2$ for $i \neq j$. Therefore v_{Z_i} and v_{Z_j} are not adjacent and so $P_D(v_{Z_i}) \cap P_D(v_{Z_j}) = \emptyset$. Thus, for $i \neq j$,

$$(2.5) \quad (P_D(v_{X_1}) \cap P_D(v_{Z_i})) \cap (P_D(v_{X_1}) \cap P_D(v_{Z_j})) = \emptyset.$$

From (2.3), (2.4), and (2.5), it follows that

$$|P_D(v_{X_1}) \setminus P_D(v_{X_2})| \geq \sum_{j=1}^{d-1} |P_D(v_{X_1}) \cap P_D(v_{Z_j})| \geq d - 1.$$

This completes the proof of the claim. \square

Now suppose that v_1 and v_2 are not adjacent in $J(n, d)$. Then v_1 and v_2 do not have a common prey in D , that is,

$$(2.6) \quad P_D(v_1) \cap P_D(v_2) = \emptyset.$$

By (2.1), (2.2) and (2.6), we have

$$k + 1 \geq |P_D(v_1) \cup P_D(v_2)| = |P_D(v_1)| + |P_D(v_2)| \geq 2d.$$

Hence $k \geq 2d - 1 > 2d - 2$.

Next suppose that v_1 and v_2 are adjacent in $J(n, d)$. Then v_1 and v_2 have at least one common prey in D , that is,

$$(2.7) \quad |P_D(v_1) \cap P_D(v_2)| \geq 1.$$

By the above claim,

$$(2.8) \quad |P_D(v_1) \setminus P_D(v_2)| \geq d - 1 \quad \text{and} \quad |P_D(v_2) \setminus P_D(v_1)| \geq d - 1.$$

Then

$$\begin{aligned} k + 1 &\geq |P_D(v_1) \cup P_D(v_2)| && \text{(by (2.2))} \\ &= |P_D(v_1) \setminus P_D(v_2)| + |P_D(v_2) \setminus P_D(v_1)| + |P_D(v_1) \cap P_D(v_2)| \\ &\geq (d - 1) + (d - 1) + 1 && \text{(by (2.7) and (2.8))} \\ &= 2d - 1. \end{aligned}$$

Hence it holds that $k \geq 2d - 2$. \blacksquare

3. A MINIMUM EDGE CLIQUE COVER OF $J(n, d)$

In this section, we build a minimum edge clique cover of $J(n, d)$.

Given a Johnson graph $J(n, d)$, we define a family \mathcal{F}_d^n of cliques of $J(n, d)$ as follows. For each $Y \in \binom{[n]}{d-1}$, we put

$$S_Y := \{v_X \mid X = Y \cup \{j\} \text{ for } j \in [n] - Y\}.$$

Note that S_Y is a clique of $J(n, d)$ with size $n - d + 1$. We let

$$(3.1) \quad \mathcal{F}_d^n := \{S_Y \mid Y \in \binom{[n]}{d-1}\}.$$

Then it is not difficult to show that \mathcal{F}_d^n is the collection of cliques of maximum size. Moreover the family \mathcal{F}_d^n is an edge clique cover of $J(n, d)$. To see why, take any edge $v_{X_1}v_{X_2}$ of $J(n, d)$. Then $|X_1 \cap X_2| = d - 1$ and both of v_{X_1} and v_{X_2} belong to the clique $S_{X_1 \cap X_2} \in \mathcal{F}_d^n$. Thus \mathcal{F}_d^n is an edge clique cover of $J(n, d)$.

We will show that \mathcal{F}_d^n is a minimum edge clique cover of $J(n, d)$. Prior to that, we present the following theorem. For two distinct cliques S and S' of a graph G , we say S and S' are *edge disjoint* if $|S \cap S'| \leq 1$.

Theorem 5. $\theta_E(J(n, d)) = \binom{n}{d-1}$ and any minimum edge clique cover of $J(n, d)$ consists of edge disjoint maximum cliques.

Proof. Let \mathcal{E} be a minimum edge clique cover for $J(n, d)$, that is, $\theta_E(J(n, d)) = |\mathcal{E}|$. Since \mathcal{F}_d^n is an edge clique cover with $|\mathcal{F}_d^n| = \binom{n}{d-1}$, we have $\theta_E(J(n, d)) \leq \binom{n}{d-1}$.

Now we show that $|\mathcal{E}| \geq \binom{n}{d-1}$. Since the size of a maximum clique is $n - d + 1$, we have $|E(S)| \leq \binom{n-d+1}{2}$ for each $S \in \mathcal{E}$ where $E(S) = \binom{S}{2}$. Therefore,

$$(3.2) \quad |E(J(n, d))| \leq \sum_{S \in \mathcal{E}} |E(S)| \leq \binom{n-d+1}{2} \times |\mathcal{E}|,$$

and the first equality holds if and only if none of two distinct cliques in \mathcal{E} have a common edge, and the second equality holds if and only if any element of \mathcal{E} is a maximum clique in $J(n, d)$.

Since the Johnson graph $J(n, d)$ is a $d(n - d)$ -regular graph and the number of vertices of $J(n, d)$ is $\binom{n}{d}$,

$$(3.3) \quad |E(J(n, d))| = \frac{1}{2}d(n - d) \times \binom{n}{d} = \binom{n - d + 1}{2} \times \binom{n}{d - 1}.$$

From (3.2) and (3.3), it follows that $\binom{n - d + 1}{2} \times \binom{n}{d - 1} \leq \binom{n - d + 1}{2} \times |\mathcal{E}|$. Thus we have $\binom{n}{d - 1} \leq |\mathcal{E}|$. Hence we can conclude that $\theta_E(J(n, d)) = \binom{n}{d - 1}$.

Furthermore, two equalities in (3.2) must hold, and therefore any minimum edge clique cover of $J(n, d)$ consists of edge disjoint maximum cliques. ■

Since $|\mathcal{F}_d^n| = \binom{n}{d - 1}$, the following corollary is an immediate consequence of Theorem 5:

Corollary 6. *The edge clique cover \mathcal{F}_d^n of $J(n, d)$ defined in (3.1) is a minimum edge clique cover of $J(n, d)$.*

4. PROOFS OF THEOREMS 1 AND 2

First, we define an order \prec on the set $\binom{[n]}{d}$ as follows. Take two distinct elements X_1 and X_2 in $\binom{[n]}{d}$. Let $X_1 = \{i_1, i_2, \dots, i_d\}$ and $X_2 = \{j_1, j_2, \dots, j_d\}$ where $i_1 < \dots < i_d$ and $j_1 < \dots < j_d$. Then we define $X_1 \prec X_2$ if there exists $t \in \{1, \dots, d\}$ such that $i_s = j_s$ for $1 \leq s \leq t - 1$ and $i_t < j_t$. It is easy to see that \prec is a total order.

Now we prove Theorem 1.

Proof of Theorem 1. As $k(J(n, 2)) \geq 2$ by Theorem 4, it remains to show $k(J(n, 2)) \leq 2$. We define a digraph D as follows:

$$V(D) = V(J(n, 2)) \cup I_2$$

where $I_2 = \{z_1, z_2\}$, and

$$A(D) = \bigcup_{i=1}^{n-2} \{(x, v_{\{i+1, i+2\}}) \mid x \in S_{\{i\}} \in \mathcal{F}_2^n\} \\ \cup \bigcup_{i=1}^2 \{(x, z_i) \mid x \in S_{\{n-2+i\}} \in \mathcal{F}_2^n\}.$$

Since the vertices of each clique in the edge clique cover \mathcal{F}_2^n has a common prey in D , it holds that $C(D) = J(n, 2) \cup I_2$. Each vertex in $S_{\{i\}}$ is denoted by v_X for some $X \in \binom{[n]}{2}$ which contains i . Then by the definition of \prec , $v_X \prec v_{\{i+1, i+2\}}$ for $i = 1, \dots, n - 2$. Thus, there exists an arc from a vertex x to a vertex y in D if and only if either $x = v_X$ and $y = v_Y$ with $X \prec Y$, or $x = v_X$ and $y = z_i$ with $X \in S_{\{n-1\}} \cup S_{\{n\}}$ and $i \in \{1, 2\}$. Therefore D is acyclic. Thus we have $k(J(n, 2)) \leq 2$ and this completes the proof. ■

Proof of Theorem 2. By Theorem 4, we have $k(J(n, 3)) \geq 4$. It remains to show $k(J(n, 3)) \leq 4$. We define a digraph D as follows:

$$V(D) = V(J(n, 3)) \cup I_4$$

where $I_4 = \{z_1, z_2, z_3, z_4\}$, and

$$\begin{aligned} A(D) = & \bigcup_{i=1}^{n-3} \bigcup_{j=i+1}^{n-2} \{(x, v_{\{i, j+1, j+2\}}) \mid x \in S_{\{i, j\}} \in \mathcal{F}_3^n\} \\ & \cup \bigcup_{i=1}^{n-3} \{(x, v_{\{i+1, i+2, i+3\}}) \mid x \in S_{\{i, n-1\}} \in \mathcal{F}_3^n\} \\ & \cup \bigcup_{i=1}^{n-4} \{(x, v_{\{i+1, i+2, i+4\}}) \mid x \in S_{\{i, n\}} \in \mathcal{F}_3^n\} \\ & \cup \bigcup_{i=1}^3 \{(x, z_i) \mid x \in S_{\{n-4+i, n\}} \in \mathcal{F}_3^n\} \\ & \cup \{(x, z_4) \mid x \in S_{\{n-2, n-1\}} \in \mathcal{F}_3^n\}. \end{aligned}$$

It is easy to check that

$$\begin{aligned} \mathcal{F}_3^n = & \{S_{\{i, j\}} \mid i = 1, \dots, n - 3; j = i + 1, \dots, n - 2\} \\ & \cup \{S_{\{i, n-1\}} \mid i = 1, \dots, n - 3\} \cup \{S_{\{i, n\}} \mid i = 1, \dots, n - 4\} \\ & \cup \{S_{\{n-3, n\}}, S_{\{n-2, n\}}, S_{\{n-1, n\}}\} \cup \{S_{\{n-2, n-1\}}\}. \end{aligned}$$

Thus $C(D) = J(n, 3) \cup I_4$. Moreover, any vertex $x \in S_{\{i, j\}}$ is denoted by v_X for some $X \in \binom{[n]}{3}$ which contains i and j . By the definition of \prec ,

$X \prec \{i, j+1, j+2\}$. In a similar manner, for x in other cliques in \mathcal{F}_3^n , we may show that $(x, y) \in A(D)$ if and only if either $x = v_X$ and $y = v_Y$ with $X \prec Y$, or $x = v_X$ and $y = z_i$ with $X \in S_{\{n-3, n\}} \cup S_{\{n-2, n\}} \cup S_{\{n-1, n\}} \cup S_{\{n-2, n-1\}}$ and $i \in \{1, 2, 3, 4\}$. Thus D is acyclic. Hence $k(J(n, 3)) \leq 4$. ■

5. CONCLUDING REMARKS

In this paper, we gave some lower bounds for the competition numbers of Johnson graphs, and computed the competition numbers of Johnson graphs $J(n, 2)$ and $J(n, 3)$. It would be natural to ask: What is the exact value of the competition number of a Johnson graph $J(n, 4)$ for $n \geq 8$? Eventually, what are the exact values of the competition numbers of the Johnson graphs $J(n, q)$ for $q \geq 5$?

REFERENCES

- [1] H.H. Cho and S.-R. Kim, *The competition number of a graph having exactly one hole*, Discrete Math. **303** (2005) 32–41.
- [2] H.H. Cho, S.-R. Kim and Y. Nam, *On the trees whose 2-step competition numbers are two*, Ars Combin. **77** (2005) 129–142.
- [3] J.E. Cohen, *Interval graphs and food webs: a finding and a problem*, Document 17696-PR, RAND Corporation (Santa Monica, CA, 1968).
- [4] J.E. Cohen, *Food webs and Niche space* (Princeton University Press, Princeton, NJ, 1978).
- [5] R.D. Dutton and R.C. Brigham, *A characterization of competition graphs*, Discrete Appl. Math. **6** (1983) 315–317.
- [6] C. Godsil and G. Royle, *Algebraic Graph Theory*, Graduate Texts in Mathematics **207** (Springer-Verlag, 2001).
- [7] S.G. Hartke, *The elimination procedure for the phylogeny number*, Ars Combin. **75** (2005) 297–311.
- [8] S.G. Hartke, *The elimination procedure for the competition number is not optimal*, Discrete Appl. Math. **154** (2006) 1633–1639.
- [9] G.T. Helleloid, *Connected triangle-free m -step competition graphs*, Discrete Appl. Math. **145** (2005) 376–383.
- [10] W. Ho, *The m -step, same-step, and any-step competition graphs*, Discrete Appl. Math. **152** (2005) 159–175.

- [11] S.-R. Kim, *The competition number and its variants*, in: Quo Vadis, Graph Theory, (J. Gimbel, J.W. Kennedy, and L.V. Quintas, eds.), Annals of Discrete Mathematics **55** (North-Holland, Amsterdam, 1993) 313–326.
- [12] S.-R. Kim, *Graphs with one hole and competition number one*, J. Korean Math. Soc. **42** (2005) 1251–1264.
- [13] S.-R. Kim and F.S. Roberts, *Competition numbers of graphs with a small number of triangles*, Discrete Appl. Math. **78** (1997) 153–162.
- [14] S.-R. Kim and Y. Sano, *The competition numbers of complete tripartite graphs*, Discrete Appl. Math. **156** (2008) 3522–3524.
- [15] J.R. Lundgren, *Food Webs, Competition Graphs, Competition-Common Enemy Graphs, and Niche Graphs*, in: Applications of Combinatorics and Graph Theory to the Biological and Social Sciences, IMH Volumes in Mathematics and Its Application **17** (Springer-Verlag, New York, 1989) 221–243.
- [16] R.J. Opsut, *On the computation of the competition number of a graph*, SIAM J. Algebraic Discrete Methods **3** (1982) 420–428.
- [17] A. Raychaudhuri and F.S. Roberts, *Generalized competition graphs and their applications*, Methods of Operations Research, **49** (Anton Hain, Königstein, West Germany, 1985) 295–311.
- [18] F.S. Roberts, *Food webs, competition graphs, and the boxicity of ecological phase space*, in: Theory and applications of graphs (Proc. Internat. Conf., Western Mich. Univ., Kalamazoo, Mich., 1976) (1978) 477–490.
- [19] F.S. Roberts and L. Sheng, *Phylogeny numbers for graphs with two triangles*, Discrete Appl. Math. **103** (2000) 191–207.
- [20] M. Sonntag and H.-M. Teichert, *Competition hypergraphs*, Discrete Appl. Math. **143** (2004) 324–329.

Received 14 April 2009

Revised 9 October 2009

Accepted 10 October 2009