

## THE WIENER NUMBER OF POWERS OF THE MYCIELSKIAN

RANGASWAMI BALAKRISHNAN AND S. FRANCIS RAJ

*Srinivasa Ramanujan Centre*  
*SASTRA University*  
*Kumbakonam–612 001, India*

**e-mail:** mathbala@satyam.net.in

**e-mail:** francisraj\_s@yahoo.com

### Abstract

The Wiener number of a graph  $G$  is defined as  $\frac{1}{2} \sum_{u,v \in V(G)} d(u,v)$ ,  $d$  the distance function on  $G$ . The Wiener number has important applications in chemistry. We determine a formula for the Wiener number of an important graph family, namely, the Mycielskians  $\mu(G)$  of graphs  $G$ . Using this, we show that for  $k \geq 1$ ,  $W(\mu(S_n^k)) \leq W(\mu(T_n^k)) \leq W(\mu(P_n^k))$ , where  $S_n$ ,  $T_n$  and  $P_n$  denote a star, a general tree and a path on  $n$  vertices respectively. We also obtain Nordhaus-Gaddum type inequality for the Wiener number of  $\mu(G^k)$ .

**Keywords:** Wiener number, Mycielskian, powers of a graph.

**2010 Mathematics Subject Classification:** 05C12.

### 1. INTRODUCTION

Let  $G$  be a simple connected undirected graph with vertex set  $V(G)$  and edge set  $E(G)$ . Then  $G$  is of order  $|V(G)|$  and size  $|E(G)|$ . Given two distinct vertices  $u, v$  of  $G$ , let  $d(u, v)$  denote the distance between  $u$  and  $v$  (= number of edges in a shortest path between  $u$  and  $v$  in  $G$ ). The Wiener number (also called Wiener index)  $W(G)$  of the graph  $G$  is defined by

$$W(G) = \frac{1}{2} \sum_{a,b \in V(G)} d(a, b) = \sum_{i=1}^D ip(i, G),$$

where  $p(i, G)$  denotes the number of pairs of vertices which are at distance  $i$  in  $G$ , and  $D$  is the diameter of  $G$ . The Wiener number is one of the oldest molecular-graph based structure-descriptors, first proposed by the American chemist Harold Wiener [13] as an aid to determine the boiling point of paraffins. Some of the recent articles in this topic are ([1, 2, 3, 4, 5, 7] and [14]).

In a search for triangle-free graphs with arbitrarily large chromatic numbers, Mycielski [11] developed an interesting graph transformation as follows. For a graph  $G = (V, E)$ , the Mycielskian of  $G$  is the graph  $\mu(G)$  with vertex set  $V \cup V' \cup \{u\}$ , where  $V' = \{x' : x \in V\}$  and is disjoint from  $V$ , and edge set  $E \cup \{xy' : xy \in E\} \cup \{y'u : y' \in V'\}$ . The vertex  $x'$  is called the twin of the vertex  $x$  (and  $x$  the twin of  $x'$ ) and the vertex  $u$  is the root of  $\mu(G)$ . In recent times, there has been an increasing interest in the study of Mycielskians, especially, in the study of their circular chromatic numbers (see, for instance, [9, 6, 8] and [10]).

Let  $H$  be a spanning connected subgraph of a (connected) graph  $G$ . Then for any pair of vertices  $u, v$  of  $G$ ,  $d_G(u, v) \leq d_H(u, v)$ . The  $k$ -th power of a graph  $G$ , denoted by  $G^k$ , is the graph with the same vertex set as  $G$  and in which two vertices are adjacent if and only if their distance in  $G$  is at most  $k$ . Clearly,  $G^1 = G$ .

The complement  $\overline{G}$  of a graph  $G$  is the graph with the same vertex set as  $G$  and in which two vertices  $u, v$  are adjacent if and only if  $u, v$  are non-adjacent in  $G$ . In 1956, Nordhaus and Gaddum [12] gave bounds for the sum of the chromatic number  $\chi(G)$  of a graph  $G$  and its complement  $\overline{G}$  as follows,

**Theorem 1.1.** *For a graph  $G$  of order  $n$ ,  $2\sqrt{n} \leq \chi(G) + \chi(\overline{G}) \leq n + 1$ .*

Zhang and Wu [15] presented the corresponding Nordhaus-Gaddum (in short NG) type inequality for the Wiener number as:

**Theorem 1.2.** *Let  $G$  be a connected graph of order  $n \geq 5$  with connected complement  $\overline{G}$ . Then  $3\binom{n}{2} \leq W(G) + W(\overline{G}) \leq \frac{n^3+3n^2+2n-6}{6}$ .*

The bounds in Theorem 1.2 are sharp.

## 2. WIENER NUMBER OF THE MYCIELSKIAN OF A GRAPH

We start this section by obtaining a formula for the Wiener number of the Mycielskian of a graph.

**Theorem 2.1.** *The Wiener number of the Mycielskian of a connected graph  $G$  of order  $n$  and size  $m$  is given by  $W(\mu(G)) = 6n^2 - n - 7m - 4p(2, G) - p(3, G)$ .*

**Proof.** By definition,

$$W(\mu(G)) = \frac{1}{2} \sum_{a, b \in V(\mu(G))} d(a, b).$$

$$\begin{aligned} \text{Hence } W(\mu(G)) &= \sum_{\substack{a=u, \\ b' \in V'}} d(a, b') + \sum_{\substack{a=u, \\ b \in V}} d(a, b) + \frac{1}{2} \sum_{a', b' \in V'} d(a', b') \\ &\quad + \frac{1}{2} \sum_{a, b \in V} d(a, b) + \sum_{\substack{a \in V, \\ b' \in V'}} d(a, b') \\ &= \sum_1 + \sum_2 + \sum_3 + \sum_4 + \sum_5 \text{ (say)}. \end{aligned}$$

One can observe that,  $\sum_1 = n$ ,  $\sum_2 = 2n$ ,  $\sum_3 = 2\binom{n}{2}$ . As distance between any pair of vertices in  $V$  is at most 4 in  $\mu(G)$ ,  $\sum_4 = \sum_{i=1}^3 ip(i, G) + 4[\binom{n}{2} - \sum_{i=1}^3 p(i, G)]$ . Now the maximum distance from any vertex in  $V$  to any vertex in  $V'$  is 3. Note that if  $ab \in E$ , then  $ab', ba' \in E(\mu(G))$ , that is, each edge of  $G$  will contribute two edges between  $V$  and  $V'$ . Also for every  $a \in V$ ,  $d(a, a') = 2$ , and for every  $a, b \in V$  such that  $d(a, b) = 2$ , we have  $d(a, b') = d(b, a') = 2$ . Thus  $\sum_5 = 2n + 2\sum_{i=1}^2 ip(i, G) + 3[n^2 - n - 2\sum_{i=1}^2 p(i, G)]$  and therefore,  $W(\mu(G)) = 6n^2 - n - 7m - 4p(2, G) - p(3, G)$ . ■

This formula comes in handy when finding the Wiener number of  $\mu(G)$  for which  $p(2, G)$  and  $p(3, G)$  are known even if the diameter of  $G$  is very large.

In [1], X. An et al. have shown that  $W(S_n^k) \leq W(T_n^k) \leq W(P_n^k)$ ,  $k \geq 1$  where  $S_n$ ,  $P_n$  and  $T_n$  denotes a star, a path and a tree other than a star and a path on  $n$  vertices. The formula mentioned in Theorem 2.1 helps us in proving that  $W(\mu(S_n^k)) \leq W(\mu(T_n^k)) \leq W(\mu(P_n^k))$  for any  $k \geq 1$ . However, this cannot be deduced from X. An's result mentioned above. In fact, there are graphs  $G$  and  $H$  with same order and size such that  $W(G) > W(H)$  and  $W(\mu(G)) < W(\mu(H))$ . For example, let  $G$  be  $C_6$  with a pendant edge attached at a pair of opposite vertices and  $H$  be  $C_7$  with a

single pendant edge, then  $W(G) = 62$  and  $W(H) = 61$  while  $W(\mu(G)) = 273$  and  $W(\mu(H)) = 275$ .

**Theorem 2.2.**  $W(\mu(S_n^k)) \leq W(\mu(T_n^k)) \leq W(\mu(P_n^k))$ ,  $k \geq 1$ .

**Proof.** By virtue of Theorem 2.1, the result in Theorem 2.2 is equivalent to  $A = 7p(1, S_n^k) + 4p(2, S_n^k) + p(3, S_n^k) \geq B = 7p(1, T_n^k) + 4p(2, T_n^k) + p(3, T_n^k) \geq C = 7p(1, P_n^k) + 4p(2, P_n^k) + p(3, P_n^k)$ .

We first prove that  $A \geq B$ . If  $k \geq 2$ , then  $S_n^k = K_n$  which implies that  $p(1, S_n^k) = \binom{n}{2} \geq \sum_{i=1}^3 p(i, T_n^k)$  and this inequality implies  $A \geq B$  (as  $7 > 4 > 1$ ). If  $k = 1$ , then  $\text{diam}(S_n) = 2$  and  $D = \text{diam}(T_n) \geq 2$ . This gives,  $p(2, S_n) = \sum_{i=2}^D p(i, T_n)$ , and therefore  $7p(1, S_n) + 4p(2, S_n) \geq 7p(1, T_n) + 4p(2, T_n) + p(3, T_n)$ . Once again,  $A \geq B$ .

Next we prove that  $B \geq C$  by induction on  $n$ .  $B \geq C$  is obvious for  $n \leq 4$ . Let  $T_n$  be a tree of order  $n \geq 5$  and let  $P_n = vv_1 \cdots v_{n-1}$  be a path of order  $n$ . Let  $P = uu_1 \cdots u_d$  be a longest path of  $T_n$  ( $d < n - 1$ ).  $u$  is then a pendant vertex of  $T_n$  and  $T_n - \{u\}$  is a tree of order  $n - 1$ . By induction hypothesis,  $B \geq C$  for  $T_n - \{u\}$  and  $P_n - \{v\}$ . Let  $p(a, i, G)$  denote the number of vertices in  $G$  that are at distance  $i$  from  $a$ . Clearly,  $p(i, T_n^k) = p(i, T_n^k - \{u\}) + p(u, i, T_n^k)$ . So it is enough to prove that  $7p(u, 1, T_n^k) + 4p(u, 2, T_n^k) + p(u, 3, T_n^k) \geq 7p(v, 1, P_n^k) + 4p(v, 2, P_n^k) + p(v, 3, P_n^k)$ .

We know that  $p(v, i, P_n^k) \leq k$  for each  $i = 1$  to  $D = \text{diam}(P_n^k)$ . If there are  $k$  vertices of  $P^k$  in  $T_n^k$  adjacent to  $u$ , then  $p(u, 1, T_n^k) \geq p(v, 1, P_n^k)$ . If not,  $u$  will be a universal vertex of  $T_n^k$  (that is, a vertex adjacent to all the other vertices of  $T_n^k$ ). Thus in any case,  $p(u, 1, T_n^k) \geq p(v, 1, P_n^k)$ .

If  $p(u, 2, T_n^k) < p(v, 2, P_n^k) \leq k$ , then  $\text{diam}(T_n^k) \leq 2$  (This is because if  $\text{diam}(T_n^k) > 2$ , then along the longest path in  $T_n^k$ , there will be  $k$  vertices which would be at distance 2 from  $u$  which is a contradiction). This gives  $p(u, 1, T_n^k) + p(u, 2, T_n^k) = (n - 1) \geq p(v, 1, P_n^k) + p(v, 2, P_n^k) + p(v, 3, P_n^k)$ , and as  $7 > 4 > 1$ ,  $7p(u, 1, T_n^k) + 4p(u, 2, T_n^k) \geq 7p(v, 1, P_n^k) + 4p(v, 2, P_n^k) + p(v, 3, P_n^k)$ .

Next if,  $p(u, 2, T_n^k) \geq p(v, 2, P_n^k)$  and  $p(u, 3, T_n^k) \geq p(v, 3, P_n^k)$  then clearly,  $B \geq C$ . Otherwise,  $\text{diam}(T_n^k) \leq 3$ , (Same argument as above) which shows that  $p(u, 1, T_n^k) + p(u, 2, T_n^k) + p(u, 3, T_n^k) = (n - 1) \geq p(v, 1, P_n^k) + p(v, 2, P_n^k) + p(v, 3, P_n^k)$  and hence  $7p(u, 1, T_n^k) + 4p(u, 2, T_n^k) + p(u, 3, T_n^k) \geq 7p(v, 1, P_n^k) + 4p(v, 2, P_n^k) + p(v, 3, P_n^k)$ . ■

It can easily be seen from the proof of Theorem 2.2 that when  $k = 1$ , we have strict inequality for  $n \geq 5$ .

**Corollary 2.3.** *If  $G$  is a connected graph of order  $n$ , then  $W(\mu(G^k)) \leq W(\mu(P_n^k))$ .*

**Proof.** Let  $T$  be a spanning tree of  $G$ . In view of Theorem 2.2, it suffices to prove that  $W(\mu(G^k)) \leq W(\mu(T^k))$ . Any pair of vertices of  $T^k$  at distance  $i$  will be at distance at most  $i$  in  $G^k$ . Therefore,  $7p(1, G^k) + 4p(2, G^k) + p(3, G^k) \geq 7p(1, T^k) + 4p(2, T^k) + p(3, T^k)$ . Thus  $W(\mu(G^k)) \leq W(\mu(P_n^k))$ . ■

### 3. NG TYPE RESULTS FOR THE WIENER NUMBER OF MYCIELSKI GRAPHS AND THEIR POWERS

When  $G$  (of order  $n$  and size  $m$ ) has no isolated vertices,  $\mu(G)$  is connected while  $\overline{\mu(G)}$  is connected always. It is easy to see that the diameter of  $\overline{\mu(G)}$  is 2 and one can establish that  $W(\overline{\mu(G)}) = 2n^2 + 2n + 3m$ .

This shows that  $W(\mu(G)) + W(\overline{\mu(G)}) = 8n^2 + n - 4m - 4p(2, G) - p(3, G)$ .

As in the proof of Theorem 2.2, one can prove the following.

**Theorem 3.1.**  $W(\mu(S_n^k)) + W(\overline{\mu(S_n^k)}) \leq W(\mu(T_n^k)) + W(\overline{\mu(T_n^k)}) \leq W(\mu(P_n^k)) + W(\overline{\mu(P_n^k)})$  for any  $k \geq 1$ .

Now  $W(\mu(G)) + W(\overline{\mu(G)})$  is maximum, when  $4m + 4p(2, G) + p(3, G)$  is least. As  $W(P_n^k) = \sum_{i=1}^{n-1} \lceil \frac{i}{k} \rceil (n-i)$  (see [1]),  $p(i, P_n^k) = \sum_{j=1}^k \{n - (k(i-1) + j)\}$  for  $i < D$ , the diameter of  $P_n^k$  and thus we see that  $4m + 4p(2, P_n^k) + p(3, P_n^k)$  is least when  $k = 1$ . From the proof of Corollary 2.3,  $W(\mu(G^k)) + W(\overline{\mu(G^k)}) \leq W(\mu(T^k)) + W(\overline{\mu(T^k)})$  where  $T$  is a spanning tree of  $G$ . Hence, for  $n \geq 3$ , we have  $W(\mu(G^k)) + W(\overline{\mu(G^k)}) \leq W(\mu(P_n^k)) + W(\overline{\mu(P_n^k)}) \leq W(\mu(P_n)) + W(\overline{\mu(P_n)}) = 8n^2 - 8n + 15$ .  $W(\mu(G)) + W(\overline{\mu(G)})$  is minimum for graphs with diameter at most two and for these graphs  $W(\mu(G)) + W(\overline{\mu(G)}) = 8n^2 + n - 4\binom{n}{2} = 6n^2 + 3n$ , and therefore,  $6n^2 + 3n \leq W(\mu(G^k)) + W(\overline{\mu(G^k)}) \leq 8n^2 - 8n + 15$ . Zhang and Wu [15] presented the NG type inequality for the Wiener number as given in Theorem 1.2. In our case, for Mycielski graphs  $|V(\mu(G))| = 2n + 1$ . Thus the corresponding inequality of Zhang and Wu [15] for graphs of order  $2n + 1$  is given by  $6n^2 + 3n \leq W(G) + W(\overline{G}) \leq \frac{8n^3 + 24n^2 + 22n}{6}$ . We can easily see that our bound for  $W(\mu(G^k)) + W(\overline{\mu(G^k)})$  is better than the bound of Zhang and Wu for  $\mu(G^k)$  as  $\frac{8n^3 + 24n^2 + 22n}{6} - (8n^2 - 8n + 15) > 0$ ,  $n \geq 3$ .

In a similar way, we might be tempted to obtain the NG type inequalities for the following sums:

- (i)  $W(\mu(G)^k) + W(\overline{\mu(G)^k})$ ,
- (ii)  $W(\mu(G)^k) + W(\overline{\mu(G)^k})$ ,
- (iii)  $W(\mu(G^k)) + W(\mu(\overline{G^k}))$ ,
- (iv)  $W(\mu(G^k)) + W(\mu(\overline{G^k}))$ .

Of these four, (i), (ii) and (iii) are uninteresting as  $\overline{G^k}$  is disconnected in most of the choices for  $G$  while  $\overline{\mu(G)^k}$  ( $k \geq 2$ ) is always disconnected (as  $u$  becomes a universal vertex in  $(\mu(G)^k)$  and diameter of  $\mu(G)$  and  $\overline{\mu(G)}$  are 4 and 2 respectively). Thus NG type inequality seems interesting only for (iv). For this, we need the following lemma due to Zhang and Wu [15].

**Lemma 3.2.** *Let  $G$  be a connected graph with connected complement. Then*

- (1) *if  $\text{diam}(G) > 3$ , then  $\text{diam}(\overline{G}) = 2$ ,*
- (2) *if  $\text{diam}(G) = 3$ , then  $\overline{G}$  has a spanning subgraph which is a double star (see Figure 3.1).*




Figure 3.1

Let  $G$  be a graph of order  $n \geq 5$  with connected complement  $\overline{G}$ . If  $\text{diam}(\overline{G}) = 2$ , we can observe the following.

- (i)  $p(2, \overline{G}) = p(1, G)$ .
- (ii)  $W(\mu(\overline{G})) = 6n^2 - n - 7\left(\binom{n}{2} - p(2, \overline{G})\right) - 4p(2, \overline{G}) = \frac{5}{2}n^2 + \frac{5}{2}n + 3p(1, G)$ .
- (iii)  $W(\mu(G)) + W(\mu(\overline{G})) = \frac{17}{2}n^2 + \frac{3}{2}n - 4p(1, G) - 4p(2, G) - p(3, G)$ . (3.1)

For  $k \geq 2$ ,  $\overline{G^k} = \overline{P_n^k} = K_n$  which implies that  $\mu(\overline{G^k}) = \mu(\overline{P_n^k})$ . Therefore, by virtue of Corollary 2.3, we get that  $W(\mu(G^k)) + W(\mu(\overline{G^k})) \leq W(\mu(P_n^k)) +$

$W(\mu(\overline{P}_n^k))$  for  $k \geq 2$ . The above inequality also holds for  $k = 1$ . This could be seen by arguments similar to those given in the proof of Theorem 2.2 and Corollary 2.3. Thus we have,

**Theorem 3.3.** *Let  $G$  be a connected graph of order  $n \geq 5$  with connected complement  $\overline{G}$ . If  $\text{diam}(\overline{G}) = 2$ , then  $W(\mu(G^k)) + W(\mu(\overline{G}^k)) \leq W(\mu(P_n^k)) + W(\mu(\overline{P}_n^k))$ .*

**Lemma 3.4.** *Let  $G$  be a connected graph of order  $n \geq 5$  with connected complement  $\overline{G}$ . Then  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) \leq W(\mu(P_n^2)) + W(\mu(\overline{P}_n^2))$ .*

**Proof.** As  $\text{diam}(\overline{P}_n) = 2$ , by using Theorem 2.1,

$$\begin{aligned} W(\mu(\overline{P}_n^2)) &= 6n^2 - n - 7p(1, \overline{P}_n^2) \\ &= 6n^2 - n - 7\binom{n}{2} = \frac{5}{2}n^2 + \frac{5}{2}n. \end{aligned}$$

For  $n = 5$ ,  $W(\mu(P_5^2)) = 6.25 - 5 - 7(4 + 3) - 4(2 + 1) = 84$ .

$$\begin{aligned} \text{For } n \geq 6, W(\mu(P_n^2)) &= 6n^2 - n - 7p(1, P_n^2) - 4p(2, P_n^2) - p(3, P_n^2) \\ &= 6n^2 - n - 14n + 21 - 8n + 28 - 2n + 11 \\ &= 6n^2 - 25n + 60. \end{aligned}$$

Hence,  $W(\mu(P_5^2)) + W(\mu(\overline{P}_5^2)) = 159$ , and

$$(3.2) \quad W(\mu(P_n^2)) + W(\mu(\overline{P}_n^2)) = \frac{17}{2}n^2 - \frac{45}{2}n + 60, \text{ for } n \geq 6.$$

By virtue of Theorem 3.3, it is enough to consider the case when,  $\text{diam}(G) = \text{diam}(\overline{G}) = 3$ . For these  $G$  and  $\overline{G}$ ,  $p(1, G) = p(2, \overline{G}) + p(3, \overline{G})$ ,  $p(1, \overline{G}) = p(2, G) + p(3, G)$  and  $p(1, G) + p(1, \overline{G}) = \binom{n}{2}$ . Now by Theorem 2.1,

$$\begin{aligned} W(\mu(G^2)) &= 6n^2 - n - 7p(1, G^2) - 4p(2, G^2) \\ &= 6n^2 - n - 7(p(1, G) + p(2, G)) - 4p(3, G) \\ &= 6n^2 - n - 7p(1, G) - 7(p(1, \overline{G}) - p(3, G)) - 4p(3, G) \\ &= 6n^2 - n - 7\binom{n}{2} + 3p(3, G). \end{aligned}$$

Thus,  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 12n^2 - 2n - 7n^2 + 7n + 3(p(3, G) + p(3, \overline{G}))$ ,

$$(3.3) \quad W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 5n^2 + 5n + 3(p(3, G) + p(3, \overline{G})).$$

As  $diam(G) = diam(\overline{G}) = 3$ , by Lemma 3.2 each of  $G$  and  $\overline{G}$  contains a double star, say,  $S_{a_1, b_1}$  and  $S_{a_2, b_2}$  (see Figure 3.1) as spanning subgraphs of  $G$  and  $\overline{G}$  respectively. Hence  $p(3, G) \leq (a_1 - 1)(b_1 - 1) = a_1b_1 - n + 1$  and  $p(3, \overline{G}) \leq (a_2 - 1)(b_2 - 1) = a_2b_2 - n + 1$ . Also,  $a_i b_i \leq \lfloor \frac{n^2}{4} \rfloor$  for  $i = 1, 2$ . Thus,

$$(3.4) \quad W(\mu(G^2)) + W(\mu(\overline{G}^2)) \leq 5n^2 - n + 6\lfloor \frac{n^2}{4} \rfloor + 6.$$

It can be seen that  $5n^2 - n + 6\lfloor \frac{n^2}{4} \rfloor + 6 < \frac{17}{2}n^2 - \frac{45}{2}n + 60$ , for  $n \geq 7$ . We now consider the remaining cases, namely 5 and 6 separately.

Case (i).  $n = 5$ .

When  $n = 5$ , by equations (3.2) and (3.3),  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 125 + 25 + 3(p(3, G) + p(3, \overline{G})) \leq 162$  and we have already seen that,  $W(\mu(P_5^2)) + W(\mu(\overline{P}_5^2)) = 159$ . We show that  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) \leq 159$ . Suppose  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 160$ , then  $p(3, G) + p(3, \overline{G}) = \frac{10}{3}$ , which is a contradiction. Similarly, we will have a contradiction when  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 161$ . Finally, if  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 162$ ; then,  $p(3, G) + p(3, \overline{G}) = \frac{12}{3} = 4$ . Since  $n = 5$  and  $diam(G) = diam(\overline{G}) = 3$ ,  $p(3, G)$  and  $p(3, \overline{G})$  cannot be greater than 2 and therefore  $p(3, G) = p(3, \overline{G}) = 2$ . There are only two graphs  $G$  of order 5 (see Figure 3.2) with the property that  $n = 5$ ,  $p(3, G) = 2$ . But for these two graphs  $p(3, \overline{G}) = 0$  which is a contradiction.




Fig 3.2



Case (ii).  $n = 6$ .

Here  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) = 210 + 3(p(3, G) + p(3, \overline{G})) \leq 234$  and  $W(\mu(P_5^2)) + W(\mu(\overline{P}_5^2)) = 231$ . Proving  $W(\mu(G^2)) + W(\mu(\overline{G}^2)) \leq 231$  is similar to case(i). In this case the graphs with the required property are as shown in Figure 3.3.




Fig 3.3

■

We now give the result for a general  $k$ .

**Theorem 3.5.** *Let  $G$  be a connected graph of order  $n \geq 5$  with connected complement  $\overline{G}$ . Then for any  $k \geq 1$ ,  $5n^2 + 5n \leq W(\mu(G^k)) + W(\mu(\overline{G}^k)) \leq W(\mu(P_n^k)) + W(\mu(\overline{P}_n^k)) \leq W(\mu(P_n)) + W(\mu(\overline{P}_n)) = \frac{17}{2}n^2 - \frac{15}{2}n + 15$ .*

**Proof.**  $W(\mu(G^k)) + W(\mu(\overline{G}^k))$  is minimum when  $G^k$  and  $\overline{G}^k$  are complete. Thus  $5n^2 + 5n \leq W(\mu(G^k)) + W(\mu(\overline{G}^k))$ . By equation 3.1 and arguments similar to that in Theorem 2.2,  $W(\mu(G)) + W(\mu(\overline{G})) \leq W(\mu(P_n)) + W(\mu(\overline{P}_n))$ . By virtue of Theorem 3.3 and Lemma 3.4, the only case left out for the upper bound to be true is when  $diam(G) = diam(\overline{G}) = 3$  and  $k \geq 3$ . In this case,  $G^k = \overline{G}^k = K_n$  and we see that  $W(\mu(G^k))$  is minimum for  $G^k = K_n$  and therefore  $W(\mu(G^k)) + W(\mu(\overline{G}^k)) \leq W(\mu(P_n^k)) + W(\mu(\overline{P}_n^k)) \leq W(\mu(P_n)) + W(\mu(\overline{P}_n)) = \frac{17}{2}n^2 - \frac{15}{2}n + 15$  (by using equation 3.1). ■

### Acknowledgement

This research was supported by the Department of Science and Technology, Government of India grant DST SR/S4/MS:234/04 dated March 31, 2006.

## REFERENCES

- [1] X. An and B. Wu, *The Wiener index of the  $k$ th power of a graph*, Appl. Math. Lett. **21** (2007) 436–440.
- [2] R. Balakrishnan and S.F. Raj, *The Wiener number of Kneser graphs*, Discuss. Math. Graph Theory **28** (2008) 219–228.
- [3] R. Balakrishnan, N. Sridharan and K.V. Iyer, *Wiener index of graphs with more than one cut vertex*, Appl. Math. Lett. **21** (2008) 922–927.
- [4] R. Balakrishnan, N. Sridharan and K.V. Iyer, *A sharp lower bound for the Wiener Index of a graph*, to appear in Ars Combinatoria.
- [5] R. Balakrishnan, K. Viswanathan and K.T. Raghavendra, *Wiener Index of Two Special Trees*, MATCH Commun. Math. Comput. Chem. **57** (2007) 385–392.
- [6] G.J. Chang, L. Huang and X. Zhu, *Circular Chromatic Number of Mycielski's graphs*, Discrete Math. **205** (1999) 23–37.
- [7] A.A. Dobrynin, I. Gutman, S. Klavžar and P. Žigert, *Wiener Index of Hexagonal Systems*, Acta Appl. Math. **72** (2002) 247–294.
- [8] H. Hajibolhassan and X. Zhu, *The Circular Chromatic Number and Mycielski construction*, J. Graph Theory **44** (2003) 106–115.
- [9] D. Liu, *Circular Chromatic Number for iterated Mycielski graphs*, Discrete Math. **285** (2004) 335–340.
- [10] Liu Hongmei, *Circular Chromatic Number and Mycielski graphs*, Acta Mathematica Scientia **26B** (2006) 314–320.
- [11] J. Mycielski, *Sur le colouriage des graphes*, Colloq. Math. **3** (1955) 161–162.
- [12] E.A. Nordhaus and J.W. Gaddum, *On complementary graphs*, Amer. Math. Monthly **63** (1956) 175–177.
- [13] H. Wiener, *Structural Determination of Paraffin Boiling Points*, J. Amer. Chem. Soc. **69** (1947) 17–20.
- [14] L. Xu and X. Guo, *Catacondensed Hexagonal Systems with Large Wiener Numbers*, MATCH Commun. Math. Comput. Chem. **55** (2006) 137–158.
- [15] L. Zhang and B. Wu, *The Nordhaus-Gaddum-type inequalities for some chemical indices*, MATCH Commun. Math. Comput. Chem. **54** (2005) 189–194.

Received 14 November 2008

Revised 8 October 2009

Accepted 20 October 2009