

## RAINBOW NUMBERS FOR SMALL STARS WITH ONE EDGE ADDED

IZOLDA GORGOL AND EWA ŁAZUKA

*Department of Applied Mathematics*  
*Lublin University of Technology*  
*Nadbystrzycka 38D, 20–618 Lublin, Poland*

**e-mail:** i.gorgol@pollub.pl

**e-mail:** e.lazuka@pollub.pl

### Abstract

A subgraph of an edge-colored graph is *rainbow* if all of its edges have different colors. For a graph  $H$  and a positive integer  $n$ , the *anti-Ramsey number*  $f(n, H)$  is the maximum number of colors in an edge-coloring of  $K_n$  with no rainbow copy of  $H$ . The *rainbow number*  $rb(n, H)$  is the minimum number of colors such that any edge-coloring of  $K_n$  with  $rb(n, H)$  number of colors contains a rainbow copy of  $H$ . Certainly  $rb(n, H) = f(n, H) + 1$ . Anti-Ramsey numbers were introduced by Erdős *et al.* [5] and studied in numerous papers.

We show that  $rb(n, K_{1,4} + e) = n + 2$  in all nontrivial cases.

**Keywords:** rainbow number, anti-Ramsey number.

**2010 Mathematics Subject Classification:** 05C55, 05C35.

### 1. INTRODUCTION

A subgraph of an edge-colored graph is *rainbow* if all of its edges have different colors. For a graph  $H$  and a positive integer  $n$ , the *anti-Ramsey number*  $f(n, H)$  is the maximum number of colors in an edge-coloring of  $K_n$  with no rainbow copy of  $H$ . The *rainbow number*  $rb(n, H)$  is the minimum number of colors such that any edge-coloring of  $K_n$  with  $rb(n, H)$  number of colors contains a rainbow copy of  $H$ . Certainly  $rb(n, H) = f(n, H) + 1$ . Anti-Ramsey numbers were introduced by Erdős *et al.* [5]. They showed that these are closely related to Turán numbers. The *Turán number*  $ex(n, \mathcal{H})$  of

$\mathcal{H}$  is the maximum number of edges of an  $n$ -vertex simple graph having no member of  $\mathcal{H}$  as a subgraph.

For a given graph  $H$  let  $\mathcal{H} = \{H - e : e \in E(H)\}$ . Erdős *et al.* [5] showed  $f(n, H) - ex(n, \mathcal{H}) = o(n^2)$ , as  $n \rightarrow \infty$ . If  $d = \min\{\chi(G) : G \in \mathcal{H}\} \geq 3$ , then by an earlier result of Erdős and Simonovits [4], we have  $ex(n, \mathcal{H}) = \frac{d-2}{d-1} \binom{n}{2} + o(n^2)$ . So the above theorem determines rainbow numbers asymptotically in that case. If  $d \leq 2$ , however, we have  $ex(n, \mathcal{H}) = o(n^2)$  and the above theorem says little about rainbow numbers. Therefore it was suggested by Erdős *et al.* [5] to study Ramsey numbers for graphs that contain an edge whose deletion leaves a bipartite subgraph and put forward two conjectures about paths and cycles.

Simonovits and Sós proved the conjecture for paths determining  $f(n, P_k)$  for  $n$  large enough [15]. As for the conjecture for cycles, they proved it for  $C_3$  by themselves. For  $C_4$  it was proved by Alon [1] and for  $C_5$  and  $C_6$  independently by Jiang and West [9] and by Schiermeyer [13] and completely solved by Montellano-Ballesteros and Neuman-Lara [12].

Moreover rainbow numbers were studied for complete bipartite graphs by Axenovich and Jiang [2], for trees by Jiang and West [10], for subdivided graphs by Jiang [7] and for complete graphs and matchings by Schiermeyer [14]. Recently cycles with an edge added were studied by Montellano-Ballesteros [11] and Gorgol [6].

The aim of the paper is to prove Theorem 4 which says that we need  $n + 2$  colors to be sure that in any coloring of the edges of  $K_n$  with this number of colors we always obtain a rainbow  $K_{1,4} + e$ .

## 2. PRELIMINARIES

Graphs considered below will always be simple. Throughout the paper we use the standard graph theory notation (see, e.g., [3]). In particular,  $G \cup H$ ,  $K_n$  and  $K_{1,r}$  stand, respectively, for disjoint sum of graphs  $G$  and  $H$ , the complete graph on  $n$  vertices and a star with  $r$  rays. A graph  $K_3$  we call a triangle. For a graph  $G$  and its subgraph  $H$  by  $G - H$  we mean a graph obtained from  $G$  by deleting all vertices of  $H$ . For a set  $S$  by  $|S|$  we denote the cardinality of  $S$ .

We will need the following theorems.

**Theorem 1** [5].  $rb(n, K_3) = n$  for  $n \geq 3$ .

**Theorem 2** [8]. *Given positive integers  $n$  and  $r$ , where  $r \leq n - 2$ ,  $rb(n, K_{1,r+1}) = \lfloor \frac{1}{2}n(r - 1) \rfloor + \lfloor \frac{n}{n-r+1} \rfloor + \varepsilon$ , where  $\varepsilon = 1$  or  $2$  if  $n$  is odd,  $r$  is even and  $\lfloor \frac{2n}{n-r+1} \rfloor$  is odd;  $\varepsilon = 1$  otherwise.*

We will also need some lemmas to conduct the inductive proof of Theorem 4. Throughout all proofs we use the following notation.  $C(G)$  is a set of colors used on the edges of a graph  $G$ ;  $C(G, H)$  is a set of colors used on the edges with one endvertex in the vertex-set of a graph  $G$  and the other in the vertex-set of a graph  $H$ ;  $C(v)$  is a set of colors used on the edges incident to a vertex  $v$  and  $c(e)$  denotes the color of the edge  $e$ .

In a graph  $K_{1,4} + e$  a vertex of degree 4 we call a center.

A vertex  $v$  is called *monochromatic* if  $|C(v)| = 1$ .

We call a color  $c$  *private* for a vertex  $v$  if all edges of color  $c$  are incident to  $v$ .

**Claim 1.** If a color  $c$  is private for two vertices  $v$  and  $w$  then an edge  $vw$  is the only edge of color  $c$ .

*Proof.* It follows immediately from the definition of a private vertex. □

**Claim 2.** An arbitrary color can be private for at most two vertices.

*Proof.* It is a straightforward consequence of Claim 1. □

For a fixed coloring of the edges of  $K_n = K$  we construct a bipartite graph  $B$  with bipartition sets  $V$  and  $C$  as follows. Let  $V = V(K)$  and  $C = C(K)$ . We put an edge between  $v \in V$  and  $c \in C$  if and only if  $c$  is private for  $v$ . Note that by Claim 2 each vertex from  $C$  has degree at most 2.

Note that by Claim 1,  $B$  cannot contain  $C_4$ .

**Lemma 1.**  $rb(n, K_{1,4} + e) = n + 2$  for  $n \in \{5, 6\}$ .

*Proof.* The lower bound follows from Theorem 2. So we have to prove the upper bound. We color the edges of  $K_n = K$  with  $n + 2$  colors arbitrarily and show that there exists a rainbow  $K_{1,4} + e$ . By Theorem 2 there exists a rainbow  $K_{1,4} = S$ . The existence of a rainbow  $K_{1,4} + e$  is obvious for  $n \in \{5, 6\}$ . ■

## 3. MAIN RESULTS

Although the next theorem is proved in a more general case in [6], we state it here with a proof to make the paper self-contained.

**Theorem 3.**  $rb(n, K_{1,3} + e) = n$  for  $n \geq 4$ .

**Proof.** By Theorem 1 we have  $rb(n, K_{1,3} + e) \geq n$  for  $n \geq 4$ . So we have to prove the opposite inequality. It is easy to check it for  $n \in \{4, 5, 6\}$ . Therefore let  $n \geq 7$ . We color the edges of  $K_n = K$  with  $n$  colors. By Theorem 1 there exists a rainbow triangle  $T$  with the set of colors  $C(T)$ . If the condition  $C(T, K_n - T) \cap (C(K) - C(T)) \neq \emptyset$  holds then there exists a rainbow  $K_{1,3} + e$ . Otherwise  $|V(K_n - T)| = n - 3$  and  $|C(K_n - T)| \geq n - 3$  and we have a rainbow copy of  $K_{1,3} + e$  in  $K_n - T$  by induction. ■

**Theorem 4.**  $rb(n, K_{1,4} + e) = n + 2$  for  $n \geq 5$ .

**Proof.** The lower bound follows from Theorem 2. So we have to prove the upper bound. The proof will be conducted by induction with respect to  $n$ . For  $n \in \{5, 6\}$  it is Lemma 1.

Therefore let  $n \geq 7$ . We color the edges of  $K_n = K$  with  $n + 2$  colors arbitrarily and construct an appropriate bipartite graph  $B$ .

If there exists a vertex  $v$  such that  $|C(K_n) - C(v)| \geq n + 1$  then  $K_n - v$  is a  $K_{n-1}$  colored with at least  $n + 1$  colors, so a rainbow  $K_{1,4} + e$  exists by induction. A contradiction. Therefore for each vertex  $v \in V(K_n)$  there exist at least two private colors. So each vertex from  $V$  has degree at least 2.

Before the next part of the proof we will show the following two facts.

**Fact 1.** If there is an isolated vertex  $c_0 \in C$  in  $B$  then  $K$  contains a rainbow  $K_{1,4} + e$ .

**Proof of Fact 1.** Assume there is an isolated vertex  $c_0 \in C$  in  $B$ , but there is not any rainbow  $K_{1,4} + e$  in  $K$ .

By Claim 2 each vertex from  $C$  has degree at most 2. By Claim 1 it means that in coloring of  $K$  the respective colors appear exactly once. Hence at least  $n - 1$  vertices from  $C$  have degree exactly 2.

For each  $v \in V$  choose exactly two private colors at  $v$ , and consider a subgraph  $B'$  of  $B$  with  $V(B') = V(B) - \{c_0\}$ , but with an edge between  $v \in V$  and  $c \in C$  if  $c$  is one of the two chosen private colors at  $v$ . Thus

$E(B') \subset E(B)$ ,  $|E(B')| = 2n \leq |E(B)| \leq 2n + 2$  and the maximal degree in  $B'$  is 2.

The above degree conditions determine the structure of  $B'$ . Namely  $B'$  consists of a path with an odd number  $\geq 1$  of vertices, starting and ending in  $C$ , and of zero or more cycle components. Without loss of generality this path could be assumed to be  $c_r v_r c_{r+1} \dots v_n c_{n+1}$ , where  $1 \leq r \leq n + 1$  ( $r = n + 1$  if and only if  $c_{n+1}$  is an isolated vertex in  $B'$ ).

Note that the graph on  $V(K)$  consisting only of the edges colored in the three or two colors  $c_0, c_r, c_{n+1}$  is connected. Thus there is some vertex  $v_i$  with at least two or three colors occurring among its edges: it is easy to see that we may choose  $v_i \notin \{v_r, v_n\}$ , if  $r \leq n$ . Hence, in fact, either  $v_i$  belongs to a 6-cycle  $v_{i-1}c_i v_i c_{i+1} v_{i+1} c'$  or to a path  $v_{i-1}c_{i-1} v_i c_i v_{i+1} c_{i+1} v_{i+2}$  in  $B'$ . In either case,  $|C(v_i)| \geq 4$ . In the case of  $C_6$ ,  $v_i$  would be a center of a rainbow  $K_{1,4} + e$ , where  $e$  would be colored with  $c'$ . Similarly in the case of the path,  $v_i$  would be a center of a rainbow  $K_{1,4} + e$ , where  $e = v_i v_{i+2}$ . Indeed, let  $c(v_i v_{i+2}) = c''$  be the color of the edge  $v_i v_{i+2}$ . It can be  $c_0$  or  $c_{n+1}$  (if  $i + 2 = n$ ) and by the choice of  $v_i$  there is another edge coming out of it of color from  $\{c_0, c_r, c_{n+1}\} \setminus \{c''\}$ . □

**Fact 2.** If  $K$  contains two disjoint rainbow triangles, then  $K$  contains a rainbow  $K_{1,4} + e$ .

**Proof of Fact 2.** Let  $T_1$  and  $T_2$  be these triangles. Note that if there is  $c \in C(T_1) \cap C(T_2)$  then indeed  $c$  is not private at any vertex in  $K$  and thus is isolated in  $B$ , whence then Fact 1 applies. Therefore we have to consider the case when  $T_1 \cup T_2$  is a rainbow  $2K_3$ .

Let  $V(T_1) = \{x, y, z\}$ ,  $V(T_2) = \{a, b, c\}$ ,  $C(T_1) = \{c_1, c_2, c_3\}$  and  $C(T_2) = \{c_4, c_5, c_6\}$ .

If  $|C(T_1) \cup C(T_2) \cup C(T_1, T_2)| \geq 8$  then Fact 2 follows from Lemma 1 so we can assume that  $|C(T_1) \cup C(T_2) \cup C(T_1, T_2)| \leq 7$  which means that there can be at most one color in  $C(T_1, T_2)$  not belonging to  $C(T_1) \cup C(T_2)$ . Let  $K' = K - (T_1 \cup T_2)$ .

Suppose that there exists an edge  $e$  between the triangles  $T_1$  and  $T_2$  of the color not belonging to  $C(T_1) \cup C(T_2)$ . Without loss of generality we can assume that  $e = xa$  and  $c(e) = c$ . Let  $C_R = C(K) - (C(T_1) \cup C(T_2) \cup \{c\})$ . Note that either we have a rainbow  $K_{1,4} + e$  or  $c(xv) \in C(T_1) \cup \{c\}$  for all vertices  $v \in V(K - T_1)$  and  $c(av) \in C(T_2) \cup \{c\}$  for all vertices  $v \in V(K - T_2)$ . If there is at least one edge between  $T_1 \cup T_2$  and  $K'$  of color from  $C_R$ , say  $yw$ , where  $w \in V(K')$ , then we obtain a rainbow  $K_{1,4} + e$ . It is the triangle

$T_1$  with edges  $ya$  and  $yw$ . Note that surely it is the case for  $n = 9$  since  $|C_R| = 4$  and  $|E(K')| = 3$ . If such an edge does not exist it means that  $n \geq 10$ ,  $C(T_1 \cup T_2, K') \subset (C(T_1) \cup C(T_2) \cup \{c\})$  and all colors from  $C_R$  are used on edges of  $K'$ . If there is a rainbow  $K_{1,3} + e$  in  $K'$  then it gives a rainbow  $K_{1,4} + e$  together with one edge coming to  $T_1 \cup T_2$ . Note that obviously it is the case for  $n = 10$  and for  $n \geq 11$  and  $C(K') = C_R$  it follows from Theorem 3. If  $|C(K')| > |C_R|$  for  $n \geq 11$  we obtain a rainbow  $K_{1,4} + e$  by induction.

Therefore we assume that  $C(T_1, T_2) \subset C(T_1) \cup C(T_2)$ . Let  $C_R = C - (C(T_1) \cup C(T_2))$ . If there is at least one edge between  $T_1 \cup T_2$  and  $K'$  of color from  $C_R$ , say  $xw$ , where  $w \in V(K')$ , then all edges coming from  $x$  to  $T_2$  are of colors from  $C(T_1)$  otherwise we get a rainbow  $K_{1,4} + e$ . As a further consequence we get that either there is a rainbow  $K_{1,4} + e$  in  $K$  or all edges coming out from  $T_2$  are of colors from  $C(T_1) \cup C(T_2)$ . In the latter case the graph  $K - T_2$  is colored with at least  $n - 1$  colors so the induction completes the proof. Note that surely it is the case for  $n \in \{7, 8, 9\}$ . So we can assume that  $n \geq 10$ ,  $C(T_1 \cup T_2, K') \subset (C(T_1) \cup C(T_2))$  and all colors from  $C_R$  are used on edges of  $K'$ . Repeating the arguments from the previous part of the proof we prove the fact.  $\square$

Now we are ready to finish the proof of Theorem 4.

By Theorem 1 there exists a rainbow triangle  $T_1$  with the vertex-set  $\{x, y, z\}$  and the set of colors  $C(T_1)$ .

Let  $K' = K - T_1$ . Note that if  $K'$  contains a rainbow triangle then  $K$  contains a rainbow  $K_{1,4} + e$  by Fact 2. Assume then it is not the case. Then  $|C(K')| \leq n - 4$  by Theorem 1.

Let  $C_R = C(K) - C(T_1)$ . Note that if there is a vertex  $v$  in  $T_1$  with  $|C(v) \cap C_R| \geq 2$ , then there is a rainbow  $K_{1,4} + e$  with center  $v$  and containing  $T_1$ . Hence the converse can be assumed.

So we are to consider only the case  $|C(K')| = n - 4$ ,  $|C(v) \cap C_R| = 1$  and the colors  $C(v) \cap C_R$  are distinct for each  $v \in \{x, y, z\}$ .

Then certainly  $|C(T_1, K') \cap C_R| = 3$  and  $C(K') \cap (C(T_1) \cup C(T_1, K')) = \emptyset$ . Now either we have a rainbow  $K_{1,4} + e$  or each edge between  $T_1$  and  $K'$  of the color from  $C_R$  comes out from a different vertex of  $T_1$ . If such an edge of color  $c$  comes out, say from  $x$ , to a vertex  $a$  which is not monochromatic in  $K'$  then  $c(ay) = c(xy)$  and  $c(za) = c(xz)$  or we have a rainbow  $K_{1,4} + e$ . But in this case we also get a rainbow  $K_{1,4} + e$ . It is a rainbow triangle  $ayz$  with two edges coming out from the vertex  $a$ .

It is easy to note that there can be at most one monochromatic vertex in  $K'$ . If there would be at least two such vertices  $a$  and  $b$  then  $C(a) \cap C(K') = C(b) \cap C(K') = \{c(ab)\}$  and so  $K' - \{a, b\}$  would be  $K_{n-5}$  colored with  $n-5$  colors, against the assumption for  $K'$ .

Hence the vertex  $a$  is monochromatic in  $K'$  and all edges between  $T_1$  and  $K'$  of the three colors in  $C(T_1, K') \cap C_R$  have  $a$  as an endpoint. Thus  $|C(a)| = 4$  and there is a rainbow  $K_{1,4} + e$  with the center  $a$  and containing an edge from  $T_1$ . ■

## REFERENCES

- [1] N. Alon, *On the conjecture of Erdős, Simonovits and Sós concerning anti-Ramsey theorems*, J. Graph Theory **7** (1983) 91–94.
- [2] M. Axenovich and T. Jiang, *Anti-Ramsey numbers for small complete bipartite graphs*, Ars Combinatoria **73** (2004) 311–318.
- [3] R. Diestel, *Graph theory* (Springer-Verlag, New York, 1997).
- [4] P. Erdős and M. Simonovits, *A limit theorem in graph theory*, Studia Sci. Math. Hungar. **1** (1966) 51–57.
- [5] P. Erdős, A. Simonovits and V. Sós, *Anti-Ramsey theorems*, in: Infinite and Finite Sets (A. Hajnal, R. Rado, and V. Sós, eds.), Colloq. Math. Soc. J. Bolyai (North-Holland, 1975) 633–643.
- [6] I. Gorgol, *On rainbow numbers for cycles with pendant edges*, Graphs and Combinatorics **24** (2008) 327–331.
- [7] T. Jiang, *Anti-Ramsey numbers for subdivided graphs*, J. Combin. Theory (B) **85** (2002) 361–366.
- [8] T. Jiang, *Edge-colorings with no large polychromatic stars*, Graphs and Combinatorics **18** (2002) 303–308.
- [9] T. Jiang and D.B. West, *On the Erdős-Simonovits-Sós conjecture about the anti-Ramsey number of a cycle*, Combin. Probab. Comput. **12** (2003) 585–598.
- [10] T. Jiang and D.B. West, *Edge-colorings of complete graphs that avoid polychromatic trees*, Discrete Math. **274** (2004) 137–145.
- [11] J.J. Montellano-Ballesteros, *Totally multicolored diamonds*, Electronic Notes in Discrete Math. **30** (2008) 231–236.
- [12] J.J. Montellano-Ballesteros and V. Neuman-Lara, *An anti-Ramsey theorem on cycles*, Graphs and Combinatorics **21** (2005) 343–354.

- [13] I. Schiermeyer, *Rainbow 5- and 6-cycles: a proof of the conjecture of Erdős, Simonovits and Sós*, preprint (TU Bergakademie Freiberg, 2001).
- [14] I. Schiermeyer, *Rainbow numbers for matchings and complete graphs*, *Discrete Math.* **286** (2004) 157–162.
- [15] M. Simonovits and V. Sós, *On restricted colorings of  $K_n$* , *Combinatorica* **4** (1984) 101–110.

Received 29 December 2008

Revised 30 October 2009

Accepted 30 October 2009