

KERNELS BY MONOCHROMATIC PATHS AND THE COLOR-CLASS DIGRAPH

HORTENSIA GALEANA-SÁNCHEZ

Instituto de Matemáticas
Universidad Nacional Autónoma de México
Area de la Investigación Científica
Ciudad Universitaria
04510, México, D.F., México
e-mail: hgaleana@matem.unam.mx

Abstract

An m -colored digraph is a digraph whose arcs are colored with m colors. A directed path is monochromatic when its arcs are colored alike.

A set $S \subseteq V(D)$ is a kernel by monochromatic paths whenever the two following conditions hold:

1. For any $x, y \in S$, $x \neq y$, there is no monochromatic directed path between them.
2. For each $z \in (V(D) - S)$ there exists a zS -monochromatic directed path.

In this paper it is introduced the concept of color-class digraph to prove that if D is an m -colored strongly connected finite digraph such that:

- (i) Every closed directed walk has an even number of color changes,
- (ii) Every directed walk starting and ending with the same color has an even number of color changes, then D has a kernel by monochromatic paths.

This result generalizes a classical result by Sands, Sauer and Woodrow which asserts that any 2-colored digraph has a kernel by monochromatic paths, in case that the digraph D be a strongly connected digraph.

Keywords: kernel, kernel by monochromatic paths, the color-class digraph.

2010 Mathematics Subject Classification: 05C20.

1. INTRODUCTION

For general concepts we refer the reader to [12] and [3]. Let D be a digraph, a set of vertices $S \subseteq V(D)$ is dominating whenever for every $w \in V(D) - S$ there exists a wS -arc in D . (The topic of domination in graphs has been deeply studied by several authors, a very complete study of this topic can be found in [13] and [14]).

Dominating independent sets in digraphs (kernels in digraphs) have found many applications in several topics of Mathematics (see for example [1, 2, 5, 6] and [15]) and they have been studied by several authors, surveys of kernels in digraphs can be found in [4] and [6]. Clearly the concept of kernel by monochromatic paths is a generalization of that of kernel.

The study of the existence of kernels by monochromatic paths in edge-colored digraphs begins with the Theorem of Sands, Sauer and Woodrow proved in [16] which asserts that every 2-colored digraph has a kernel by monochromatic paths. Sufficient conditions for the existence of kernels by monochromatic paths in edge-colored digraphs have been obtained mainly in nearly tournaments and they ask for the monochromaticity or quasi-monochromaticity of small subdigraphs (due to the difficulty of the problem), see for example [8, 9, 10, 11, 7, 17] and [18].

In this paper we give a different approach to obtain sufficient conditions for the existence of a kernel by monochromatic paths in an edge-colored digraph. We introduce the concept of color-class digraph of an m -colored digraph D and study some structural properties of that digraph which imply that D possesses a kernel by monochromatic paths. As a consequence it is obtained a wide generalization of the classical result of Sands, Sauer and Woodrow in the case that the digraph D be strongly connected.

2. THE COLOR-CLASS DIGRAPH OF AN m -COLORED DIGRAPH D

In this section the color-class digraph of an m -colored digraph D is defined; it is proved that some structural properties of this digraph allow us to consider that the m -colored digraph D is essentially 2-colored and we can conclude that D has a kernel by monochromatic paths.

Definition. Let D be an m -colored digraph. The color-class digraph of D denoted $\mathcal{C}_C(D)$ is defined as follows:

$V(\mathcal{C}_C(D)) = \{\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m\}$ where \mathcal{C}_i is the subdigraph of D whose arcs

are the arcs of D colored i and the vertices of \mathcal{C}_i ; are those vertices of D which are initial endpoints or terminal endpoints of the arcs colored i ; \mathcal{C}_i will be called the color-class i of D (Notice that since D is an m -colored digraph, we have $\mathcal{C}_i \neq \emptyset$ for each $1 \leq i \leq m$).

$(\mathcal{C}_i, \mathcal{C}_j) \in A(\mathcal{C}_C(D))$ if and only if there exists two arcs namely $f = (u, v) \in A(D)$ colored i and $g = (v, w) \in A(D)$ colored j .

Observe that $\mathcal{C}_C(D)$ may allow isolated vertices.

Lemma 2.1. *Let D be an m -colored digraph. If D is a strongly connected digraph, then $\mathcal{C}_C(D)$ is a strongly connected digraph.*

Proof. Let $\mathcal{C}_i, \mathcal{C}_j$ be two different vertices of $\mathcal{C}_C(D)$. Since D is an m -colored digraph, there exist $f = (u, v) \in A(\mathcal{C}_i)$ and $g = (z, w) \in A(\mathcal{C}_j)$. If $v = z$ then $(\mathcal{C}_i, \mathcal{C}_j)$ is a $\mathcal{C}_i\mathcal{C}_j$ -directed path in $\mathcal{C}_C(D)$. If $v \neq z$ then we have that there exists a vz -directed path contained in D (because D is an strongly connected digraph). Let $T = (v = u_1, u_1, u_2, \dots, u_{n-1} = z)$ and $P = (u_0 = u, v) \cup T \cup (u_{n-1} = z, u_n = w)$, $P = (u_0 = u, u_1 = v, u_2, u_3, \dots, u_{n-1} = z, u_n = w)$. Take $u_{i_1}, u_{i_2}, \dots, u_{i_k}$ the vertices of P where a color change occurs. So the walk P has k color changes and $(u, T, u_{i_1}) \subseteq \mathcal{C}_i$, $(u_{i_1}, T, u_{i_2}) \subseteq \mathcal{C}_{r_2}$, $(u_{i_2}, T, u_{i_3}) \subseteq \mathcal{C}_{r_3}, \dots, (u_{i_{k-1}}, T, u_{i_k}) \subseteq \mathcal{C}_{r_k}$, $(u_{i_k}, P, w) \subseteq \mathcal{C}_j$ for some $\{r_2, \dots, r_k\} \subseteq \{1, 2, \dots, m\}$. Clearly we have that $\hat{P} = (\mathcal{C}_i, \mathcal{C}_{r_2}, \mathcal{C}_{r_3}, \mathcal{C}_{r_4}, \dots, \mathcal{C}_{r_k}, \mathcal{C}_j)$ is a $\mathcal{C}_i\mathcal{C}_j$ -directed walk in $\mathcal{C}_C(D)$. Therefore there exists a $\mathcal{C}_i\mathcal{C}_j$ -directed path in $\mathcal{C}_C(D)$. ■

Lemma 2.2. *Let D be an m -colored digraph with color classes $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m$ such that neither the pair $(\mathcal{C}_1, \mathcal{C}_2)$ nor $(\mathcal{C}_2, \mathcal{C}_1)$ are arcs of the color-class digraph. And, let \hat{D} the $(m - 1)$ -colored digraph obtained from D by assigning color 1 to each arc of D colored 2 (Thus the arcs of D colored 2 are now colored 1 in \hat{D} , the rest of the arcs of D remain the same). For any $u, v \in V(D) = V(\hat{D})$, $u \neq v$; there exists a uv -monochromatic directed path in D if and only if there exists a uv -monochromatic directed path in \hat{D} .*

Proof. First notice that the digraph \hat{D} is the same as D except that the arcs of \mathcal{C}_2 in D are now colored 1 in \hat{D} ; the color classes of \hat{D} are $\mathcal{C}'_1, \mathcal{C}'_2, \dots, \mathcal{C}'_{m-1}$ where $\mathcal{C}'_1 = \mathcal{C}_1 \cup \mathcal{C}_2$ and $\mathcal{C}'_j = \mathcal{C}_{j+1}$.

First suppose that there exists a uv -monochromatic directed path contained in D and let P be such that a path. Thus $P \subseteq \mathcal{C}_i$ for some $i \in \{1, 2, \dots, m\}$; for $i \in \{3, \dots, m\}$ we have $\mathcal{C}_i = \mathcal{C}'_{i-1}$ and for $i \in \{1, 2\}$ we have $\mathcal{C}_i \subseteq \mathcal{C}'_1$ thus $P \subseteq \mathcal{C}'_j$ for some $j \in \{1, 2, \dots, m - 1\}$ which means that P is a uv -monochromatic directed path in \hat{D} .

Now suppose that T is a uv -monochromatic directed path in \widehat{D} . Thus $T \subseteq \mathcal{C}'_j$ for some $j \in \{1, \dots, m-1\}$. When $j \in \{2, \dots, m-1\}$ we have $\mathcal{C}'_j = \mathcal{C}_{j+1}$ and T is a uv -monochromatic directed path in D . So, suppose $T \subseteq \mathcal{C}'_1 = \mathcal{C}_1 \cup \mathcal{C}_2$; when $T \subseteq \mathcal{C}_1$ or $T \subseteq \mathcal{C}_2$ we have that T is a uv -monochromatic directed path in D . Henceforth we have $T \subseteq \mathcal{C}_1 \cup \mathcal{C}_2$, $T \not\subseteq \mathcal{C}_1$ and $T \not\subseteq \mathcal{C}_2$; assume without loss of generality that T starts in \mathcal{C}_1 . Let $T = (u_0, u_1, \dots, u_n)$ and $g = (u_i, u_{i+1})$ the first arc of T belonging to \mathcal{C}_2 ; hence $f = (u_{i-1}, u_i) \in A(\mathcal{C}_1)$ and it follows from the definition of $\mathcal{C}_C(D)$ that $(\mathcal{C}_1, \mathcal{C}_2) \in A(\mathcal{C}_C(D))$ contradicting our assumption. We conclude that T is a uv -monochromatic directed path in D . ■

Corollary 2.3. *Let D be an m -colored digraph and \widehat{D} the $(m-1)$ -colored digraph obtained from D as in the hypothesis of Lemma 2.2. A set $N \subseteq V(D) = V(\widehat{D})$ is a kernel by monochromatic paths of D if and only if it is a kernel by monochromatic paths of \widehat{D} .*

Theorem 2.4 (Sands, Sauer and Woodrow [16]). *If D is a 1-colored (monochromatic digraph) or D is a 2-colored digraph, then D has a kernel by monochromatic paths.*

This theorem will be useful to prove the next theorem which is the main result of this section

Theorem 2.5. *Let D be an m -colored digraph. If $\mathcal{C}_C(D)$ is a bipartite digraph, then D has a kernel by monochromatic paths.*

Proof. We proceed by induction on $|V(\mathcal{C}_C(D))|$ (i.e., we proceed by induction on m).

For $m = 1$ or $m = 2$ the result follows directly from Theorem 2.4. Suppose that if D' is an $(m-1)$ -colored digraph such that $\mathcal{C}_C(D')$ is bipartite (i.e., that $|V(\mathcal{C}_C(D'))| = m-1$), then D' has a kernel by monochromatic paths, for $m \geq 3$.

Let D be an m -colored digraph, and let V_1, V_2 the bipartition of $V(\mathcal{C}_C(D))$ which witnesses that $\mathcal{C}_C(D)$ is bipartite; so V_1 (resp. V_2) is an independent set in $\mathcal{C}_C(D)$. Since $m \geq 3$, $m = |V(\mathcal{C}_C(D))|$ we have $|V_1| \geq 2$ or $|V_2| \geq 2$; without loss of generality assume that $|V_1| \geq 2$ and let $\mathcal{C}_1, \mathcal{C}_2 \in V_1$. Consider \widehat{D} the $(m-1)$ -colored digraph obtained from D as in the hypothesis of Lemma 2.2. Clearly $\mathcal{C}_C(\widehat{D})$ is the digraph obtained from $\mathcal{C}_C(D)$ by identifying the vertices \mathcal{C}_1 and \mathcal{C}_2 . Since $\mathcal{C}_C(D)$ is bipartite, we have that $\mathcal{C}_C(\widehat{D})$ is also bipartite. Thus it follows from the inductive hypothesis that

\widehat{D} has a kernel by monochromatic paths; let N be such a kernel. Henceforth it follows from Corollary 2.3 that N is a kernel by monochromatic paths of D . ■

3. KERNELS BY MONOCHROMATIC PATHS

In this section we study a condition on D which implies that $\mathcal{C}_C(D)$ is bipartite which from Theorem 2.5 implies that D has a kernel by monochromatic paths.

Theorem 3.1. *Let D be a strongly connected m -colored digraph. If D satisfies the two following conditions:*

- (a) *Every closed directed walk in D possesses an even number of color changes.*
- (b) *Every directed walk starting and ending in arcs of the same color has an even number of color changes.*

Then every directed cycle in $\mathcal{C}_C(D)$ has an even length.

Proof. Assume by contradiction that $\gamma = (\mathcal{C}_0, \mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_{2n}, \mathcal{C}_0)$ is an odd directed cycle in $\mathcal{C}_C(D)$. Where i is the color associated to \mathcal{C}_i . Then, from the definition of $\mathcal{C}_C(D)$ we have that there exists arcs $f_i = (x_i, y_i), f'_i = (x'_i, y'_i)$ both colored i for $i \in \{0, 1, \dots, 2n\}$ such that $y'_i = x_{i+1}, y'_{2n} = x_0$. That means $f'_0 = (x'_0, y'_0)$ is colored 0 and $f_1 = (y'_0 = x_1, y_1)$ is colored 1; $f'_1 = (x'_1, y'_1)$ is colored 1 and $f_2 = (y'_1 = x_2, y_2)$ is colored 2; $f'_2 = (x'_2, y'_2)$ is colored 2 and $f_3 = (y'_2 = x_3, y_3)$ is colored 3; in general $f'_i = (x'_i, y'_i)$ is colored i and $f_{i+1} = (y'_i = x_{i+1}, y_{i+1})$ is colored $i + 1$ and $f'_{2n} = (x'_{2n}, y'_{2n})$ is colored $2n$ and $f_0 = (y'_{2n} = x_0, y_0)$ is colored 0. Since D is a strongly connected digraph; there exists a directed path, namely T_i from y_i to x'_i for each $i \in \{0, 1, \dots, 2n\}$. Thus we have the directed walks $W_i = (x_i, y_i) \cup T_i \cup (x'_i, y'_i)$ starting in f_i and ending in f'_i ; since f_i and f'_i are both colored i we have that W_i has an even number of color changes; for each $i \in \{0, 1, \dots, 2n\}$. Now consider the closed directed walk $W = \bigcup_{i=0}^{2n} W_i$ clearly the color changes of W are those of each W_i and those that occur in x_i for each $i \in \{0, 1, \dots, 2n\}$. Hence the number of color changes of W is odd contradicting our assumption. (See Figure 1.) ■

Theorem 3.2 [3]. *Let D be a strongly connected digraph; D is a bipartite digraph if and only if each directed cycle of D has an even length.*




Figure 1.

Theorem 3.3. *Let D be a strongly connected m -colored digraph. If D satisfies the two following conditions:*

- (a) *Every closed directed walk in D possesses an even number of color changes,*
- (b) *Every directed walk starting and ending in arcs of the same color has an even number of color changes.*

Then D has a kernel by monochromatic paths.

Proof. It follows from Theorem 3.1 that every directed cycle of $\mathcal{C}_C(D)$ has an even length. From Lemma 2.1 $\mathcal{C}_C(D)$ is a strongly connected digraph. Thus from Theorem 3.2 we have that $\mathcal{C}_C(D)$ is a bipartite digraph. Hence we conclude by Theorem 2.5 that D has a kernel by monochromatic paths. ■

As a direct consequence of Theorem 3.3 we have the following two corollaries.

Corollary 3.4. *Let D be a strongly connected m -colored digraph. If D satisfies the two following conditions:*

- (a) *Every closed directed walk is 1-colored or 2-colored;*

- (b) *Every directed walk starting and ending in arcs colored alike is 1-colored or 2-colored.*

Then D has a kernel by monochromatic paths.

Corollary 3.5. *If D is a strongly connected 2-colored digraph, then D has a kernel by monochromatic paths.*

Clearly Theorem 3.3 is a wide generalization of Theorem 2.4 in the case that D is a strongly connected digraph

4. APPLICATIONS

Let D be an m -colored digraph, and let $C = \{c_1, c_2, \dots, c_m\}$ the set of colors used to color $A(D)$.

Denote by $\xi(v) = \{c_i \in C \mid \text{there exists an arc colored } c_i \text{ incident with } v\}$ ($\xi(v)$ are the colors that appear in arcs incident from (or toward) v).

(I) Let D be an m -colored digraph such that:

- (i) $|\xi(v)| \leq 2$ for each $v \in V(D)$.
- (ii) There exists a fixed color c_i such that $c_i \in \xi(v)$ for each $v \in V(D)$.

Then D has a kernel by monochromatic paths.

Proof. Clearly the $\mathcal{C}_C(D)$ is bipartite. ■

(II) Let D be an m -colored digraph such that:

- (i) $|\xi(v)| \leq 2$ for each $v \in V(D)$.
- (ii) There exist two fixed colors c_i, c_j such that $|\{c_i, c_j\} \cap \xi(v)| = 1$ for each $v \in V(D)$.

Then D has a kernel by monochromatic paths.

Proof. $\mathcal{C}_C(D)$ is bipartite. ■

(III) Let H be a digraph possibly with loops and let D be a digraph whose arcs are colored with the vertices of H . A directed walk (path), W in D is an H -walk (H -path) if the consecutive color encountered on W form a directed walk in H . A set $N \subseteq V(D)$ is an H -kernel if no two vertices of N have an H -path between them and any $u \in V(D) \setminus N$ reaches some $v \in N$ on an H -path. The concept of H -walk was first introduced by Linek and Sands (1996). This concept was studied later by several authors.

Since $V(\mathcal{C}_C(D)) \subseteq V(H)$, the question that we can do us is the the next: What structure or substructures must $\mathcal{C}_C(D)$ have respect to the digraph H in order to ensure the existence of H -kernels in D ?

This questions will be studied in a forthcoming paper (H -kernels, Hortensia Galeana-Sánchez and Rocío Sánchez López).

Acknowledgement

The author wish to thank the anonymous referees for many suggestions which improved the rewriting of this paper.

REFERENCES

- [1] J.M. Le Bars, *Counterexample of the 0 – 1 law for fragments of existential second-order logic; an overview*, Bull. Symbolic Logic **9** (2000) 67–82.
- [2] J.M. Le Bars, *The 0 – 1 law fails for frame satisfiability of propositional model logic*, Proceedings of the 17th Symposium on Logic in Computer Science (2002) 225–234.
- [3] C. Berge, *Graphs* (North-Holland, Amsterdam, 1985).
- [4] E. Boros and V. Gurvich, *Perfect graphs, kernels and cores of cooperative games*, Discrete Math. **306** (2006) 2336–2354.
- [5] A.S. Fraenkel, *Combinatorial game theory foundations applied to digraph kernels*, Electronic J. Combin. **4** (2) (1997) #R10.
- [6] A.S. Fraenkel, *Combinatorial games: selected bibliography with a succinct gourmet introduction*, Electronic J. Combin. **14** (2007) #DS2.
- [7] G. Hahn, P. Ille and R. Woodrow, *Absorbing sets in arc-coloured tournaments*, Discrete Math. **283** (2004) 93–99.
- [8] H. Galeana-Sánchez, *On monochromatic paths and monochromatic cycles in edge coloured tournaments*, Discrete Math. **156** (1996) 103–112.
- [9] H. Galeana-Sánchez, *Kernels in edge-coloured digraphs*, Discrete Math. **184** (1998) 87–99.
- [10] H. Galeana-Sánchez and R. Rojas-Monroy, *A counterexample to a conjecture on edge-coloured tournaments*, Discrete Math. **282** (2004) 275–276.
- [11] H. Galeana-Sánchez and R. Rojas-Monroy, *On monochromatic paths and monochromatic 4-cycles in edge coloured bipartite tournaments*, Discrete Math. **285** (2004) 313–318.
- [12] G. Gutin and J. Bang-Jensen, *Digraphs: Theory, Algorithms and Applications* (Springer-Verlag, London, 2001).

- [13] T.W. Haynes, T. Hedetniemi and P.J. Slater, *Domination in Graphs (Advanced Topics)*, Marcel Dekker Inc., 1998.
- [14] T.W. Haynes, T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker Inc., 1998).
- [15] J. von Leeuwen, *Having a Grundy Numbering is NP-complete*, Report 207 Computer Science Department, University Park, PA, 1976, Pennsylvania State University.
- [16] B. Sands, N. Sauer and R. Woodrow, *On monochromatic paths in edge-coloured digraphs*, *J. Combin. Theory (B)* **33** (1982) 271–275.
- [17] I. Włoch, *On imp-sets and kernels by monochromatic paths in duplication*, *Ars Combin.* **83** (2007) 93–99.
- [18] I. Włoch, *On kernels by monochromatic paths in the corona of digraphs*, *Cent. Eur. J. Math.* **6** (2008) 537–542.

Received 24 November 2009

Revised 2 December 2010

Accepted 27 January 2011