

## SOME RESULTS ON SEMI-TOTAL SIGNED GRAPHS <sup>1</sup>

DEEPA SINHA AND PRAVIN GARG

*Centre for Mathematical Sciences*  
*Banasthali University*  
*Banasthali–304022, Rajasthan, India*

**e-mail:** [deepa.sinha2001@yahoo.com](mailto:deepa.sinha2001@yahoo.com)  
[garg.praavin@gmail.com](mailto:garg.praavin@gmail.com)

### Abstract

A *signed graph* (or *sigraph* in short) is an ordered pair  $S = (S^u, \sigma)$ , where  $S^u$  is a graph  $G = (V, E)$ , called the *underlying graph* of  $S$  and  $\sigma : E \rightarrow \{+, -\}$  is a function from the edge set  $E$  of  $S^u$  into the set  $\{+, -\}$ , called the *signature* of  $S$ . The  $\times$ -*line sigraph* of  $S$  denoted by  $L_{\times}(S)$  is a sigraph defined on the line graph  $L(S^u)$  of the graph  $S^u$  by assigning to each edge  $ef$  of  $L(S^u)$ , the product of signs of the adjacent edges  $e$  and  $f$  in  $S$ . In this paper, first we define *semi-total line sigraph* and *semi-total point sigraph* of a given sigraph and then characterize balance and consistency of semi-total line sigraph and semi-total point sigraph.

**Keywords:** sigraph, semi-total line sigraph, semi-total point sigraph, balanced sigraph, consistent sigraph.

**2010 Mathematics Subject Classification:** Primary: 05C22;  
Secondary: 05C75.

### 1. INTRODUCTION

For standard terminology and notation in graph theory we refer Harary [14] and West [21] and Zaslavsky [22, 23] for sigraphs. Throughout the text, we consider finite, undirected graph with no loops or multiple edges.

---

<sup>1</sup>Research supported by the Department of Science and Technology (Govt. of India), New Delhi, India under the Project SR/S4/MS: 409/06.

A *signed graph* (or *sigraph* in short; see [7, 11]) is an ordered pair  $S = (S^u, \sigma)$ , where  $S^u$  is a graph  $G = (V, E)$ , called the *underlying graph* of  $S$  and  $\sigma : E \rightarrow \{+, -\}$  is a function from the edge set  $E$  of  $S^u$  into the set  $\{+, -\}$ , called the *signature* of  $S$ . Alternatively, the sigraph can be written as  $S = (V, E, \sigma)$ , with  $V, E, \sigma$  in the above sense. Let  $E^+(S) = \{e \in E(G) : \sigma(e) = +\}$  and  $E^-(S) = \{e \in E(G) : \sigma(e) = -\}$ . The elements of  $E^+(S)$  and  $E^-(S)$  are called *positive* and *negative* edges of  $S$ , respectively. A sigraph is said to be *homogeneous* if all its edges are of the same sign and *heterogeneous* otherwise.

A sigraph  $S$  is called a *regular sigraph* if the number of positive edges,  $d^+(v)$  incident at a vertex  $v$  in  $S$ , is independent of the choice of  $v$  in  $S$  and the number of negative edges,  $d^-(v)$  incident at a vertex  $v$  in  $S$  is also independent of the choice of  $v$  in  $S$ , i.e.,  $S$  is a sigraph of order  $n$  and regular of degree  $k = i + j$ , where  $i = d^+(v)$  is the positive degree of  $v$  in  $S$  and  $j = d^-(v)$  is the negative degree of  $v$  in  $S$ .

For a sigraph  $S$ , Behzad and Chartrand [7] defined its *line sigraph*  $L(S)$  as the sigraph in which the edges of  $S$  are represented as vertices, two of these vertices are defined adjacent whenever the corresponding edges in  $S$  have a vertex in common and any such edge  $ef$  is defined to be negative whenever both  $e$  and  $f$  are negative edges in  $S$ . In [12], the author introduced a variation of the above standard notion of line sigraph  $L(S)$  of a given sigraph  $S$  as follows:  $L_{\times}(S)$  is a sigraph defined on the line graph  $L(S^u)$  of the graph  $S^u$  by assigning to each edge  $ef$  of  $L(S^u)$ , the product of signs of the adjacent edges  $e$  and  $f$  of  $S$ .  $L_{\times}(S)$  is called the  $\times$ -*line sigraph* of  $S$ .

A path in a sigraph  $S$  is said to be *all-negative* if each of its edge is negative. A cycle in a sigraph  $S$  is said to be *all-positive*(*all-negative*) if each of its edge is positive (negative). A cycle in a sigraph  $S$  is said to be *positive* if it contains an even number of negative edges. A given sigraph  $S$  is said to be *balanced* if every cycle in  $S$  is positive, i.e., it contains an even number of negative edges [4, 10, 13]. A spectral characterization of balanced sigraphs was given by Acharya [2]. Harary and Kabell [15, 16] developed a simple algorithm to get balanced sigraphs and also enumerated them. The following important lemma on balanced sigraph is given by Zaslavsky.

**Lemma 1** [24]. *A signed graph in which every chordless cycle is positive, is balanced.*

A *marked signed graph* is an ordered pair  $S_{\mu} = (S, \mu)$ , where  $S = (S^u, \sigma)$  is a sigraph and  $\mu : V(S^u) \rightarrow \{+, -\}$  is a function from the vertex set  $V(S^u)$

of  $S^u$  into the set  $\{+, -\}$ , called a *marking* of  $S$ . A cycle  $Z$  in  $S_\mu$  is said to be *consistent* if it contains an even number of negative vertices. A given sigraph  $S$  is said be *consistent* if every cycle in it is consistent [8, 9]. To this end, we define the following *canonical marking* on  $S$ : for each vertex  $v \in V(S)$ ,

$$\mu(v) = \prod_{e \in E_v} \sigma(e)$$

where  $E_v$  is set of edges  $e$  incident at  $v$  in  $S$ .

The *semi-total line graph*  $T_1(G)$  of a graph  $G$  [19] is the graph whose vertex set is  $V(G) \cup E(G)$  where  $V(G)$  and  $E(G)$  are vertex set and edge set of  $G$ , respectively and in  $T_1(G)$  two vertices are adjacent if and only if (i) they are adjacent edges in  $G$  (ii) one is a vertex and the other is an edge in  $G$  incident to it.




Figure 1. Showing  $T_1(S)$  and  $T_2(S)$  of a sigraph  $S$ .

The *semi-total point graph*  $T_2(G)$  of a graph  $G$  [19] is the graph whose vertex set is  $V(G) \cup E(G)$  where  $V(G)$  and  $E(G)$  are vertex set and edge set of  $G$ , respectively and in  $T_2(G)$  two vertices are adjacent if and only if (i) they

are adjacent vertices in  $G$ , (ii) one is a vertex and the other is an edge in  $G$  incident to it.

Let  $S = (V, E, \sigma)$  be any sigraph. Its *semi-total line sigraph*  $T_1(S)$  [as shown in Figure 1] has  $T_1(S^u)$  as its underlying graph and for any edge  $uv$  of  $T_1(S^u)$

$$\sigma_{T_1}(uv) = \begin{cases} \sigma(u)\sigma(v) & \text{if } u, v \in E, \\ \sigma(v) & \text{if } u \in V \text{ and } v \in E. \end{cases}$$

Let  $S = (V, E, \sigma)$  be any sigraph. Its *semi-total point sigraph*  $T_2(S)$  [as shown in Figure 1] has  $T_2(S^u)$  as its underlying graph and for any edge  $uv$  of  $T_2(S^u)$

$$\sigma_{T_2}(uv) = \begin{cases} \sigma(uv) & \text{if } u, v \in V, \\ \sigma(u) \prod_{e \in E_v} \sigma(e) & \text{if } u \in E \text{ and } v \in V. \end{cases}$$

We observe that the  $\times$ -line sigraph,  $L_\times(S)$  is an induced subsigraph of  $T_1(S)$  and  $S$  is an induced subsigraph of  $T_2(S)$ .

## 2. BALANCED SEMI-TOTAL LINE SIGRAPH

In this section, we obtain a characterization of balanced semi-total line sigraph.

**Theorem 2** [6]. *The  $\times$ -line sigraph  $L_\times(S)$  of a sigraph  $S$  is a balanced sigraph.*

**Theorem 3** [18]. *A sigraph  $S = (S^u, \sigma)$  is balanced if and only if there exists a marking  $\mu$  of its vertices such that for each edge  $uv$  in  $S$  one has  $\sigma(uv) = \mu(u)\mu(v)$ .*

**Theorem 4.** *The semi-total line sigraph  $T_1(S)$  of a sigraph  $S$  is a balanced sigraph.*

**Proof.** By the definition of  $T_1(S)$ , it contains  $L_\times(S)$  as induced subsigraph, triangles formed by the adjacent edges  $e$  and  $f$  in  $S$  and the vertex  $v$  such that  $e \cap f = \{v\}$  and cycles formed by the symmetric difference of these triangles and cycles in  $L_\times(S)$ . Since  $L_\times(S)$  is a balanced sigraph due to Theorem 2, we have to only show that triangles and cycles formed as above in  $T_1(S)$  are positive.

*Case (i).* Suppose  $e$  and  $f$  both are positive adjacent edges with  $v$  as their common vertex in  $S$ , then  $ef$  will be a positive edge in  $T_1(S)$ . Now by the definition of  $T_1(S)$ , the triangle formed by the vertices  $e$ ,  $f$  and  $v$  does not contain any negative edge. Thus, such triangles are positive.

*Case (ii).* Suppose  $e$  and  $f$  both are negative adjacent edges with  $v$  as their common vertex in  $S$ , then  $ef$  will be a positive edge in  $T_1(S)$ . Now by the definition of  $T_1(S)$ , the triangle formed by the vertices  $e$ ,  $f$  and  $v$  contain two negative edges. Thus, such triangles are positive too.

*Case (iii).* Suppose  $e$  and  $f$  are edges of opposite parity and they are adjacent with  $v$  as their common vertex in  $S$ , then  $ef$  will be a negative edge in  $T_1(S)$ . Now by the definition of  $T_1(S)$ , the triangle formed by the vertices  $e$ ,  $f$  and  $v$  contain two negative edges. Thus, such triangles are also positive.

Now, due to Lemma 1, it follows that  $T_1(S)$  is a balanced sigraph. Hence the theorem. ■

### 3. CONSISTENT SEMI-TOTAL LINE SIGRAPH

Beineke and Harary [8, 9] were the first to pose the problem of characterizing consistent marked graphs, which was subsequently settled by Acharya [1, 2] and Hoede [17]. Acharya and Sinha obtained consistency of sigraphs that satisfy certain sigraph equations in [20, 5]. In this section, first we define a  $\mu_1$ -marking and then obtain a characterization of  $\mu_1$ -consistent semi-total line sigraph.

For any sigraph  $S = (S^u, \sigma)$ , we define  $\mu_1$ -marking in semi-total line sigraph  $T_1(S)$  as  $\mu_1 : V(T_1(S)) \rightarrow \{+, -\}$  such that

$$\mu_1(v_i) = \prod_{e_j \in E_{v_i}} \sigma(e_j)$$

and

$$\mu_1(e_i) = \sigma(e_i).$$

**Theorem 5** [17]. *A marked graph  $G_\mu$  is consistent if and only if for any spanning tree  $T$  of  $G$  all fundamental cycles with respect to  $T$  are consistent and all common paths of pairs of those fundamental cycles have end vertices carrying the same marks.*

**Theorem 6.** *The semi-total line sigraph  $T_1(S)$  of a sigraph  $S$  is  $\mu_1$ -consistent if and only if the following conditions hold in  $S$ :*

- (i) *each cycle  $Z$  in  $S$  is homogeneous and positive,*
- (ii) *if  $d(v) \geq 3$ , then  $d^-(v) = 0$  for every vertex  $v$  in  $S$ .*

**Proof.** *Necessity:* Suppose  $T_1(S)$  of a sigraph  $S$  is  $\mu_1$ -consistent. Since  $L_\times(S)$  is an induced subsigraph of  $T_1(S)$  and  $T_1(S)$  is  $\mu_1$ -consistent, it follows that  $L_\times(S)$  is  $\mu_1$ -consistent. Let  $Z$  be a cycle in  $S$  and  $v \in V(Z)$ . Let  $d(v) = 2$ . If possible  $d^-(v) = 1$ , then let a positive edge  $e$  and a negative edge  $f$  be incident at  $v$ . Due to  $\mu_1$ -marking in  $T_1(S)$ , there is a  $\mu_1$ -consistent cycle  $Z_1$  having one positively marked vertex  $e$  and two negatively marked vertices  $v$  and  $f$  in  $T_1(S)$ . Let  $Z_2$  be a  $\mu_1$ -consistent cycle in  $L_\times(S)$  having the edge  $ef$ . Now, taking the symmetric difference of the edge sets of  $Z_1$  and  $Z_2$ , we get a  $\mu_1$ -inconsistent cycle in  $T_1(S)$ , since the end vertices of the common edge  $ef$  are oppositely marked. Thus, a contradiction to the assumption that  $T_1(S)$  is  $\mu_1$ -consistent. That means, each cycle  $Z$  in  $S$  is homogeneous. Again, the edges of each cycle  $Z$  in  $S$  create a cycle in  $L_\times(S)$  and each cycle in  $L_\times(S)$  has an even number of negatively marked vertices. So, each cycle  $Z$  in  $S$  has an even number of negative edges. That means, each cycle  $Z$  in  $S$  is positive. Thus, (i) follows.

If for a vertex  $v$  in  $S$ ,  $d^-(v) \geq 3$ , then any of the three negative edges incident to  $v$  will form a  $\mu_1$ -inconsistent triangle in  $L_\times(S)$ , a contradiction that  $L_\times(S)$  is  $\mu_1$ -consistent. So,  $d^-(v) < 3$ . Now, if  $d(v) > 3$ , then  $d^-(v)$  being equal to one or two would contradict the fact that  $L_\times(S)$  is  $\mu_1$ -consistent. If  $d(v) = 3$ , then  $d^-(v)$  being equal to one again contradicts the fact that  $L_\times(S)$  is  $\mu_1$ -consistent. If  $d(v) = 3$  and  $d^-(v) = 2$ , then let a positive edge  $e$  and two negative edges  $f$  and  $g$  be incident on  $v$ . Now due to  $\mu_1$ -marking in  $T_1(S)$ , there is a  $\mu_1$ -inconsistent cycle  $Z$  having two positively marked vertices  $v$  and  $e$  and one negatively marked vertex  $f$  in  $T_1(S)$ , a contradiction to the assumption that  $T_1(S)$  is  $\mu_1$ -consistent. Thus, (ii) follows.

*Sufficiency:* Suppose both the conditions (i) and (ii) hold for a given sigraph  $S$ . We have to show that  $T_1(S)$  is  $\mu_1$ -consistent. By the definition of  $T_1(S)$ , it contains  $L_\times(S)$  as an induced subsigraph, triangles due to the adjacent edges  $e$  and  $f$  in  $S$  and the vertex  $v$  such that  $e \cap f = \{v\}$  and cycles formed by the symmetric difference of these triangles and cycles in  $L_\times(S)$ . By these conditions,  $S$  is either an all-negative cycle of even length

or the sigraph containing all-positive cycles and the end vertices of induced all-negative path do not lie on any cycle.

*Case (i).* Suppose  $S$  is an all-negative cycle of even length. That means,  $L_{\times}(S)$  has an even number of negatively marked vertices. That means,  $L_{\times}(S)$  is  $\mu_1$ -consistent. Now we have to see the  $\mu_1$ -consistency of the triangles formed by the edges of  $S$  and  $L_{\times}(S)$ . Suppose both  $e$  and  $f$  are negative adjacent edges with  $v$  as their common vertex in  $S$ . Then due to the  $\mu_1$ -marking of  $T_1(S)$ ,  $\mu_1(e) = \mu_1(f) = -$  and  $\mu_1(v) = +$ . Hence, the triangle formed by the vertices  $e$ ,  $f$  and  $v$  in  $T_1(S)$  contains two negatively marked vertices  $e$  and  $f$ . That means, such triangles are  $\mu_1$ -consistent. Now, since the vertices  $e$  and  $f$  in  $T_1(S)$  have the same marks, so due to Theorem 5, cycles formed by the symmetric difference of these triangles and cycles in  $L_{\times}(S)$  will be  $\mu_1$ -consistent. Hence  $T_1(S)$  is  $\mu_1$ -consistent.

*Case (ii).* Suppose  $S$  is the graph containing all-positive cycles and by the condition (ii), such cycles will be adjacent with positive edges only. That means the end vertices of induced all-negative path do not lie on any cycle and due to condition (ii), the end vertices of these induced all-negative path are of degree two. Let  $e$  and  $f$  be positive and negative adjacent edges, respectively with  $v$  as their common vertex in  $S$ , then by the  $\mu_1$ -marking of  $T_1(S)$ , there is a  $\mu_1$ -consistent cycle  $Z$  having one positively marked vertex  $e$  and two negatively marked vertices  $v$  and  $f$  in  $T_1(S)$ . Again, let both  $e$  and  $f$  be negative adjacent edges with  $v$  as their common vertex in  $S$ . Then, by the  $\mu_1$ -marking of  $T_1(S)$ , there is a consistent cycle  $Z$  having one positively marked vertex  $v$  and two negatively marked vertices  $e$  and  $f$  in  $T_1(S)$ . Hence  $T_1(S)$  is  $\mu_1$ -consistent. ■

#### 4. BALANCED SEMI-TOTAL POINT SIGGRAPH

In this section, we define a  $\mu_1$ -marking in semi-total point siggraph and obtain a characterization of balanced semi-total point siggraph.

For any siggraph  $S = (S^u, \sigma)$ , we define  $\mu_1$ -marking in semi-total point siggraph  $T_2(S)$  as  $\mu_1 : V(T_2(S)) \rightarrow \{+, -\}$  such that  $\mu_1(v_i) = \prod_{e_j \in E_{v_i}} \sigma(e_j)$  and  $\mu_1(e_i) = \sigma(e_i)$ .

**Theorem 7.** *The semi-total point siggraph  $T_2(S)$  of a siggraph  $S$  is balanced if and only if the following conditions hold in  $S$ :*

- (i) if  $e$  is a positive edge in  $S$  and  $u, v$  are the end vertices of  $e$ , then the number of negative edges incident at  $u$  and  $v$  are of the same parity,
- (ii) if  $e$  is a negative edge in  $S$  and  $u, v$  are the end vertices of  $e$ , then the number of negative edges incident at  $u$  and  $v$  are of the opposite parity.

**Proof.** *Necessity:* Suppose  $T_2(S)$  is a balanced sigraph, then every cycle in  $T_2(S)$  must have an even number of negative edges. The vertex  $e$  being adjacent to the vertices  $u$  and  $v$  in  $T_2(S)$ , where  $uv$  is an edge  $e$  in  $S$ , we get a triangle  $Z$  in  $T_2(S)$  due to the vertices  $u, v$  and  $e$  which is balanced due to hypothesis. Now,

*Case (i).* If  $e$  is a positive edge in  $Z$ , then the edges  $ue$  and  $ve$  must be of the same parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) = \prod_{e_j \in E_v} \sigma(e_j),$$

whence, the number of negative edges incident at  $u$  and  $v$  are of the same parity. Thus, (i) follows.

*Case (ii).* If  $e$  is a negative edge in  $Z$ , then the edges  $ue$  and  $ve$  must be of the opposite parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) \neq \prod_{e_j \in E_v} \sigma(e_j),$$

whence, the number of negative edges incident at  $u$  and  $v$  are of the opposite parity. Thus, (ii) follows.

*Sufficiency:* Suppose conditions (i) and (ii) hold for a given sigraph  $S$ . We have to show that  $T_2(S)$  is a balanced sigraph. Let  $e$  be an edge in  $S$  whose end vertices are  $u$  and  $v$ . By the definition of  $T_2(S)$ ,  $T_2(S)$  contains  $S$  as an induced subgraph, triangles due to the vertices  $u, v$  and  $e$  and the cycles due to the symmetric difference of these triangles and cycles in  $S$ .

By condition (i) and (ii), the sign of each edge in  $S$  is the product of  $\mu_1$ -marking of corresponding end vertices in  $S$ . So, using Theorem 3,  $S$  is balanced. Now, we have to only show that the triangles and cycles formed as above in  $T_2(S)$  are also positive.

By condition (i),  $e$  is a positive edge in  $S$  whose end vertices are  $u$  and  $v$  and the number of negative edges incident at  $u$  and  $v$  are of the same parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) = \prod_{e_j \in E_v} \sigma(e_j)$$

and

$$\sigma(e) = +.$$

Hence, by the definition of  $T_2(S)$ , the triangle due to the vertices  $u, v$  and  $e$ , contains either no negative edge or two negative edges. Thus, such triangles are positive.

By condition (ii),  $e$  is a negative edge in  $S$ , whose end vertices are  $u$  and  $v$  and the number of negative edges incident at  $u$  and  $v$  are of the opposite parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) \neq \prod_{e_j \in E_v} \sigma(e_j)$$

and

$$\sigma(e) = -.$$

Hence, by the definition of  $T_2(S)$ , the triangle due to the vertices  $u, v$  and  $e$ , contains two negative edges. Thus, such triangles are positive.

Thus, due to Lemma 1, it follows that  $T_1(S)$  is a balanced sigraph. Hence the theorem. ■

**Corollary 8.** *The semi-total point sigraph  $T_2(S)$  of a regular heterogeneous sigraph  $S$  is not balanced.*

### 5. CONSISTENT SEMI-TOTAL POINT SIGRAPH

In this section, we obtain a characterization of  $\mu_1$ -consistent semi-total point sigraph.

**Theorem 9.** *The semi-total point sigraph  $T_2(S)$  of a sigraph  $S = (S^u, \sigma)$  is  $\mu_1$ -consistent if and only if the following conditions hold in  $S$ :*

- (i) *if  $e$  is a positive edge in  $S$  and  $u, v$  are the end vertices of  $e$ , then the vertices  $u$  and  $v$  are of the same parity,*
- (ii) *if  $e$  is a negative edge in  $S$  and  $u, v$  are the end vertices of  $e$ , then the vertices  $u$  and  $v$  are of the opposite parity,*
- (iii) *each cycle  $Z$  in  $S$  is all-positive and if for any  $v \in V(Z)$*

$$\prod_{e \in E_v} \sigma(e) = -,$$

*then  $Z$  is of even length.*

**Proof. Necessity:** Suppose  $T_2(S)$  is  $\mu_1$ -consistent, then every cycle in  $T_2(S)$  must have an even number of negative vertices. Since  $T_2(S)$  has  $S$  as an induced subgraph and  $T_2(S)$  is  $\mu_1$ -consistent, it follows that the induced  $S$  in  $T_2(S)$  is  $\mu_1$ -consistent. Now, the vertex  $e$  being adjacent to the vertices  $u$  and  $v$  in  $T_2(S)$ , where  $uv$  is an edge  $e$  in  $S$ , we get a triangle  $Z$  in  $T_2(S)$  due to the vertices  $u, v$  and  $e$  which is  $\mu_1$ -consistent by hypothesis.

*Case (i).* Let  $e$  be a positive edge in  $Z$  and  $\mu_1(e) = \sigma(e)$ , then the number of negative edges incident at  $u$  and  $v$  are of the same parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) = \prod_{e_j \in E_v} \sigma(e_j).$$

This implies, the vertices  $u$  and  $v$  are of the same parity. Thus, (i) follows.

*Case (ii).* Let  $e$  be a negative edge in  $Z$  and  $\mu_1(e) = \sigma(e)$ , then the number of negative edges incident at  $u$  and  $v$  are of the opposite parity. That means,

$$\prod_{e_i \in E_u} \sigma(e_i) \neq \prod_{e_j \in E_v} \sigma(e_j).$$

This implies, the vertices  $u$  and  $v$  are of the opposite parity. Thus, (ii) follows.

Now, let  $e$  be a negative edge contained in a cycle  $Z$  in  $S$  and  $u, v$  are the end vertices of  $e$ , then due to condition (ii),  $u$  and  $v$  are of the opposite parity. That means,

$$\mu_1(u) \neq \mu_1(v)$$

and

$$\mu_1(e) = \sigma(e)$$

whence, we get a triangle  $Z_1$  in  $T_2(S)$  due to the vertices  $u, v$  and  $e$  which is  $\mu_1$ -consistent by hypothesis. Let  $Z_2$  be the consistent cycle in  $T_2(S)$  containing the edge  $e = uv$ . Now, if we take the symmetric difference of  $Z_1$  and  $Z_2$ , then by Theorem 5,  $T_2(S)$  will not be  $\mu_1$ -consistent, a contradiction of our hypothesis. Thus,  $e$  can not be contained in any cycle in  $S$ . This implies, each cycle  $Z$  in  $S$  is all-positive. Now, let for any  $v \in V(Z)$

$$\prod_{e \in E_v} \sigma(e) = -.$$

Since  $Z$  is all-positive, then by condition (i), for each  $v \in V(Z)$

$$\prod_{e \in E_v} \sigma(e) = -,$$

whence, for each  $v \in V(Z)$

$$\mu_1(v) = -.$$

Since  $S$  is an induced subgraph of  $T_2(S)$ , the cycle  $Z$  will be the cycle in  $T_2(S)$  and  $Z$  will be  $\mu_1$ -consistent due to hypothesis. It follows that,  $Z$  is of even length. Thus (iii) follows.

*Sufficiency:* Suppose conditions (i), (ii) and (iii) hold for a given sigraph  $S$ . We have to show that  $T_2(S)$  is  $\mu_1$ -consistent. Let  $e$  be an edge in  $S$  whose end vertices are  $u$  and  $v$ . By the definition of  $T_2(S)$ ,  $T_2(S)$  contains  $S$  as an induced subgraph, triangles due to the vertices  $u, v$  and  $e$  and cycles due to the symmetric difference of these triangles and cycles in  $S$ .

By the condition (i),  $e$  is a positive edge in  $S$  whose end vertices are  $u$  and  $v$  and the vertices  $u$  and  $v$  are of the same parity. That means,

$$\mu_1(u) = \mu_1(v)$$

and

$$\mu_1(e) = \sigma(e) = +.$$

By the definition of  $T_2(S)$ , the triangle due to the vertices  $u, v$  and  $e$ , contains either no negatively marked vertex or two negatively marked vertices. Therefore, such triangles are  $\mu_1$ -consistent.

By the condition (ii),  $e$  is a negative edge in  $S$  whose end vertices are  $u$  and  $v$  and the vertices  $u$  and  $v$  are of the opposite parity. That means,

$$\mu_1(u) \neq \mu_1(v)$$

and

$$\mu_1(e) = \sigma(e) = -.$$

By the definition of  $T_2(S)$ , the triangle due to the vertices  $u, v$  and  $e$ , contains two negatively marked vertices. Therefore, such triangles are  $\mu_1$ -consistent.

By the condition (iii), each cycle  $Z$  in  $S$  is all-positive. Let  $e$  be a positive edge of such a cycle and  $u, v$  be the end vertices of  $e$ . Then, by condition (i) and the definition of  $T_2(S)$ , we get a  $\mu_1$ -consistent triangle  $Z_1$

due to the vertices  $u$ ,  $v$  and  $e$ , containing the edge  $e$ . Let  $Z_2$  be the  $\mu_1$ -consistent cycle in  $S$  containing the edge  $e$ . Now, we take the symmetric difference of  $Z_1$  and  $Z_2$ . Then, by Theorem 5, we get a  $\mu_1$ -consistent cycle. Since cycles in  $S$  are cycles in  $T_2(S)$ ,  $Z$  will be the cycle in  $T_2(S)$  and if for any  $v \in V(Z)$

$$\prod_{e \in E_v} \sigma(e) = -$$

then  $Z$  is of even length. Thus,  $Z$  is a  $\mu_1$ -consistent cycle. Hence the theorem. ■

**Corollary 10.** *The semi-total point sigraph  $T_2(S)$  of a regular heterogeneous sigraph  $S$  is not  $\mu_1$ -consistent.*

**Corollary 11.** *The semi-total point sigraph  $T_2(S)$  of a heterogeneous cycle  $S$  is not  $\mu_1$ -consistent.*

### Acknowledgement

The authors express their gratitude to Dr. B.D. Acharya for encouraging them to obtain more results on any given idea or theme as also for his constant readiness for discussions, which almost invariably yielded deeper insights. His rigorous efforts in going through the paper have helped the authors to bring the paper in the present form.

### REFERENCES

- [1] B.D. Acharya, *A characterization of consistent marked graphs*, National Academy, Science Letters, India **6** (1983) 431–440.
- [2] B.D. Acharya, *A spectral criterion for cycle balance in networks*, J. Graph Theory **4** (1981) 1–11.
- [3] B.D. Acharya, *Some further properties of consistent marked graphs*, Indian J. Pure Appl. Math. **15** (1984) 837–842.
- [4] B.D. Acharya and M. Acharya, *New algebraic models of a social system*, Indian J. Pure Appl. Math. **17** (1986) 150–168.
- [5] B.D. Acharya, M. Acharya and D. Sinha, *Characterization of a signed graph whose signed line graph is  $s$ -consistent*, Bull. Malays. Math. Sci. Soc. **32** (2009) 335–341.

- [6] M. Acharya,  $\times$ -line sigraph of a sigraph, J. Combin. Math. Combin. Comp. **69** (2009) 103–111.
- [7] M. Behzad and G.T. Chartrand, *Line coloring of signed graphs*, Element der Mathematik, **24** (1969) 49–52.
- [8] L.W. Beineke and F. Harary, *Consistency in marked graphs*, J. Math. Psychol. **18** (1978) 260–269.
- [9] L.W. Beineke and F. Harary, *Consistent graphs with signed points*, Riv. Math. per. Sci. Econom. Sociol. **1** (1978) 81–88.
- [10] D. Cartwright and F. Harary, *Structural Balance: A generalization of Heiders Theory*, Psych. Rev. **63** (1956) 277–293.
- [11] G.T. Chartrand, *Graphs as Mathematical Models* (Prindle, Weber and Schmid, Inc., Boston, Massachusetts, 1977).
- [12] M.K. Gill, *Contribution to some topics in graph theory and its applications*, Ph.D. Thesis, (Indian Institute of Technology, Bombay, 1983).
- [13] F. Harary, *On the notion of balanced signed graphs*, Mich. Math. J. **2** (1953) 143–146.
- [14] F. Harary, *Graph Theory* (Addison-Wesley Publ. Comp., Reading, Massachusetts, 1969).
- [15] F. Harary and J.A. Kabell, *A simple algorithm to detect balance in signed graphs*, Math. Soc. Sci. **1** (1980/81) 131–136.
- [16] F. Harary and J.A. Kabell, *Counting balanced signed graphs using marked graphs*, Proc. Edinburgh Math. Soc. **24** (1981) 99–104.
- [17] C. Hoede, *A characterization of consistent marked graphs*, J. Graph Theory **16** (1992) 17–23.
- [18] E. Sampathkumar, *Point-signed and line-signed graphs*, Karnatak Univ. Graph Theory Res. Rep. **1** (1973), also see Abstract No. 1 in Graph Theory Newsletter **2** (1972), Nat. Acad. Sci.-Letters **7** (1984) 91–93.
- [19] E. Sampathkumar and S.B. Chikkodimath, *Semitotal graphs of a graph-I*, J. Karnatak Univ. Sci. **XVIII** (1973) 274–280.
- [20] D. Sinha, *New frontiers in the theory of signed graph*, Ph.D. Thesis (University of Delhi, Faculty of Technology, 2005).
- [21] D.B. West, *Introduction to Graph Theory* (Prentice-Hall, India Pvt. Ltd., 1996).
- [22] T. Zaslavsky, *A mathematical bibliography of signed and gain graphs and allied areas*, Electronic J. Combinatorics #DS8 (vi+151pp)(1999)

- [23] T. Zaslavsky, *Glossary of signed and gain graphs and allied areas*, II Edition, Electronic J. Combinatorics, #DS9(1998).
- [24] T. Zaslavsky, *Signed analogs of bipartite graphs*, Discrete Math. **179** (1998) 205–216.

Received 11 October 2009  
Revised 30 September 2010  
Accepted 1 October 2010