

BOUNDS ON THE SIGNED 2-INDEPENDENCE NUMBER IN GRAPHS

LUTZ VOLKMANN

Lehrstuhl II für Mathematik
RWTH-Aachen University
52056 Aachen, Germany

e-mail: volkm@math2.rwth-aachen.de

Abstract

Let G be a finite and simple graph with vertex set $V(G)$, and let $f : V(G) \rightarrow \{-1, 1\}$ be a two-valued function. If $\sum_{x \in N[v]} f(x) \leq 1$ for each $v \in V(G)$, where $N[v]$ is the closed neighborhood of v , then f is a signed 2-independence function on G . The weight of a signed 2-independence function f is $w(f) = \sum_{v \in V(G)} f(v)$. The maximum of weights $w(f)$, taken over all signed 2-independence functions f on G , is the signed 2-independence number $\alpha_s^2(G)$ of G .

In this work, we mainly present upper bounds on $\alpha_s^2(G)$, as for example $\alpha_s^2(G) \leq n - 2\lceil \Delta(G)/2 \rceil$, and we prove the Nordhaus-Gaddum type inequality $\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n + 1$, where n is the order and $\Delta(G)$ is the maximum degree of the graph G . Some of our theorems improve well-known results on the signed 2-independence number.

Keywords: bounds, signed 2-independence function, signed 2-independence number, Nordhaus-Gaddum type result.

2010 Mathematics Subject Classification: 05C69.

1. INTRODUCTION

Domination and independence in graphs are well studied. The literature on this subject has been surveyed and detailed in the two books by Haynes, Hedetniemi and Slater [2, 3].

All graphs considered are undirected, simple and finite. The vertex set and edge set of a graph G are denoted by $V(G)$ and $E(G)$. The *order* $n = n(G)$ and *size* $q = q(G)$ of a graph G is the number of vertices and edges, respectively. The *open neighborhood* of $v \in V(G)$ is $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$ and the *closed neighborhood* of v is $N_G[v] = N_G(v) \cup \{v\}$. The *degree* of v in G ,

denoted by $d_G(v)$, is the cardinality of $N_G(v)$. We write $\Delta(G)$ and $\delta(G)$ for the *maximum* and *minimum degree* of G . If the graph G is clear from context, we simply use $N(v)$, $N[v]$, $d(v)$, Δ and δ instead of $N_G(v)$, $N_G[v]$, $d_G(v)$, $\Delta(G)$ and $\delta(G)$, respectively. For two disjoint subsets A and B of $V(G)$, let $e(A, B)$ denote the number of edges between A and B . A graph G is *r-partite* with vertex classes V_1, V_2, \dots, V_r if $V(G) = V_1 \cup V_2 \cup \dots \cup V_r$, $V_i \cap V_j = \emptyset$ whenever $1 \leq i < j \leq r$, and no edge joins two vertices in the same class. The subgraph of G induced by A is denoted by $G[A]$. The complete graph of order n is denoted by K_n . A graph is *K_{r+1} -free* if it does not contain the complete graph K_{r+1} as a subgraph. The *complement* of a graph G is denoted by \overline{G} .

For a two-valued function $f : V(G) \rightarrow \{-1, 1\}$, the *weight* of f is $w(f) = \sum_{v \in V(G)} f(v)$. For a subset $A \subseteq V(G)$, we define $f(A) = \sum_{v \in A} f(v)$ and so $w(f) = f(V(G))$. For a vertex v in $V(G)$, we denote $f(N[v])$ by $f[v]$ for notational convenience. The function f is defined in [1] to be a *signed dominating function* of G if $f[v] = f(N[v]) \geq 1$ for every $v \in V(G)$. The *signed domination number* of G is the minimum weight of a signed dominating function on G .

The function $f : V(G) \rightarrow \{-1, 1\}$ is defined in [7] to be a *signed 2-independence function* on G if $f[v] = f(N[v]) \leq 1$ for every $v \in V(G)$. The *signed 2-independence number* $\alpha_s^2(G)$ of G is the maximum weight of a signed 2-independence function on G . Hence the signed 2-independence number is a certain dual to the signed domination number of a graph. Results on the signed 2-independence number can be found in [4, 5, 7].

In this paper we continue the investigations of the signed 2-independence number. We mainly present upper bounds on $\alpha_s^2(G)$ for general graphs and K_{r+1} -free graphs. In addition, we prove the Nordhaus-Gaddum type inequality $\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n + 1$. Some of our results improve known bounds on the signed 2-independence numbers of graphs given by Henning [4] in 2002 and Shan, Sohn and Kang [5] in 2003.

Zelinka [7] determined the signed 2-independence number of complete graphs, and he established a sharp upper bound on $\alpha_s^2(G)$ for regular graphs G .

Theorem 1 [7]. *If G is isomorphic to the complete graph K_n , then $\alpha_s^2(G) = 0$ when n is even and $\alpha_s^2(G) = 1$ when n is odd.*

Theorem 2 [7]. *If G is an r -regular graph of order n , then $\alpha_s^2(G) \leq n/(r + 1)$ when r is even and $\alpha_s^2(G) \leq 0$ when r is odd.*

2. MAIN RESULTS

Theorem 3. *If G is a graph of order n , then*

$$2 - n \leq \alpha_s^2(G) \leq n - 2 \left\lceil \frac{\Delta}{2} \right\rceil.$$

Proof. Let $w \in V(G)$ be a vertex of maximum degree $d(w) = \Delta$, and let f be a signed 2-independence function on G for which $f(V(G)) = \alpha_s^2(G)$. We define the two sets $P = \{v \in V(G) \mid f(v) = 1\}$ and $M = \{v \in V(G) \mid f(v) = -1\}$. If $|P| = p$ and $|M| = m$, then $n = p + m$ and $\alpha_s^2(G) = p - m = n - 2m$.

Assume first that $f(w) = 1$ and therefore $w \in P$. The condition $f[w] \leq 1$ leads to the inequality $|N(w) \cap P| - |N(w) \cap M| \leq 0$, and since w is a vertex of maximum degree, we have $|N(w) \cap P| + |N(w) \cap M| = \Delta$. Combining the last two inequalities, we deduce that $m \geq |N(w) \cap M| \geq \lceil \Delta/2 \rceil$, and this yields to

$$\alpha_s^2(G) = n - 2m \leq n - 2 \left\lceil \frac{\Delta}{2} \right\rceil.$$

Assume second that $f(w) = -1$ and so $w \in M$. As $f[w] \leq 1$ and $d(w) = \Delta$, we obtain $|N(w) \cap P| - |N(w) \cap M| \leq 2$ and $|N(w) \cap P| + |N(w) \cap M| = \Delta$. Combining these two inequalities, we conclude that

$$m \geq |N(w) \cap M| + 1 = \frac{2|N(w) \cap M| + 2}{2} \geq \frac{\Delta}{2}$$

and thus $m \geq \lceil \Delta/2 \rceil$. This implies $\alpha_s^2(G) = n - 2m \leq n - 2 \lceil \Delta/2 \rceil$ as above, and the upper bound on $\alpha_s^2(G)$ is proved.

For the first inequality define $f : V(G) \rightarrow \{-1, 1\}$ by $f(v) = 1$ for an arbitrary vertex $v \in V(G)$ and $f(x) = -1$ for each vertex $x \in V(G) - \{v\}$. Obviously, f is a signed 2-independence function on G of weight $2 - n$ and thus $\alpha_s^2(G) \geq 2 - n$. ■

If G is isomorphic to the star $K_{1,\Delta}$, then

$$\alpha_s^2(G) = n - 2 \left\lceil \frac{\Delta}{2} \right\rceil,$$

and therefore the upper bound on $\alpha_s^2(G)$ in Theorem 3 is sharp.

Corollary 4. *If G is a graph of order n , then $\alpha_s^2(G) = n$ if and only if G is the empty graph.*

Proof. If G is the empty graph, then $f : V(G) \rightarrow \{-1, 1\}$ with $f(v) = 1$ for each vertex $v \in V(G)$ is a signed 2-independence function on G of weight n and thus $\alpha_s^2(G) = n$.

Conversely, assume that $\alpha_s^2(G) = n$. If we suppose that G is not the empty graph, then $\Delta \geq 1$, and Theorem 3 leads to the contradiction $n = \alpha_s^2(G) \leq n - 2$. Therefore G is the empty graph, and the proof is complete. ■

Obviously, $\alpha_s^2(K_2) = 0 = n - 2$, and therefore equality in the left inequality of Theorem 3 is achieved. However, if G is a graph of order $n \geq 3$, then the next result improves the lower bound in Theorem 3.

Theorem 5. *If G is a graph of order $n \geq 3$, then $\alpha_s^2(G) \geq 4 - n$.*

Proof. If G has two non-adjacent vertices u and v , then $f : V(G) \rightarrow \{-1, 1\}$ with $f(u) = f(v) = 1$ and $f(x) = -1$ for each $x \in V(G) - \{u, v\}$ is a signed 2-independence function on G of weight $4 - n$ and thus $\alpha_s^2(G) \geq 4 - n$. Otherwise, G is the complete graph. If $n \geq 4$, then it follows from Theorem 1 that $\alpha_s^2(G) \geq 0 \geq 4 - n$, and if $n = 3$, then Theorem 1 implies that $\alpha_s^2(G) = 1 = 4 - n$. ■

As an application of Theorems 1, 2, 3 and Corollary 4, we will prove the following Nordhaus-Gaddum type result.

Theorem 6. *If G is a graph of order n , then*

$$\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n + 1$$

with equality if and only if n is odd and $G = K_n$ or $\overline{G} = K_n$.

Proof. Theorem 3 implies that

$$\begin{aligned} \alpha_s^2(G) + \alpha_s^2(\overline{G}) &\leq n - \Delta(G) + n - \Delta(\overline{G}) \\ (1) \qquad \qquad \qquad &= n - \Delta(G) + n - (n - \delta(G) - 1) \\ &= n + 1 - \Delta(G) + \delta(G), \end{aligned}$$

and the desired bound follows, since $\delta(G) - \Delta(G) \leq 0$. If n is odd and $G = K_n$ or $\overline{G} = K_n$, then we deduce from Theorem 1 and Corollary 4 that $\alpha_s^2(G) + \alpha_s^2(\overline{G}) = n + 1$.

If $\Delta(G) - \delta(G) \geq 1$, then the inequality chain (1) leads to $\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n$.

Assume now that $\Delta(G) = \delta(G) = \delta$, implying that G is δ -regular.

Assume first that n is even. If δ is even, then $\delta(\overline{G}) = n - \delta - 1$ is odd, and Theorem 2 implies that $\alpha_s^2(\overline{G}) \leq 0$ and thus $\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n$. If δ is odd, then Theorem 2 implies that $\alpha_s^2(G) \leq 0$ and thus $\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq n$.

Finally assume that n is odd. The handshaking lemma implies that δ and $\delta(\overline{G})$ are even. If $\delta = 0$ or $\delta(\overline{G}) = 0$, then $\overline{G} = K_n$ or $G = K_n$ and thus $\alpha_s^2(G) + \alpha_s^2(\overline{G}) = n + 1$. In the remaining case that $\delta \geq 2$ and $\delta(\overline{G}) \geq 2$, Theorem 2 shows that

$$\alpha_s^2(G) + \alpha_s^2(\overline{G}) \leq \frac{n}{\delta + 1} + \frac{n}{\delta(\overline{G}) + 1} \leq \frac{2n}{3} < n,$$

and the proof of Theorem 6 is complete. ■

The following upper bound on $\alpha_s^2(G)$ was obtained by Henning [4] in 2002.

Theorem 7 [4]. *If G is a connected graph of order $n \geq 2$ and size q , then*

$$\alpha_s^2(G) \leq \frac{4q - n}{5}.$$

We now improve the bound in Theorem 7.

Theorem 8. *If G is a connected graph of order $n \geq 2$ and size q , then*

$$\alpha_s^2(G) \leq \frac{4q + (2 - \delta - 2\lceil \delta/2 \rceil)n}{2 + \delta + 2\lceil \delta/2 \rceil}.$$

Proof. Let f be a signed 2-independence function on G for which $f(V(G)) = \alpha_s^2(G)$, and let P, M, p and m be defined as in the proof of Theorem 3. Then $n = p + m$ and $\alpha_s^2(G) = p - m = 2p - n$. The condition $f[v] \leq 1$ implies that $|N(v) \cap P| \leq |N(v) \cap M|$ for $v \in P$ and $|N(v) \cap P| \leq |N(v) \cap M| + 2$ for $v \in M$. Thus we obtain

$$\delta \leq d(v) = |N(v) \cap P| + |N(v) \cap M| \leq 2|N(v) \cap M|$$

and so $|N(v) \cap M| \geq \lceil \frac{\delta}{2} \rceil$ for each $v \in P$. Hence we deduce that

$$(2) \quad e(P, M) = \sum_{v \in P} |N(v) \cap M| \geq p \left\lceil \frac{\delta}{2} \right\rceil = (n - m) \left\lceil \frac{\delta}{2} \right\rceil.$$

In addition, we have

$$(3) \quad \begin{aligned} e(P, M) &= \sum_{v \in M} |N(v) \cap P| \leq \sum_{v \in M} (|N(v) \cap M| + 2) \\ &= 2|E(G[M])| + 2m. \end{aligned}$$

Combining (2) and (3), we find that

$$(4) \quad 2|E(G[M])| \geq p \left\lceil \frac{\delta}{2} \right\rceil - 2m.$$

Furthermore, we deduce from (2) that

$$(5) \quad \begin{aligned} e(P, M) + |E(G[P])| &= \sum_{v \in P} |N(v) \cap M| + \frac{1}{2} \sum_{v \in P} |N(v) \cap P| \\ &= \frac{1}{2} \sum_{v \in P} |N(v) \cap M| + \frac{1}{2} \sum_{v \in P} |N(v) \cap M| \\ &\quad + \frac{1}{2} \sum_{v \in P} |N(v) \cap P| \\ &= \frac{1}{2} \sum_{v \in P} |N(v)| + \frac{1}{2} \sum_{v \in P} |N(v) \cap M| \\ &\geq \frac{1}{2} p \delta + \frac{1}{2} p \left\lceil \frac{\delta}{2} \right\rceil. \end{aligned}$$

According to (4) and (5), we have

$$\begin{aligned} 2q &= 2e(P, M) + 2|E(G[P])| + 2|E(G[M])| \\ &\geq p\delta + 2p \left\lceil \frac{\delta}{2} \right\rceil - 2m = p \left(2 + \delta + 2 \left\lceil \frac{\delta}{2} \right\rceil \right) - 2n. \end{aligned}$$

Hence

$$p \leq \frac{2q + 2n}{2 + \delta + 2\lceil\delta/2\rceil},$$

and so we obtain the desired bound as follows

$$\alpha_s^2(G) = 2p - n \leq \frac{4q + (2 - \delta - 2\lceil\delta/2\rceil)n}{2 + \delta + 2\lceil\delta/2\rceil}.$$

■

Note that

$$\frac{4q + (2 - \delta - 2\lceil\delta/2\rceil)n}{2 + \delta + 2\lceil\delta/2\rceil} \leq \frac{4q - n}{5}$$

for $\delta \geq 1$, and therefore Theorem 8 is an improvement of Theorem 7.

For the next result, we use the famous theorem of Turán [6].

Theorem 9 [6]. *Let $r \geq 1$ be an integer. If G is a K_{r+1} -free graph of order n , then*

$$|E(G)| \leq \frac{r-1}{2r}n^2.$$

Theorem 10. *If G is a K_{r+1} -free graph of order n with $r \geq 2$ and minimum degree $\delta \geq 1$, then*

$$\alpha_s^2(G) \leq n + \frac{r(2 + \lceil\delta/2\rceil)}{r-1} - \sqrt{\left(\frac{r(2 + \lceil\delta/2\rceil)}{r-1}\right)^2 + \frac{4rn\lceil\delta/2\rceil}{r-1}}.$$

Proof. Let f, P, M, p and m be defined as in the proof of Theorem 3. By (2) and (3), we obtain

$$(6) \quad (n - m) \lceil\delta/2\rceil \leq e(P, M) \leq 2|E(G[M])| + 2m.$$

Since G is K_{r+1} -free, the induced subgraph $G[M]$ is also K_{r+1} -free, and hence it follows from Theorem 9 that $|E(G[M])| \leq (r - 1)m^2/2r$. Using (6), we obtain

$$(n - m) \lceil\delta/2\rceil \leq e(P, M) \leq \frac{r-1}{r}m^2 + 2m$$

and so

$$m^2 + \frac{r}{r-1}(2 + \lceil\delta/2\rceil)m - \frac{r}{r-1}n\lceil\delta/2\rceil \geq 0.$$

This yields

$$m \geq -\frac{r}{2(r-1)}(2 + \lceil\delta/2\rceil) + \sqrt{\left(\frac{r}{2(r-1)}(2 + \lceil\delta/2\rceil)\right)^2 + \frac{r}{r-1}n\lceil\delta/2\rceil},$$

and we obtain the desired bound as follows

$$\alpha_s^2(G) = n - 2m \leq n + \frac{r(2 + \lceil \delta/2 \rceil)}{r-1} - \sqrt{\left(\frac{r(2 + \lceil \delta/2 \rceil)}{r-1}\right)^2 + \frac{4rn\lceil \delta/2 \rceil}{r-1}}.$$

■

Since

$$\frac{r(2 + \lceil \delta/2 \rceil)}{r-1} - \sqrt{\left(\frac{r(2 + \lceil \delta/2 \rceil)}{r-1}\right)^2 + \frac{4rn\lceil \delta/2 \rceil}{r-1}} \leq \frac{3r}{r-1} - \sqrt{\left(\frac{3r}{r-1}\right)^2 + \frac{4rn}{r-1}}$$

for $\delta \geq 1$, the next known result is an immediate consequence of Theorem 10.

Corollary 11 [5]. *If G is an r -partite graph of order n with $r \geq 2$, then*

$$\alpha_s^2(G) \leq n + \frac{3r}{r-1} - \sqrt{\left(\frac{3r}{r-1}\right)^2 + \frac{4rn}{r-1}}.$$

REFERENCES

- [1] J.E. Dunbar, S.T. Hedetniemi, M.A. Henning and P.J. Slater, *Signed domination in graphs*, in: Graph Theory, Combinatorics, and Applications (John Wiley and Sons, Inc. 1, 1995) 311–322.
- [2] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, Inc., New York, 1998).
- [3] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Domination in Graphs, Advanced Topics* (Marcel Dekker, Inc., New York, 1998).
- [4] M.A. Henning, *Signed 2-independence in graphs*, *Discrete Math.* **250** (2002) 93–107. doi:10.1016/S0012-365X(01)00275-8
- [5] E.F. Shan, M.Y. Sohn and L.Y. Kang, *Upper bounds on signed 2-independence numbers of graphs*, *Ars Combin.* **69** (2003) 229–239.
- [6] P. Turán, *On an extremal problem in graph theory*, *Math. Fiz. Lapok* **48** (1941) 436–452.
- [7] B. Zelinka, *On signed 2-independence numbers of graphs*, manuscript.

Received 21 February 2012

Revised 3 September 2012

Accepted 3 September 2012

