

PACKING TREES INTO n -CHROMATIC GRAPHS

ANDRÁS GYÁRFÁS

Alfréd Rényi Institute of Mathematics

e-mail: gyarf. andras@mta.renyi.hu

Abstract

We show that if a sequence of trees T_1, T_2, \dots, T_{n-1} can be packed into K_n then they can be also packed into any n -chromatic graph.

Keywords: tree packing.

2010 Mathematics Subject Classification: 05C70.

Let T_i denote a tree with i edges. The author's tree packing conjecture [3] states that K_n has an edge disjoint decomposition into any given sequence T_1, T_2, \dots, T_{n-1} of trees. Gerbner, Keszegh and Palmer extended the conjecture by replacing K_n with an arbitrary n -chromatic graph (Conjecture 2 in [1]). Here we show that the extended conjecture follows from the original one. We need the following, perhaps folklore result (Theorem 1 in [4]) and for convenience we include its simple proof.

Lemma 1. *Let G be a k -chromatic graph with a proper k -coloring with k distinct colors. Suppose T is a tree on k vertices and each vertex of T is labeled with a different color from the same set of k colors. Then G contains a subtree that is label-isomorphic to T (labeled exactly the same way as T).*

Proof. The proof is by induction, the base step is trivial for $k = 1$. Let S_t denote the vertices of color t in a proper k -coloring of a k -chromatic graph G with a k -element color set C . Select a leaf vertex P in a tree T labeled with the k colors of C , suppose its label is i and assume P is adjacent in T with vertex Q labeled with j . Let A denote the set of vertices in S_j adjacent to at least one vertex of S_i . Observe that A is nonempty, otherwise G would be $(k - 1)$ -chromatic. The subgraph $G^* \subset G$ obtained by removing S_i and $S_j - A$ from $V(G)$ is $(k - 1)$ -chromatic, since the removed vertices form an independent set. Also, G^* is colored with the color set $C - i$. By induction, G^* contains a label-isomorphic copy of the tree $T - P$, its vertex with color j is in A , thus adjacent to a vertex in S_i , extending $T - P$ to a label-isomorphic copy of T . ■

Theorem 2. *Suppose that K_n has an edge disjoint decomposition into a given sequence T_1, T_2, \dots, T_{n-1} of trees and G is an n -chromatic graph. Then G contains edge disjoint copies of T_1, T_2, \dots, T_{n-1} .*

Proof. Let S_1, S_2, \dots, S_n be a partition of $V(G)$ into independent sets where G is an n -chromatic graph and color all vertices of S_i with color i . By assumption the complete graph on vertex set $V = \{1, 2, \dots, n\}$ can be decomposed into T_1, T_2, \dots, T_{n-1} . Let G_i be the subgraph of G induced by

$$\bigcup_{j \in V(T_i)} S_j.$$

The graph G_i is obviously $(i + 1)$ -chromatic since it has $i + 1$ color classes and a proper coloring of G_i with at most i colors could be obviously extended to a proper coloring of G with at most $n - 1$ colors, leading to a contradiction. Applying Lemma 1 to G_i , we find a copy F_i of T_i labeled exactly the same way as T_i is labeled in K_n . Repeating this for $i = 1, 2, \dots, n - 1$, we obtain edge disjoint copies of F_1, F_2, \dots, F_{n-1} in G , in fact they are not only edge disjoint but the union of their edge sets takes exactly one edge from each bipartite graph $\{[S_i, S_j] : 1 \leq i < j \leq n\}$. ■

Theorem 2 allows to transfer known tree-packing results from K_n to n -chromatic graphs. In particular, since any sequence of trees containing only paths and stars are known to be packable to K_n ([3, 5]), we get the following, conjectured in [2].

Corollary 3. Any sequence T_1, T_2, \dots, T_{n-1} of stars and paths is packable into any n -chromatic graph.

REFERENCES

- [1] D. Gerbner, B. Keszegh and C. Palmer, *Generalizations of the tree packing conjecture*, Discuss. Math. Graph Theory **32** (2012) 569–582.
doi:10.7151/dmgt.1628
- [2] D. Gerbner, B. Keszegh and C. Palmer, *Red-blue alternating paths*, Third Emléktábla Workshop, p.25.
<http://www.renyi.hu/emlektab/index/booklet.html>
- [3] A. Gyárfás and J. Lehel, *Packing trees of different order into K_n* , Combinatorics, Proc. Fifth Hungarian Coll. Keszthely, 1976, Vol II. North Holland. Colloq. Math. Soc. J. Bolyai **18** 463–469.
- [4] A. Gyárfás, E. Szemerédi and Zs. Tuza, *Induced subtrees in graphs of large chromatic number*, Discrete Math. **30** (1980) 235–244.
doi:10.1016/0012-365X(80)90230-7

- [5] S. Zaks and C.L. Liu, *Decomposition of graphs into trees*, Proceedings of 8-th Southeastern Conference on Combinatorics, Graph Theory and Computing, Louisiana State Univ., Baton Rouge, La. Util. Math., Congr. Numer. **XIX** (1977) 643–654.

Received 13 June 2013

Accepted 26 June 2013