

## ON TWIN EDGE COLORINGS OF GRAPHS

ERIC ANDREWS, LAARS HELENIUS

DANIEL JOHNSTON, JONATHON VERWYS

AND

PING ZHANG

*Department of Mathematics*  
*Western Michigan University*  
*Kalamazoo, MI 49008, USA*

**e-mail:** ping.zhang@wmich.edu

### Abstract

A twin edge  $k$ -coloring of a graph  $G$  is a proper edge coloring of  $G$  with the elements of  $\mathbb{Z}_k$  so that the induced vertex coloring in which the color of a vertex  $v$  in  $G$  is the sum (in  $\mathbb{Z}_k$ ) of the colors of the edges incident with  $v$  is a proper vertex coloring. The minimum  $k$  for which  $G$  has a twin edge  $k$ -coloring is called the twin chromatic index of  $G$ . Among the results presented are formulas for the twin chromatic index of each complete graph and each complete bipartite graph.

**Keywords:** edge coloring, vertex coloring, factorization.

**2010 Mathematics Subject Classification:** 05C15, 05C70.

### 1. INTRODUCTION

In 1968, Rosa [13] introduced a vertex labeling that induces an *edge-distinguishing labeling* defined by subtracting labels. In particular, for a graph  $G$  of size  $m$ , a vertex labeling (an injective function)  $f : V(G) \rightarrow \{0, 1, \dots, m\}$  was called a  $\beta$ -valuation by Rosa if the induced edge labeling  $f' : E(G) \rightarrow \{1, 2, \dots, m\}$  defined by  $f'(uv) = |f(u) - f(v)|$  was bijective. In 1972 Golomb [8] called a  $\beta$ -valuation a *graceful labeling* and a graph possessing a graceful labeling a *graceful graph*. It is this terminology that has become standard. Much research has been done on

graceful graphs. A popular conjecture in graph theory, due to Anton Kotzig and Gerhard Ringel, is the following.

**The Graceful Tree Conjecture.** *Every nontrivial tree is graceful.*

In 1991 Gnana Jothi [7] introduced a concept that, in a certain sense, reverses the roles of vertices and edges in graceful labelings (see also [6]). For a connected graph  $G$  of order  $n \geq 3$ , let  $f : E(G) \rightarrow \mathbb{Z}_n$  be an edge labeling of  $G$  that induces a bijective function  $f' : V(G) \rightarrow \mathbb{Z}_n$  defined by  $f'(v) = \sum_{e \in E_v} f(e)$  for each vertex  $v$  of  $G$ , where  $E_v$  is the set of edges of  $G$  incident with a vertex  $v$ . Such a labeling  $f$  is called a *modular edge-graceful labeling*, while a graph possessing such a labeling is called *modular edge-graceful* (see [10]). Verifying a conjecture by Gnana Jothi on trees, Jones, Kolasinski and Zhang [11] showed not only that every tree of order  $n \geq 3$  is modular edge-graceful if and only if  $n \not\equiv 2 \pmod{4}$  but a connected graph of order  $n \geq 3$  is modular edge-graceful if and only if  $n \not\equiv 2 \pmod{4}$ . These concepts have been studied in greater detail by Jones [9]. A generalization of this concept has been introduced recently by Anholcer, Cichacz and Milanič in [2].

Prior to Jothi's paper, an edge labeling (with positive integers) of a connected graph  $G$  was introduced in 1986 [3] for the purpose of producing a weighted graph whose degrees (obtained by adding the weights of the incident edges of each vertex) were distinct. Such a weighted graph was called *irregular*. This concept was later looked at in another manner. For the set  $\mathbb{N}$  of positive integers, an edge coloring  $c : E(G) \rightarrow \mathbb{N}$ , where adjacent edges may be colored the same, is said to be *vertex-distinguishing* if the coloring  $c' : V(G) \rightarrow \mathbb{N}$  induced by  $c$  and defined by  $c'(v) = \sum_{e \in E_v} c(e)$  has the property that  $c'(x) \neq c'(y)$  for every two distinct vertices  $x$  and  $y$  of  $G$ . The research in [3] dealt with minimizing the largest color assigned to the edges of the graph to produce an irregular graph. Vertex-distinguishing edge colorings have received increased attention during the past 25 years (see [5, pp. 370-385]).

A *neighbor-distinguishing coloring* of a graph  $G$  is a coloring in which every pair of adjacent vertices of  $G$  are colored differently. Such a coloring is more commonly called a *proper vertex coloring*. The minimum number of colors needed in a proper vertex coloring of a graph  $G$  is the chromatic number of  $G$  and denoted by  $\chi(G)$ . A number of neighbor-distinguishing vertex colorings different from standard proper colorings have been introduced in the literature (see [5, pp. 383-391], for example).

In 2005 non-proper edge colorings of graphs were studied that induce a proper vertex coloring [1]. In particular, for  $k \in \mathbb{N}$ , let  $c : E(G) \rightarrow \{1, 2, \dots, k\}$  be an edge coloring of  $G$  (where adjacent edges may be assigned the same color). A vertex coloring  $c' : V(G) \rightarrow \mathbb{N}$  is defined where  $c'(v)$  is the sum of the colors of the edges incident with  $v$ . If  $c'$  is a proper vertex coloring of  $G$ , then  $c$  is called a

*neighbor-distinguishing edge coloring* of  $G$  (see [5, p. 385]). A major conjecture in this area is the following [12].

**The 1-2-3 Conjecture.** *For every connected graph  $G$  of order at least 3, there exists a neighbor-distinguishing edge coloring of  $G$  using only the colors 1, 2, 3.*

Among the various edge colorings studied in graph theory, the best known and most studied are proper edge colorings. In a *proper edge coloring* of a graph  $G$ , each edge of  $G$  is assigned a color from a given set of colors where adjacent edges are colored differently. The minimum number of colors needed in a proper edge coloring of  $G$  is called the *chromatic index* of  $G$  and is denoted by  $\chi'(G)$ . The classic theorem in this connection is due to Vizing [14] who proved that  $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$  for every nonempty graph  $G$ .

A related and also well-studied graph coloring is the so-called *total coloring* of a graph  $G$  that assigns colors to both the vertices and edges of  $G$  so that not only the vertex coloring and edge coloring are proper but no vertex and an incident edge are assigned the same color. The minimum number of colors required for a total coloring of  $G$  is the *total chromatic number* of  $G$ , denoted by  $\chi''(G)$ . It then follows that  $\chi''(G) \geq \Delta(G) + 1$ , where  $\Delta(G)$  is the maximum degree of  $G$ . A well-known conjecture in this area is due independently to Behzad and Vizing (see [5, p. 282]).

**The Total Coloring Conjecture.** *For every graph  $G$ ,  $\chi''(G) \leq 2 + \Delta(G)$ .*

Inspired by the graph colorings described above, we introduce a proper edge coloring of a graph that induces a proper vertex coloring where the colors belong to  $\mathbb{Z}_k$  for some integer  $k \geq 2$ . We refer to the books [4, 5] for graph theory notation and terminology not described in this paper. All graphs under consideration here are connected graphs of order at least 3.

## 2. TWIN CHROMATIC INDEX

For a connected graph  $G$  of order at least 3, a proper edge coloring  $c : E(G) \rightarrow \mathbb{Z}_k$  for some integer  $k \geq 2$  is sought for which the induced vertex coloring  $c' : V(G) \rightarrow \mathbb{Z}_k$  defined by

$$c'(v) = \sum_{e \in E_v} c(e) \text{ in } \mathbb{Z}_k,$$

(where the indicated sum is computed in  $\mathbb{Z}_k$ ) results in a proper vertex coloring of  $G$ . We refer to such a coloring as a *twin edge  $k$ -coloring* or simply a *twin edge coloring* of  $G$ . The minimum  $k$  for which  $G$  has a twin edge  $k$ -coloring is called the *twin chromatic index* of  $G$  and is denoted by  $\chi'_t(G)$ . Since a twin edge coloring

is not only a proper edge coloring of  $G$  but induces a proper vertex coloring of  $G$ , it follows that

$$\chi'_t(G) \geq \max\{\chi(G), \chi'(G)\}.$$

Since  $\max\{\chi(G), \chi'(G)\} = \chi'(G)$  except when  $G$  is a complete graph of even order, we have  $\chi'_t(G) \geq \chi'(G)$  except possibly when  $G$  is a complete graph of even order.

While  $\chi'_t(G)$  does not exist if  $G$  is the connected graph of order 2, every connected graph of order at least 3 has a twin edge coloring. To see this, let  $G$  be a connected graph of size  $m \geq 2$ . If  $m = 2$ , then assign the colors 1 and 2 in  $\mathbb{Z}_3$  to the two edges of  $G$ . If  $m \geq 3$ , then assign the  $m$  elements  $0, 1, 2, 4, \dots, 2^{m-2} \in \mathbb{Z}_{2^{m-1}}$  to the  $m$  edges of  $G$  in a one-to-one manner so that the color 0 is assigned to a pendant edge if  $G$  has such an edge. Hence the sets of edges colored by nonzero elements in  $\mathbb{Z}_{2^{m-1}}$  that are incident with every two adjacent vertices are distinct. Since the base 2 representations of the colors of these vertices are different, it follows that adjacent vertices are assigned distinct colors in  $\mathbb{Z}_{2^{m-1}}$ . Thus, this coloring is a twin edge coloring. This observation yields the following.

**Proposition 2.1.** *If  $G$  is a connected graph of order at least 3 and size  $m$ , then  $\chi'_t(G)$  exists. Furthermore,  $\chi'_t(G) \leq 2^{m-1}$  if  $m \geq 3$ .*

To illustrate the concept of twin edge colorings, we determine the twin chromatic indexes of two familiar classes of graphs, namely paths and cycles. We begin with paths.

**Proposition 2.2.** *If  $P_n$  is a path of order  $n \geq 3$ , then  $\chi'_t(P_n) = 3$ .*

**Proof.** Let  $P_n = (v_1, v_2, \dots, v_n)$  be a path of order  $n \geq 3$  where  $e_i = v_i v_{i+1}$  for  $i = 1, 2, \dots, n-1$ . Since  $\chi'(P_n) = 2$ , it follows that  $\chi'_t(P_n) \geq \chi'(P_n) = 2$ . First, we show that  $\chi'_t(P_n) \neq 2$ . Let  $c$  be a proper edge coloring of  $P_n$  using the colors of  $\mathbb{Z}_2$ . Then  $c(e_i) = 1 \in \mathbb{Z}_2$  for some  $i \in \{1, 2, \dots, n-1\}$  and so  $c(e_{i-1}) = 0$  if  $i \geq 2$  and  $c(e_{i+1}) = 0$  if  $i \leq n-2$ . However then,  $c'(v_i) = c'(v_{i+1}) = 1$  and so  $c$  is not a twin edge 2-coloring. Thus, as claimed,  $\chi'_t(P_n) \geq 3$ . It remains to show that  $P_n$  has a twin edge 3-coloring. A coloring  $c : E(P_n) \rightarrow \mathbb{Z}_3$  is defined as follows.

• For  $n \equiv 0 \pmod{3}$  or  $n \equiv 1 \pmod{3}$ , let  $c(e_j) = r$  if  $j \equiv r \pmod{3}$  for  $r = 0, 1, 2$ . For example, if  $n = 6$ , then  $(c(e_1), c(e_2), \dots, c(e_5)) = (1, 2, 0, 1, 2)$ ; while if  $n = 7$ , then  $(c(e_1), c(e_2), \dots, c(e_6)) = (1, 2, 0, 1, 2, 0)$ . If  $n \equiv 0 \pmod{3}$ , then for  $1 \leq i \leq n$ ,

$$(1) \quad c'(v_i) = \begin{cases} 0 & \text{if } i \equiv 2 \pmod{3}, \\ 1 & \text{if } i \equiv 1 \pmod{3}, \\ 2 & \text{if } i \equiv 0 \pmod{3}. \end{cases}$$

If  $n \equiv 1 \pmod{3}$ , then  $c'(v_i)$  is given in (1) for  $1 \leq i \leq n - 1$  and  $c'(v_n) = 0$ . Hence  $(c'(v_1), c'(v_2), \dots, c'(v_6)) = (1, 0, 2, 1, 0, 2)$  and  $(c'(v_1), c'(v_2), \dots, c'(v_7)) = (1, 0, 2, 1, 0, 2, 0)$ .

• For  $n \equiv 2 \pmod{3}$ , let  $c(e_j) = 2 + r$  if  $j \equiv r \pmod{3}$  for  $r = 0, 1, 2$ . Then  $c'(v_1) = c'(v_n) = 0$  and for  $2 \leq i \leq n - 1$ ,

$$c'(v_i) = \begin{cases} 0 & \text{if } i \equiv 0 \pmod{3}, \\ 1 & \text{if } i \equiv 2 \pmod{3}, \\ 2 & \text{if } i \equiv 1 \pmod{3}. \end{cases}$$

For example, if  $n = 8$ , then  $(c(e_1), c(e_2), \dots, c(e_7)) = (0, 1, 2, 0, 1, 2, 0)$  and  $(c'(v_1), c'(v_2), \dots, c'(v_8)) = (0, 1, 0, 2, 1, 0, 2, 0)$ . Therefore,  $\chi'_t(P_n) \geq 3$  and so  $\chi'_t(P_n) = 3$  for  $n \geq 3$ . ■

To determine the twin chromatic indexes of cycles, the following observation will be useful.

**Observation 2.3.** *If a connected graph  $G$  contains two adjacent vertices of degree  $\Delta(G)$ , then  $\chi'_t(G) \geq 1 + \Delta(G)$ .*

**Proposition 2.4.** *If  $C_n$  is a cycle of order  $n \geq 3$ , then*

$$\chi'_t(C_n) = \begin{cases} 3 & \text{if } n \equiv 0 \pmod{3}, \\ 4 & \text{if } n \not\equiv 0 \pmod{3} \text{ and } n \neq 5, \\ 5 & \text{if } n = 5. \end{cases}$$

**Proof.** Let  $C_n = (v_1, v_2, \dots, v_n, v_{n+1} = v_1)$  where  $e_i = v_i v_{i+1}$  for  $i = 1, 2, \dots, n$  and  $e_{n+1} = e_1$ . By Observation 2.3,  $\chi'_t(C_n) \geq 3$ . First, suppose that  $n \equiv 0 \pmod{3}$  and so  $n = 3k$  for some positive integer  $k$ . Define the coloring  $c : E(C_n) \rightarrow \mathbb{Z}_3$  by  $c(e_i) \equiv 2 + r \pmod{3}$  if  $i \equiv r \pmod{3}$  for  $r = 0, 1, 2$ . Then for  $1 \leq i \leq n$ ,

$$c'(v_i) = \begin{cases} 0 & \text{if } i \equiv 0 \pmod{3}, \\ 1 & \text{if } i \equiv 2 \pmod{3}, \\ 2 & \text{if } i \equiv 1 \pmod{3}. \end{cases}$$

For example, if  $n = 6$ , then  $(c(e_1), c(e_2), \dots, c(e_6)) = (0, 1, 2, 0, 1, 2)$  and  $(c'(v_1), c'(v_2), \dots, c'(v_6)) = (2, 1, 0, 2, 1, 0)$ . Hence  $\chi'_t(C_n) = 3$  when  $n \equiv 0 \pmod{3}$ .

Next, suppose that  $n \not\equiv 0 \pmod{3}$  and  $n \neq 5$ . First, we make an observation, namely, if  $c$  is a twin edge coloring of  $C_n$  and  $|i - j| = 2$ , then  $c(e_i) \neq c(e_j)$ . Suppose, say, that  $c(e_1) = c(e_3)$ . However then,  $c'(v_2) = c(e_1) + c(e_2) = c(e_2) + c(e_3) = c'(v_3)$ , which is impossible. This implies that if  $n \not\equiv 0 \pmod{3}$ , then  $\chi'_t(C_n) \geq 4$ . To show that  $\chi'_t(C_n) \leq 4$ , define the coloring  $c : E(C_n) \rightarrow \mathbb{Z}_4$  as follows.

• For  $n \equiv 1 \pmod{3}$ , let  $c(e_i) \equiv 2 + r \pmod{3}$  if  $i \equiv r \pmod{3}$  for  $r = 0, 1, 2$  and  $1 \leq i \leq n-1$  and  $c(e_n) = 3$ . Then  $c'(v_1) = 3, c'(v_n) = 1$  and for  $2 \leq i \leq n-1$ ,

$$(2) \quad c'(v_i) = \begin{cases} 1 & \text{if } i \equiv 2 \pmod{3}, \\ 2 & \text{if } i \equiv 1 \pmod{3}, \\ 3 & \text{if } i \equiv 0 \pmod{3}. \end{cases}$$

(In particular,  $c'(v_2) = 1$  and  $c'(v_{n-1}) = 3$ .) For example, if  $n = 7$ , then  $(c(e_1), c(e_2), \dots, c(e_7)) = (0, 1, 2, 0, 1, 2, 3)$  and  $(c'(v_1), c'(v_2), \dots, c'(v_7)) = (3, 1, 3, 2, 1, 3, 1)$ . Hence  $\chi'_t(C_n) = 4$  when  $n \equiv 1 \pmod{3}$ .

• Let  $n \equiv 2 \pmod{3}$  and  $n \geq 8$ . If  $n = 8$ , let  $(c(e_1), c(e_2), \dots, c(e_8)) = (0, 1, 2, 3, 0, 1, 2, 3)$ ; while if  $n \geq 11$ , let  $c(e_i) \equiv 2 + r \pmod{3}$  if  $i \equiv r \pmod{3}$  for  $r = 0, 1, 2$  and  $1 \leq i \leq n-9$  and let  $(c(e_{n-8}), c(e_{n-7}), \dots, c(e_n)) = (0, 1, 2, 3, 0, 1, 2, 3)$ .

Consequently, if  $n = 8$ , then  $(c'(v_1), c'(v_2), \dots, c'(v_8)) = (3, 1, 3, 1, 3, 1, 3, 1)$ ; while if  $n \geq 11$ , then  $c'(v_1) = 3, c'(v_i)$  is the same as in (2) for  $2 \leq i \leq n-9$  and  $(c'(v_{n-8}), c'(v_{n-7}), \dots, c'(v_n)) = (3, 1, 3, 1, 3, 1, 3, 1)$ . For example, if  $n = 11$ , then  $(c(e_1), c(e_2), \dots, c(e_{11})) = (0, 1, 2, 0, 1, 2, 3, 0, 1, 2, 3)$  and  $(c'(v_1), c'(v_2), \dots, c'(v_{11})) = (3, 1, 3, 2, 1, 3, 1, 3, 1, 3, 1)$ . Hence  $\chi'_t(C_n) = 4$  when  $n \equiv 2 \pmod{3}$ .

Finally, we show that  $\chi'_t(C_5) = 5$ . We have already observed that  $\chi'_t(C_5) \geq 3$ . Let  $C_5 = (v_0, v_1, v_2, v_3, v_4, v_5 = v_0)$  and let  $c : E(C_5) \rightarrow \mathbb{Z}_5$  be defined by  $c(v_i v_{i+1}) = i$  for  $0 \leq i \leq 4$ . Since  $c'(v_0) = 4, c'(v_1) = 1, c'(v_2) = 3, c'(v_3) = 0$  and  $c'(v_4) = 2$ , it follows that  $c$  is a twin edge 5-coloring of  $C_5$  and so  $\chi'_t(C_5) \leq 5$ . We now show that  $\chi'_t(C_5) \geq 5$ . Suppose that there is a twin edge  $k$ -coloring where  $k = 3$  or  $k = 4$ . Then some element  $a \in \mathbb{Z}_k$  must be used twice, say  $c(v_0 v_1) = c(v_2 v_3) = a$ . Suppose that  $c(v_1 v_2) = b$ , where  $b \neq a$ . Then  $c'(v_1) = c'(v_2) = a + b$ , which is a contradiction. Thus,  $\chi'_t(C_5) = 5$ . ■

### 3. COMPLETE GRAPHS

We now investigate twin edge colorings of complete graphs  $K_n$  starting with the case  $n$  being odd. The following observation will be useful later.

**Observation 3.1.** *Let  $n \geq 2$  be an integer. If  $n$  is odd, then  $\binom{n}{2} = 0$  in  $\mathbb{Z}_n$  and if  $n$  is even, then  $\binom{n}{2} = \frac{n}{2}$  in  $\mathbb{Z}_n$ .*

**Lemma 3.2.** *If  $n \geq 3$  is an odd integer, then  $\chi'_t(K_n) = n$ .*

**Proof.** By Observation 2.3,  $\chi'_t(K_n) \geq 1 + \Delta(K_n) = n$ . To show that  $\chi'_t(K_n) \leq n$ , let  $V(K_n) = \{v_0, v_1, \dots, v_{n-1}\}$  and arrange the vertices  $v_0, v_1, \dots, v_{n-1}$  consecutively in a regular  $n$ -gon and join every two vertices by a straight line segment, producing  $K_n$ . For each  $i$  ( $0 \leq i \leq n-1$ ), assign to  $v_{i-1} v_{i+1}$  and those edges

parallel to  $v_{i-1}v_{i+1}$  the color  $i$ . Then  $v_i$  has the color  $\binom{n}{2} - i$ , resulting in a proper vertex coloring of  $K_n$ . Thus  $\chi'_t(K_n) = n$ . ■

When  $n \geq 4$  is even, however,  $\chi'_t(K_n) \neq n$ .

**Lemma 3.3.** *If  $n \geq 4$  is an even integer, then  $\chi'_t(K_n) \geq n + 1$ .*

**Proof.** Since  $\chi'_t(K_n) \geq 1 + \Delta(K_n) = n$  by Observation 2.3, it remains to show that  $\chi'_t(K_n) \neq n$ . Assume, to the contrary, that  $\chi'_t(K_n) = n$ . Then there is a proper edge coloring of  $K_n$  using the colors in  $\mathbb{Z}_n$  that results in a proper vertex coloring of  $K_n$ . Since every vertex of  $K_n$  has degree  $n - 1$ , the edges incident with each vertex of  $K_n$  are colored with an  $(n - 1)$ -element subset of  $\mathbb{Z}_n$ . For example, if  $v$  is a vertex of  $K_n$ , then there is exactly one element  $a \in \mathbb{Z}_n$  that is not used in coloring the edges incident with  $v$ . Consequently, at most  $\frac{n}{2} - 1$  edges of  $K_n$  are colored  $a$ , implying that there exists some other vertex  $u$  of  $K_n$  none of whose incident edges are colored  $a$ . However then,  $c'(u) = c'(v) = \binom{n}{2} - a$ , which is impossible since  $u$  and  $v$  are adjacent in  $K_n$ . Thus  $\chi'_t(K_n) \geq n + 1$ . ■

If  $n \geq 4$  is an even integer, then either  $n \equiv 0 \pmod{4}$  or  $n \equiv 2 \pmod{4}$ . We consider these two situations, beginning with  $n \equiv 0 \pmod{4}$ .

**Lemma 3.4.** *If  $n \geq 4$  is an integer with  $n \equiv 0 \pmod{4}$ , then  $\chi'_t(K_n) = n + 1$ .*

**Proof.** By Lemma 3.3, it suffices to show that  $K_n$  has a twin edge  $(n + 1)$ -coloring. Let  $V(K_n) = \{v_0, v_1, \dots, v_{n-1}\}$  and arrange the vertices  $v_0, v_1, \dots, v_{n-1}$  consecutively in a regular  $n$ -gon and join every two vertices by a straight line segment, thereby producing  $K_n$ .

Since  $n \equiv 0 \pmod{4}$  and  $n \geq 4$ , it follows that  $n = 4k$  for some positive integer  $k$ . For  $k = 1$ , the coloring  $c : E(K_4) \rightarrow \mathbb{Z}_5$  defined by  $c(v_0v_1) = c(v_2v_3) = 0$ ,  $c(v_0v_2) = 1$ ,  $c(v_0v_3) = 2$ ,  $c(v_1v_2) = 3$  and  $c(v_1v_3) = 4$  is a twin edge 5-coloring of  $K_4$  and so we may assume that  $k \geq 2$ . First, let  $M_0, M_1, \dots, M_{2k-1}$  be  $2k$  pairwise edge-disjoint matchings of size  $2k - 1$  in  $K_{4k}$  where each matching  $M_i$  ( $0 \leq i \leq 2k - 1$ ) consists of those  $2k - 1$  edges perpendicular to  $v_i v_{2k+i}$ . Then  $H = K_{4k} - \left(\bigcup_{i=0}^{2k-1} M_i\right)$  is therefore a  $(2k)$ -regular graph. The graph  $H$  has a 1-factorization  $\{F_1, F_2, \dots, F_{2k}\}$  where  $F_i$  ( $1 \leq i \leq 2k$ ) consists of the edge  $v_i v_{i+1}$  and those edges parallel to  $v_i v_{i+1}$ . Let  $X_1 = \{v_0 v_{2k-1}, v_1 v_{2k-2}, \dots, v_{k-1} v_k\}$  and  $X'_1 = \{v_{2k} v_{4k-1}, v_{2k+1} v_{4k-2}, \dots, v_{3k-1} v_{3k}\}$ . Thus  $|X_1| = |X'_1| = k$  and  $E(F_{k-1}) = X_1 \cup X'_1$ . Define a coloring  $c : E(K_{4k}) \rightarrow \mathbb{Z}_{4k+1}$  as follows. If  $k = 2$ , let

$$c(e) = \begin{cases} 0 & \text{if } e \in X'_1, \\ i - 1 & \text{if } e \in E(F_i) \text{ where } 2 \leq i \leq 2k, \\ 2k & \text{if } e \in X_1, \\ 2k + j + 1 & \text{if } e \in M_j \text{ where } 0 \leq j \leq 2k - 1. \end{cases}$$

If  $k \geq 3$ , let

$$c(e) = \begin{cases} 0 & \text{if } e \in X'_1, \\ i & \text{if } e \in E(F_i) \text{ where } 1 \leq i \leq k-2, \\ i-1 & \text{if } e \in E(F_i) \text{ where } k \leq i \leq 2k, \\ 2k & \text{if } e \in X_1, \\ 2k+j+1 & \text{if } e \in M_j \text{ where } 0 \leq j \leq 2k-1. \end{cases}$$

Then  $c$  is a proper edge coloring. For  $0 \leq i \leq 2k-1$ ,

$$c'(v_i) = \left[ \binom{4k+1}{2} - 2k \right] - (2k+i+1) + 2k = -(2k+i+1) \text{ in } \mathbb{Z}_{4k+1};$$

while for  $2k \leq i \leq 4k-1$ ,

$$c'(v_i) = \left[ \binom{4k+1}{2} - 2k \right] - (i+1) + 0 = -(2k+i+1) \text{ in } \mathbb{Z}_{4k+1}.$$

Thus  $(c'(v_0), c'(v_1), \dots, c'(v_{4k-1})) = (2k, 2k-1, \dots, 1, 0, 4k, 4k-1, \dots, 2k+2)$ . That is, each color in  $\mathbb{Z}_{4k+1}$  (except  $2k+1$ ) is used exactly once. Therefore,  $c' : V(K_{4k}) \rightarrow \mathbb{Z}_{4k+1}$  is a proper vertex coloring of  $G$  and so  $\chi'_t(K_n) = n+1$ . ■

**Lemma 3.5.** *If  $n \geq 6$  is an integer with  $n \equiv 2 \pmod{4}$ , then  $\chi'_t(K_n) = n+1$ .*

**Proof.** Since  $\chi'_t(K_n) \geq n+1$  by Lemma 3.3, it suffices to show that  $K_n$  has a twin edge  $(n+1)$ -coloring when  $n \geq 6$  with  $n \equiv 2 \pmod{4}$ . Let  $n = 4k+2$  for some positive integer  $k$  and let  $V(K_{4k+2}) = \{v_0, v_1, \dots, v_{4k+1}\}$ . Arrange the vertices  $v_1, v_2, \dots, v_{4k+1}$  consecutively in a regular  $(4k+1)$ -gon, place  $v_0$  in the center of the  $(4k+1)$ -gon and then join every two vertices by a straight line segment, thereby producing  $K_{4k+2}$ .

Let  $\mathcal{F} = \{F_1, F_2, \dots, F_{4k+1}\}$  be the 1-factorization of  $K_{4k+2}$ , in which  $F_i$  is the 1-factor of  $K_{4k+2}$  that consists of the edge  $v_0v_{2k+1+i}$  and the  $2k$  edges perpendicular to  $v_0v_{2k+1+i}$  when  $1 \leq i \leq 2k$  and  $F_i$  consists of the edge  $v_0v_{i-2k}$  and the  $2k$  edges perpendicular to  $v_0v_{i-2k}$  where  $2k+1 \leq i \leq 4k+1$ . Also, let  $M_i = E(F_i)$  ( $1 \leq i \leq 4k+1$ ) denote the perfect matching of  $K_{4k+2}$  resulting from  $F_i$ . Observe that the edge  $v_iv_{i+1}$  belongs to  $M_i$  for  $1 \leq i \leq 4k$  and  $v_{4k+1}v_1 \in M_{4k+1}$ .

We now define an edge coloring  $c_1$  (described below) that assigns the  $4k+1$  colors in  $\mathbb{Z}_{4k+3} - \{0, 1\}$  to the  $4k+1$  matchings  $M_1, M_2, \dots, M_{4k+1}$  such that

- (i)  $c_1$  assigns exactly one color to all edges in  $M_i$  for each  $i$  ( $1 \leq i \leq 4k+1$ ) and
- (ii)  $c_1(e) \neq c_1(f)$  if  $e \in M_i, f \in M_j$  where  $i \neq j$ .



- For an even integer  $i$  with  $2 \leq i \leq 4k$ , let

$$c_1(e) = \begin{cases} (2k + 3) - i & \text{if } e \in M_i \text{ and } 2 \leq i \leq 2k, \\ i - 2k & \text{if } e \in M_i \text{ and } 2k + 2 \leq i \leq 4k. \end{cases}$$

- For  $i = 1$  or  $i = 2k + 1$ , let

$$c_1(e) = \begin{cases} 2k + 3 & \text{if } e \in M_1, \\ 2k + 2 & \text{if } e \in M_{2k+1}. \end{cases}$$

• For the remaining  $2k - 1$  matchings  $M_3, M_5, \dots, M_{2k-1}$  and  $M_{2k+3}, M_{2k+5}, \dots, M_{4k+1}$ , the coloring  $c_1$  assigns the remaining  $2k - 1$  colors  $2k + 4, 2k + 5, \dots, 4k + 2$  to these matchings in an arbitrary way such that distinct colors are assigned to the edges in distinct matchings.

Hence, the  $2k$  colors  $2, 3, \dots, 2k + 1$  in  $\mathbb{Z}_{4k+3}$  are used to color the edges in the  $2k$  matchings  $M_2, M_4, \dots, M_{4k}$ ; while the  $2k + 1$  colors  $2k + 2, 2k + 3, \dots, 4k + 2$  in  $\mathbb{Z}_{4k+3}$  are used to color the edges in  $2k + 1$  matchings  $M_1, M_3, \dots, M_{4k+1}$ . Therefore,  $c_1$  is a proper edge coloring of  $K_{4k+2}$ . Since the colors 0 and 1 are not used,

$$c'_1(v) = 2 + 3 + \dots + (4k + 2) = \binom{4k+3}{2} - 0 - 1$$

for each  $v \in V(K_{4k+2})$ .

Next, we define a new edge coloring  $c : E(K_{4k+2}) \rightarrow \mathbb{Z}_{4k+3}$  from the coloring  $c_1$  as follows. First, we partition  $M_{2k+1}$  into two sets  $X$  and  $Y$  where

$$\begin{aligned} X &= \{v_i v_{4k+3-i} : i \text{ is odd and } 3 \leq i \leq 2k + 1\}, \\ Y &= \{v_0 v_1\} \cup \{v_i v_{4k+3-i} : i \text{ is even and } 2 \leq i \leq 2k\}. \end{aligned}$$

For each  $e \in E(K_{4k+2})$ , let

$$(3) \quad c(e) = \begin{cases} 0 & \text{if } e \in \{v_i v_{i+1} : i \text{ is even and } 0 \leq i \leq 4k\}, \\ 1 & \text{if } e \in \{v_1 v_2\} \cup X, \\ c_1(e) & \text{otherwise.} \end{cases}$$

Let  $b = \binom{4k+3}{2} - 1$ , where then  $b = -1$  in  $\mathbb{Z}_{4k+3}$  and let  $c' : V(K_{4k+2}) \rightarrow \mathbb{Z}_{4k+3}$  be the vertex coloring induced by  $c$ . Then

- For  $i = 0, 1, 2$ ,

$$\begin{aligned} c'(v_0) &= b - (2k + 2), \\ c'(v_1) &= b - (2k + 2) - (2k + 3) + 1 = b - 1, \\ c'(v_2) &= b - (2k + 3) - (2k + 1) + 1 = b - 0. \end{aligned}$$

- For  $3 \leq i \leq 2k + 1$ ,

$$c'(v_i) = \begin{cases} b - (2k + 3 - i) & \text{if } i \text{ is even,} \\ b - (2k + 2) - (2k + 4 - i) + 1 = b - (2 - i) & \text{if } i \text{ is odd.} \end{cases}$$

- For  $2k + 2 \leq i \leq 4k + 1$ ,

$$c'(v_i) = \begin{cases} b - (2k + 2) - (i - 2k) + 1 = b - (i + 1) & \text{if } i \text{ is even,} \\ b - (i - 1 - 2k) & \text{if } i \text{ is odd.} \end{cases}$$

For each  $i$  with  $0 \leq i \leq 4k + 1$ , let  $c'(v_i) = b - a_i$  for  $0 \leq i \leq 4k + 1$ . If  $s_{c'} = (a_0, a_1, \dots, a_{4k+1})$  (where  $a_i = b - c'(v_i)$  for  $0 \leq i \leq 4k + 1$ ), then

$$\begin{aligned} s_{c'} = & (2k + 2, 1, 0, 4k + 2, 2k - 1, 4k, 2k - 3, \dots, 2k + 8, 5, \\ & 2k + 6, 3, 2k + 4 = b - c'(v_{2k+1}), 2k + 3, 2, \\ & 2k + 5, 4, \dots, 4k - 1, 2k - 2, 4k + 1, 2k). \end{aligned}$$

For example, the sequences  $s_{c'}$  for  $n = 6, 10, 14$  are the following:

$$\begin{aligned} & (4, 1, 0, 6, 5, 2) \text{ for } n = 6 \text{ and } k = 1, \\ & (6, 1, 0, 10, 3, 8, 7, 2, 9, 4) \text{ for } n = 10 \text{ and } k = 2, \\ & (8, 1, 0, 14, 5, 12, 3, 10, 9, 2, 11, 4, 13, 6) \text{ for } n = 14 \text{ and } k = 3. \end{aligned}$$

In conclusion, we observe that  $\{c'(v) : v \in V(K_{4k+2})\} = \{b - i : 0 \leq i \leq 4k + 2, i \neq 2k + 1\}$  and so  $c$  is a twin edge  $(4k + 3)$ -coloring of  $K_{4k+2}$ . ■

In summary, we have the following.

**Theorem 3.6.** *For each integer  $n \geq 3$ ,*

$$\chi'_t(K_n) = \begin{cases} n & \text{if } n \text{ is odd,} \\ n + 1 & \text{if } n \text{ is even.} \end{cases}$$

#### 4. COMPLETE BIPARTITE GRAPHS

In this section we determine the twin chromatic indexes of the complete bipartite graphs  $K_{a,b}$  where  $1 \leq a \leq b$ , beginning with stars. For a star  $K_{1,b}$  ( $b \geq 2$ ), a twin edge coloring is the same as a modular edge-graceful labeling (see [9]) and so we have the following result (see also Lemma 2 in [2]).

**Proposition 4.1.** *If  $K_{1,b}$  is a star of order  $b \geq 2$ , then*

$$\chi'_t(K_{1,b}) = \begin{cases} b + 1 & \text{if } b \not\equiv 1 \pmod{4}, \\ b + 2 & \text{if } b \equiv 1 \pmod{4}. \end{cases}$$

We now determine  $\chi'_t(K_{a,b})$  where  $2 \leq a \leq b$  and  $b \in \{a, a + 1\}$ .

**Lemma 4.2.** *If  $a \geq 2$  and  $b$  are integers with  $b \in \{a, a + 1\}$ , then  $\chi'_t(K_{a,b}) = a + 2$ .*

**Proof.** Let  $U = \{u_1, u_2, \dots, u_a\}$  and  $V = \{v_1, v_2, \dots, v_b\}$  be the partite sets of  $K_{a,b}$ . We consider two cases, according to whether  $b = a$  or  $b = a + 1$ .

*Case 1.  $b = a$ .* By Observation 2.3,  $\chi'_t(K_{a,a}) \geq a + 1$ . First, we show that  $\chi'_t(K_{a,a}) \neq a + 1$ . Suppose that  $\chi'_t(K_{a,a}) = a + 1$ . Then there is a twin edge  $(a + 1)$ -coloring  $c$  of  $K_{a,a}$  using the colors in  $\mathbb{Z}_{a+1}$ . Hence  $c$  assigns exactly  $a$  colors to the  $a$  incident edges of each vertex of  $K_{a,a}$ . Consider  $u_1$  and let  $t \in \mathbb{Z}_{a+1}$  such that  $c$  assigns the colors in  $\mathbb{Z}_{a+1} - \{t\}$  to the edges incident with  $u_1$  (and so no edge incident with  $u_1$  is colored  $t$ ). We claim that for each vertex  $v_j$  ( $1 \leq j \leq a$ ), there is an edge incident with  $v_j$  that is colored  $t$ ; for otherwise, we may assume that no edge incident with  $v_1$  is colored  $t$ . However then,  $c$  assigns the colors in  $\mathbb{Z}_{a+1} - \{t\}$  to the edges incident with  $v_1$  and so  $c'(u_1) = c'(v_1)$ , which is impossible. Thus, as claimed, there is an edge incident with  $v_j$  that is colored  $t$  for  $j = 1, 2, \dots, a$ . Hence there are at least  $a$  edges of  $K_{a,a}$  that are colored  $t$ . Since no edge incident with  $u_1$  is colored  $t$ , it follows that at least two edges colored  $t$  are incident with the same vertex in  $V$ , which is a contradiction. Therefore,  $\chi'_t(K_{a,a}) \neq a + 1$  and so  $\chi'_t(K_{a,a}) \geq a + 2$ .

Next, we show that  $K_{a,a}$  has a twin edge  $(a + 2)$ -coloring. Since  $K_{a,a}$  is bipartite and  $a$ -regular,  $K_{a,a}$  is 1-factorable. Let  $\{F_0, F_1, \dots, F_{a-1}\}$  be a 1-factorization of  $K_{a,a}$  where

$$E(F_i) = \{u_j v_{j+i} : 1 \leq j \leq a\} \text{ for } 0 \leq i \leq a - 1$$

(all subscripts are expressed as integers modulo  $a$ ). For example,  $E(F_0) = \{u_j v_j : 1 \leq j \leq a\}$ ,  $E(F_1) = \{u_j v_{j+1} : 1 \leq j \leq a\}$  and  $E(F_{a-1}) = \{u_j v_{j+(a-1)} : 1 \leq j \leq a\}$ . We consider two cases, according to whether  $a$  is odd or  $a$  is even.

*Subcase 1.1  $a$  is odd.* Then  $a = 2k + 1$  for some positive integer  $k$ . Let  $M_a$  and  $M_{a+1}$  be the following matchings in  $K_{a,a}$ :

$$M_a = \{u_1 v_1, u_3 v_2, u_4 v_4, u_6 v_6, \dots, u_{2k} v_{2k}\},$$

$$M_{a+1} = \{u_1 v_2, u_3 v_3, u_4 v_5, u_6 v_7, \dots, u_{2k} v_{2k+1}\}.$$

Thus  $|M_a| = |M_{a+1}| = k + 1$ . For each  $i$  with  $0 \leq i \leq a - 1$ , let  $M_i = E(F_i) - (M_a \cup M_{a+1})$ . Define a proper edge coloring  $c : E(K_{a,a}) \rightarrow \mathbb{Z}_{a+2}$  by  $c(e) = i$  if  $e \in M_i$  for  $0 \leq i \leq a + 1$ . Since  $c'(u) = \binom{a}{2}$  or  $c'(u) = \binom{a}{2} - 4$  if  $u \in U$  and  $c'(v) = \binom{a}{2} - 1$  or  $c'(v) = \binom{a}{2} - 2$  if  $v \in V$ , it follows that  $c$  is a twin edge  $(a + 2)$ -coloring. Therefore,  $\chi'_t(K_{a,a}) = a + 2$ .

*Subcase 1.2  $a$  is even.* Then  $a = 2k \geq 2$  for some positive integer  $k$ . Let  $M_a$  and  $M_{a+1}$  be the following matchings in  $K_{a,a}$ :

$$M_a = \{u_1 v_1, u_3 v_3, \dots, u_{2k-1} v_{2k-1}\},$$

$$M_{a+1} = \{u_1 v_2, u_3 v_4, \dots, u_{2k-1} v_{2k}\}.$$

Thus  $|M_a| = |M_{a+1}| = k$ . For each  $i$  with  $0 \leq i \leq a - 1$ , let  $M_i = E(F_i) - (M_a \cup M_{a+1})$ . Define a proper edge coloring  $c : E(K_{a,a}) \rightarrow \mathbb{Z}_{a+2}$  by  $c(e) = i$  if  $e \in M_i$  for  $0 \leq i \leq a + 1$ . Since  $c'(u) = \binom{a}{2}$  or  $c'(u) = \binom{a}{2} - 4$  if  $u \in U$  and  $c'(v) = \binom{a}{2} - 2$  if  $v \in V$ , it follows that  $c$  is a twin edge  $(a+2)$ -coloring. Therefore,  $\chi'_t(K_{a,a}) = a + 2$ .

*Case 2.*  $b = a + 1$ . Since  $\Delta(K_{a,a+1}) = a + 1$ , it follows that  $\chi'_t(K_{a,a+1}) \geq a + 1$ . First, we show that  $\chi'_t(K_{a,a+1}) \neq a + 1$ . Suppose that  $\chi'_t(K_{a,a+1}) = a + 1$ . Then there is a twin edge  $(a + 1)$ -coloring  $c$  of  $K_{a,a+1}$  using colors in  $\mathbb{Z}_{a+1}$ . Since  $\deg u_i = a + 1$  for  $1 \leq i \leq a$ , it follows that  $c$  assigns all colors in  $\mathbb{Z}_{a+1}$  to the  $a + 1$  edges incident with each vertex  $u_i$ . Thus,  $a$  edges in  $K_{a,a+1}$  are colored 0. Since  $|V| = a + 1$ , some vertex in  $V$  is not incident with any edge colored 0, say  $v_1$ . Consequently,  $c$  assigns the  $a$  colors in  $\mathbb{Z}_{a+1} - \{0\}$  to the  $a$  edges incident with  $v_1$ . However then,  $c'(v_1) = c'(u_i) = \binom{a+1}{2}$  for  $1 \leq i \leq a$ , which is impossible. Therefore,  $\chi'_t(K_{a,a+1}) \neq a + 1$  and  $\chi'_t(K_{a,a+1}) \geq a + 2$ .

Next, we show that  $K_{a,a+1}$  has a twin edge  $(a + 2)$ -coloring. Define a proper edge coloring  $c : E(K_{a,a+1}) \rightarrow \mathbb{Z}_{a+2}$  using only the colors in  $\mathbb{Z}_{a+2} - \{0\}$  as follows. For each  $i$  with  $1 \leq i \leq a$ , let  $c(u_i v_{i+j}) = j + 1$  for each  $j$  with  $0 \leq j \leq a$ . In particular,  $c(u_i v_i) = 1$  for  $1 \leq i \leq a$ . Thus,  $c'(u_i) = \binom{a+2}{2}$  for  $1 \leq i \leq a$ . Furthermore,  $c'(v_j) = \binom{a+2}{2} - (j + 1)$  for  $1 \leq j \leq a$  and  $c'(v_{a+1}) = \binom{a+2}{2} - 1$ . Since  $c'(v_j) \neq \binom{a+2}{2}$  in  $\mathbb{Z}_{a+2}$  for  $1 \leq j \leq a + 1$ , it follows that  $c'$  is a proper vertex coloring of  $K_{a,a+1}$ . Therefore,  $\chi'_t(K_{a,a+1}) = a + 2$ . ■

Finally, we determine  $\chi'_t(K_{a,b})$  for all integers  $a$  and  $b$  with  $a \geq 2$  and  $b \geq a + 2$ .

**Lemma 4.3.** *If  $a \geq 2$  and  $b$  are integers with  $b \geq a + 2$ , then  $\chi'_t(K_{a,b}) = b$ .*

**Proof.** Since  $\chi'_t(K_{a,b}) \geq \chi'(K_{a,b}) = \Delta(K_{a,b}) = b$ , it suffices to show that  $K_{a,b}$  has a twin edge  $b$ -coloring. Let  $U = \{u_1, u_2, \dots, u_a\}$  and  $V = \{v_1, v_2, \dots, v_b\}$  be the partite sets of  $K_{a,b}$ . Suppose that

$$\sum_{i=0}^{a-1} i = \binom{a}{2} \equiv k \pmod{b} \text{ and } \sum_{i=0}^{b-1} i = \binom{b}{2} \equiv \ell \pmod{b},$$

where  $0 \leq k, \ell \leq b - 1$ . We consider two cases, depending on whether  $a$  and  $b$  are relatively prime.

*Case 1.*  $a$  and  $b$  are not relatively prime. Then  $d = \gcd(a, b) \geq 2$  and  $b = pd$  for some  $p \in \mathbb{N}$ . For  $0 \leq i \leq d - 1$ , let  $X_i = \{i, i + a, i + 2a, \dots, i + (p - 1)a\}$  be a subset of  $\mathbb{Z}_b$ . In fact,  $X_0, X_1, \dots, X_{d-1}$  are the cosets of the subgroup  $X_0 = \{0, a, 2a, \dots, (p - 1)a\}$  in the group  $\mathbb{Z}_b$ . Hence  $\mathcal{X} = \{X_0, X_1, \dots, X_{d-1}\}$  is a partition of  $\mathbb{Z}_b$ . Next, let  $X, X' \in \mathcal{X}$  such that  $k \in X$  and  $\ell \in X'$ . We define a coloring  $c : E(K_{a,b}) \rightarrow \mathbb{Z}_b$ , according to whether  $X \neq X'$  or  $X = X'$ .

*Subcase 1.1.*  $X \neq X'$ . For  $1 \leq i \leq a$  and  $1 \leq j \leq b$ , define  $c(u_i v_j) = i + j - 2 \in \mathbb{Z}_b$ . Then  $c'(u_i) = \ell \in X'$  for  $1 \leq i \leq a$  and  $c'(v_j) = k + (j - 1)a \in X$  for  $1 \leq j \leq b$ . Since  $X \cap X' = \emptyset$ , it follows that  $c'(u_i) \neq c'(v_j)$  for all  $i, j$  with  $1 \leq i \leq a$  and  $1 \leq j \leq b$ . Thus  $c'$  is a proper vertex coloring.

*Subcase 1.2.*  $X = X'$ . Since  $d \geq 2$ , it follows that  $k + 1 \notin X$ , say  $k + 1 \in X'' \in \mathcal{X}$ . For  $1 \leq i \leq a - 1$  and  $1 \leq j \leq b$ , define  $c(u_i v_j) = i + j - 2 \in \mathbb{Z}_b$ . Furthermore, define  $c(u_a v_j) = a + j - 1 \in \mathbb{Z}_b$  for  $1 \leq j \leq b$ . Then  $c'(u_i) = \ell \in X$  for  $1 \leq i \leq a$  and  $c'(v_j) = (k + 1) + (j - 1)a \in X''$  for  $1 \leq j \leq b$ . Since  $X \cap X'' = \emptyset$ , it follows that  $c'(u_i) \neq c'(v_j)$  for all  $i, j$  with  $1 \leq i \leq a$  and  $1 \leq j \leq b$ . Thus  $c'$  is a proper vertex coloring.

*Case 2.*  $a$  and  $b$  are relatively prime. Note that  $\mathbb{Z}_b = \{\ell, \ell + a, \dots, \ell + (b - 1)a\}$ . We start with a proper edge coloring  $c_1 : E(K_{a,b}) \rightarrow \mathbb{Z}_b$  defined by  $c_1(u_i v_j) = i + j - 2$  for  $1 \leq i \leq a$  and  $1 \leq j \leq b$ . Then  $c'_1(u_i) = \ell$  for  $1 \leq i \leq a$  and  $c'_1(v_j) = k + (j - 1)a \in X$  for  $1 \leq j \leq b$ . Since  $a$  and  $b$  are relatively prime,  $\{c'_1(v_j) : 1 \leq j \leq b\} = \mathbb{Z}_b$ . Therefore, there exists exactly one integer  $t$  with  $1 \leq t \leq b$  such that  $c'_1(v_t) = \ell$ . Thus  $c'_1$  is not a proper vertex coloring. We now produce a twin edge  $b$ -coloring  $c$  from  $c_1$  as follows. Let  $r = \lceil a/2 \rceil$  and  $s = t + \lfloor (b - 1)/2 \rfloor$  in  $\mathbb{Z}_b$ , where then  $1 \leq s \leq b$  and  $s \neq t$ , and let  $c$  be the coloring obtained from  $c_1$  by interchanging the colors of the edges  $u_r v_t$  and  $u_r v_s$  in  $c_1$ ; that is,

$$c(e) = \begin{cases} c_1(e) & \text{if } e \in E(K_{a,b}) - \{u_r v_t, u_r v_s\}, \\ c_1(u_r v_s) & \text{if } e = u_r v_t, \\ c_1(u_r v_t) & \text{if } e = u_r v_s. \end{cases}$$

We show that  $c'(u_i) = \ell$  for  $1 \leq i \leq a$  and  $c'(v_j) \neq \ell$  for  $1 \leq j \leq b$ .

By the defining property of  $c$ , it follows that  $c'(u_i) = c'_1(u_i) = \ell$  and  $c'(v_j) = c'_1(v_j) \neq \ell$  for  $1 \leq j \leq b$  and  $j \neq s, t$ . Thus, it remains to show that  $c'(v_t) \neq \ell$  and  $c'(v_s) \neq \ell$ . Since  $\ell = k + (t - 1)a$  and  $s = t + \lfloor (b - 1)/2 \rfloor$ , it follows that

$$\begin{aligned} c'(v_t) &= c'_1(v_t) - c_1(u_r v_t) + c_1(u_r v_s) = \ell - (r + t - 2) + (r + s - 2) \\ &= \ell - t + s = \ell - t + [t + \lfloor (b - 1)/2 \rfloor] = \ell + \lfloor (b - 1)/2 \rfloor \\ c'(v_s) &= c'_1(v_s) - c_1(u_r v_s) + c_1(u_r v_t) = [k + (s - 1)a] - (r + s - 2) + (r + t - 2) \\ &= [k + (s - 1)a] - s + t = [k + (s - 1)a] - \lfloor (b - 1)/2 \rfloor \\ &= k + (t + \lfloor (b - 1)/2 \rfloor - 1)a - \lfloor (b - 1)/2 \rfloor = \ell + (a - 1)\lfloor (b - 1)/2 \rfloor. \end{aligned}$$

We consider two cases, according to whether  $b$  is odd or  $b$  is even.

*Subcase 2.1.*  $b$  is odd. Then  $\lfloor (b - 1)/2 \rfloor = \frac{b-1}{2}$ . We claim that

$$(4) \quad c'(v_t) = \ell + \frac{b-1}{2} \neq \ell \text{ in } \mathbb{Z}_b,$$

$$(5) \quad c'(v_s) = \ell + (a-1)\frac{b-1}{2} \neq \ell \text{ in } \mathbb{Z}_b.$$

Since  $b$  is odd,  $\ell = 0$  in  $\mathbb{Z}_b$  by Observation 3.1, while  $\frac{b-1}{2} \neq 0$  in  $\mathbb{Z}_b$ , which implies that (4) holds. To verify (5), we show that  $\frac{b-1}{2}(a-1) \not\equiv 0 \pmod{b}$ . If this were not the case, then  $\frac{b-1}{2}(a-1) = bx$  for some integer  $x$ . This implies that  $2bx = (b-1)(a-1) = a-1$  in  $\mathbb{Z}_b$  or  $a-1 \equiv 0 \pmod{b}$ . However then,  $b \mid (a-1)$ , which is impossible.

*Subcase 2.2.*  $b$  is even. Then  $\lfloor (b-1)/2 \rfloor = \frac{b}{2} - 1$  and  $\ell = \frac{b}{2}$  in  $\mathbb{Z}_b$  by Observation 3.1. Since  $a$  and  $b$  are relatively prime, it follows that  $a \geq 3$  is odd and so  $b \geq a + 3 \geq 6$ . We claim that

$$(6) \quad c'(v_t) = \ell + \left(\frac{b}{2} - 1\right) = b - 1 \neq \ell \text{ in } \mathbb{Z}_b,$$

$$(7) \quad c'(v_s) = \ell + \left(\frac{b}{2} - 1\right)(a-1) \neq \ell \text{ in } \mathbb{Z}_b.$$

Since  $\ell = \frac{b}{2}$  in  $\mathbb{Z}_b$  and  $b-1 \neq \frac{b}{2}$  in  $\mathbb{Z}_b$ , it follows that (6) holds. To verify (7), we show that  $(\frac{b}{2} - 1)(a-1) \not\equiv 0 \pmod{b}$ . If this were not the case, then  $\frac{b-2}{2}(a-1) = bx$  for some positive integer  $x$ . Since  $b$  is even,  $b = 2y$  for some integer  $y \geq 3$ . Then  $a = 2\frac{xy}{y-1} + 1$ . Since  $a$  is an integer and  $y \geq 3$ , it follows that  $(y-1) \nmid y$  and so  $(y-1) \mid x$ . Let  $x = (y-1)z$  for some positive integer  $z$ . However then,  $a = 2yz + 1 = bz + 1$ , which is impossible.

Thus  $c'$  is a proper vertex coloring of  $K_{a,b}$  and so  $\chi'_t(K_{a,b}) = b$ . ■

In summary, we have the following.

**Theorem 4.4.** *For positive integers  $a$  and  $b$  with  $a \leq b$ ,*

$$\chi'_t(K_{a,b}) = \begin{cases} b & \text{if } b \geq a + 2 \text{ and } a \geq 2, \\ b + 1 & \text{if either } a = 1 \text{ and } b \not\equiv 1 \pmod{4} \text{ or } b = a + 1 \geq 3, \\ b + 2 & \text{if either } a = 1 \text{ and } b \equiv 1 \pmod{4} \text{ or } b = a \geq 2. \end{cases}$$

For every connected graph  $G$  for which the twin chromatic index has been determined, we have seen that  $\chi'_t(G) = \Delta(G) + i$  for some  $i \in \{0, 1, 2, 3\}$ . This leads us to conclude this paper by stating the following problem.

**Problem 4.5.** *Is  $\chi'_t(G) \leq \Delta(G) + 3$  for every connected graph  $G$  of order at least 3?*

### Acknowledgments

We are grateful to Professor Gary Chartrand for suggesting the concept of twin edge colorings to us and kindly providing useful information on this topic. Furthermore, we thank the referees whose valuable suggestions resulted in an improved paper.

## REFERENCES

- [1] L. Addario-Berry, R.E.L. Aldred, K. Dalal and B.A. Reed, *Vertex colouring edge partitions*, J. Combin. Theory (B) **94** (2005) 237–244.  
doi:10.1016/j.jctb.2005.01.001
- [2] M. Anholcer, S. Cichacz and M. Milanič, *Group irregularity strength of connected graphs*, J. Comb. Optim., to appear.  
doi:10.1007/s10878-013-9628-6
- [3] G. Chartrand, M.S. Jacobson, J. Lehel, O.R. Oellermann, S. Ruiz and F. Saba, *Irregular networks*, Congr. Numer. **64** (1988) 187–192.
- [4] G. Chartrand, L. Lesniak and P. Zhang, *Graphs & Digraphs: 5th Edition* (Chapman & Hall/CRC, Boca Raton, FL, 2010).
- [5] G. Chartrand and P. Zhang, *Chromatic Graph Theory* (Chapman & Hall/CRC, Boca Raton, FL, 2008).  
doi:10.1201/9781584888017
- [6] J.A. Gallian, *A dynamic survey of graph labeling*, Electron. J. Combin. **16** (2013) #DS6.
- [7] R.B. Gnana Jothi, *Topics in Graph Theory*, Ph.D. Thesis, Madurai Kamaraj University (1991).
- [8] S.W. Golomb, *How to number a graph*, in: *Graph Theory and Computing* R.C. Read (Ed.), (Academic Press, New York, 1972) 23–37.
- [9] R. Jones, *Modular and Graceful Edge Colorings of Graphs*, Ph.D. Thesis, Western Michigan University (2011).
- [10] R. Jones, K. Kolasinski, F. Fujie-Okamoto and P. Zhang, *On modular edge-graceful graphs*, Graphs Combin. **29** (2013) 901–912.  
doi:10.1007/s00373-012-1147-1
- [11] R. Jones, K. Kolasinski and P. Zhang, *A proof of the modular edge-graceful trees conjecture*, J. Combin. Math. Combin. Comput. **80** (2012) 445–455.
- [12] M. Karoński, T. Łuczak and A. Thomason, *Edge weights and vertex colours*, J. Combin. Theory (B) **91** (2004) 151–157.  
doi:10.1016/j.jctb.2003.12.001
- [13] A. Rosa, *On certain valuations of the vertices of a graph*, in: *Theory of Graphs, Proc. Internat. Symposium Rome 1966* (Gordon and Breach, New York 1967) 349–355.
- [14] V.G. Vizing, *On an estimate of the chromatic class of a  $p$ -graph*, Diskret. Analiz. **3** (1964) 25–30 (in Russian).

Received 19 July 2013

Revised 13 September 2013

Accepted 16 September 2013