

NEW BOUNDS ON THE SIGNED TOTAL DOMINATION NUMBER OF GRAPHS

SEYYED MEHDI HOSSEINI MOGHADDAM

Sama Technical and Vocational Training College
Islamic Azad University, Qom Branch, Qom, Iran

e-mail: sm.hosseini1980@yahoo.com

DOOST ALI MOJDEH

Department of Mathematics
University of Mazandaran, Babolsar, Iran

e-mail: damojdeh@umz.ac.ir

BABAK SAMADI

Department of Mathematics
Arak University, Arak, Iran

e-mail: b-samadi@araku.ac.ir

AND

LUTZ VOLKMANN

Lehrstuhl II für Mathematik
RWTH Aachen University, 52056 Aachen, Germany

e-mail: volkm@math2.rwth-aachen.de

Abstract

In this paper, we study the signed total domination number in graphs and present new sharp lower and upper bounds for this parameter. For example by making use of the classic theorem of Turán [8], we present a sharp lower bound on K_{r+1} -free graphs for $r \geq 2$. Applying the concept of total limited packing we bound the signed total domination number of G with $\delta(G) \geq 3$ from above by $n - 2 \left\lfloor \frac{2\rho_o(G) + \delta - 3}{2} \right\rfloor$. Also, we prove that $\gamma_{st}(T) \leq n - 2(s - s')$ for any tree T of order n , with s support vertices and s' support vertices of degree two. Moreover, we characterize all trees attaining this bound.

Keywords: open packing, signed total domination number, total limited packing, tuple total domination number.

2010 Mathematics Subject Classification: 05C69, 05C70.

1. INTRODUCTION

Let $G = (V, E)$ be a graph with vertex set $V = V(G)$ of order n and edge set $E = E(G)$. The minimum and maximum degree of G are denoted by $\delta = \delta(G)$ and $\Delta = \Delta(G)$, respectively. For a vertex $v \in V$, $N(v)$ is the open neighborhood of v , which is the set of vertices adjacent to v and $N[v] = N(v) \cup \{v\}$ is the closed neighborhood of v . Let $[A, B]$ be the set of edges with end points in both A and B . The set of leaves and support vertices of a tree T are denoted by $L(T)$ and $S(T)$, respectively. Also, we consider L_u as the set of all leaves adjacent to the support vertex u . We use [9] as a reference for terminology and notation which are not defined here.

A set $S \subseteq V$ is a *total dominating set* if each vertex in V is adjacent to at least one vertex in S . The *total domination number* $\gamma_t(G)$ is the minimum cardinality of a total dominating set. In [4], Henning and Slater studied the concept of open packing in graphs. A subset $B \subseteq V(G)$ is an *open packing* in G if for every distinct vertices $u, v \in B$, $N(u) \cap N(v) = \emptyset$. The *open packing number*, $\rho_o(G)$, is the maximum cardinality of an open packing in G .

A generalization of total domination titled k -tuple total domination has been studied by Henning and Kazemi in [3] (this concept had been studied by Zhao *et al.* [11] as total k -domination). A subset $S \subseteq V$ is a *k -tuple total dominating set* in G if $|N(v) \cap S| \geq k$, for all $v \in V(G)$. The *k -tuple total domination number*, $\gamma_{\times k, t}(G)$, is the smallest number of vertices in a k -tuple total dominating set.

Gallant *et al.* [1] introduced the concept of limited packing. They exhibited some real-world applications of it to network security, NIMBY, market saturation and codes. In fact, a set of vertices $B \subseteq V$ is called a *k -limited packing set* in G provided that for all $v \in V(G)$, we have $|N[v] \cap B| \leq k$. The *k -limited packing number*, denoted $L_k(G)$, is the largest number of vertices in a k -limited packing set. We can consider the concept of limited packing as the dual of tuple domination in a graph. For more information the reader can consult [6].

The above discussions give us a motivation to introduce the concept of total limited packing in graphs. Let G be a graph, and $k \in \mathbb{N}$. A set of vertices $L \subseteq V(G)$ is called a *k -total limited packing* in G provided that for all $v \in V(G)$, we have $|N(v) \cap L| \leq k$. The *k -total limited packing number*, denoted $L_{k, t}(G)$, is the largest number of vertices in a k -total limited packing set. We can consider total limited packing first as a generalization of open packing, second as a dual of tuple total domination and third as a total version of limited packing. In fact,

one can apply total limited packing to the subjects that we consider for limited packing as applications.

Let $S \subseteq V$. For a real-valued function $f : V \rightarrow \mathbb{R}$ we define $f(S) = \sum_{v \in S} f(v)$. Also, $f(V)$ is the weight of f . A *signed total dominating function*, abbreviated STDF, of G is defined in [10] as a function $f : V \rightarrow \{-1, 1\}$ such that $f(N(v)) \geq 1$ for every $v \in V$. The *signed total domination number*, abbreviated STDN, of G is $\gamma_{st}(G) = \min\{f(V) | f \text{ is a STDF of } G\}$. There exist some real-world applications of the signed total domination. For example, the author in [2] applied this concept to model networks of people or organizations in which global decisions must be made.

In this paper, we continue the study of the concept of the signed total domination in graphs. The authors noted that most of the existing bounds on $\gamma_{st}(G)$ are lower bounds. In Section 2, we prove that $\gamma_{st}(G) \leq n - 2 \lfloor \frac{2\rho_o(G) + \delta - 3}{2} \rfloor$, for a graph G of order n with $\delta(G) \geq 3$. Also, we show that $n - 2(s - s')$ is an upper bound of the signed total domination number of a tree T of order n with s support vertices and s' support vertices of degree two. Furthermore, we characterize all trees attaining this bound. In Section 3, we give some lower bounds on the signed total domination number of graphs. As an application of the well-known theorem of Turán [8] we give a lower bound on this parameter for K_{r+1} -free graphs and conclude the lower bounds given in [7] for r -partite graphs and for triangle-free graphs as special cases.

2. UPPER BOUNDS

First we apply the concept of total limited packing to obtain a sharp upper bound on $\gamma_{st}(G)$. It is easy to check that $\rho_o(K_n) = 1$, $\gamma_{st}(K_{2n}) = 2$ and $\gamma_{st}(K_{2n+1}) = 3$, for all positive integers n . We make use of them to show that the following bound is sharp.

Theorem 1. *Let G be a graph of order n and $\delta \geq 3$. Then*

$$\gamma_{st}(G) \leq n - 2 \left\lfloor \frac{2\rho_o(G) + \delta - 3}{2} \right\rfloor,$$

and this bound is sharp.

Proof. Let L be a maximum $\lfloor \frac{\delta-1}{2} \rfloor$ -total limited packing set in G . Define $f : V \rightarrow \{-1, 1\}$ by

$$f(v) = \begin{cases} -1 & \text{if } v \in L, \\ 1 & \text{if } v \in V - L. \end{cases}$$

Since L is a $\lfloor \frac{\delta-1}{2} \rfloor$ -total limited packing, $|N(v) \cap (V - L)| = \deg(v) - |N(v) \cap L| \geq \delta - \lfloor \frac{\delta-1}{2} \rfloor$. Therefore, for every vertex $v \in V$, we have $f(N(v)) = |N(v) \cap (V - L)|$

$-|N(v) \cap L| \geq \delta - \lfloor \frac{\delta-1}{2} \rfloor - \lfloor \frac{\delta-1}{2} \rfloor \geq 1$. Therefore f is a STDF of G with weight $n - 2|L| = n - 2L_{\lfloor \frac{\delta-1}{2} \rfloor, t}(G)$. This shows that

$$(1) \quad \gamma_{st}(G) \leq n - 2L_{\lfloor \frac{\delta-1}{2} \rfloor, t}(G).$$

Now let L be a maximum $\lfloor \frac{\delta-1}{2} \rfloor$ -total limited packing in G . We claim that $L \neq V$. If $L = V$ and $u \in V$ such that $\deg(u) = \Delta$, then $\Delta = |N(u) \cap L| \leq \lfloor \frac{\delta-1}{2} \rfloor$, a contradiction. Now let $u \in V - L$. It is easy to check that $|N(v) \cap (L \cup \{u\})| \leq \lfloor \frac{\delta-1}{2} \rfloor + 1$ for each $v \in V$. Therefore $L \cup \{u\}$ is a $(\lfloor \frac{\delta-1}{2} \rfloor + 1)$ -total limited packing in G . Hence $1 + L_{\lfloor \frac{\delta-1}{2} \rfloor, t}(G) = |L \cup \{u\}| \leq L_{(\lfloor \frac{\delta-1}{2} \rfloor + 1), t}(G)$. If we continue this process we finally arrive at

$$L_{\lfloor \frac{\delta-1}{2} \rfloor, t}(G) \geq 1 + L_{(\lfloor \frac{\delta-1}{2} \rfloor - 1), t}(G) \geq \dots \geq \left\lfloor \frac{\delta-1}{2} \right\rfloor - 1 + L_{1, t}(G),$$

and since $L_{1, t}(G) = \rho_o(G)$, we have

$$(2) \quad L_{\lfloor \frac{\delta-1}{2} \rfloor, t}(G) \geq \left\lfloor \frac{\delta-1}{2} \right\rfloor - 1 + \rho_o(G).$$

Inequalities (1) and (2) give the desired upper bound. Moreover, this bound is sharp. It is sufficient to consider the complete graph K_n , when $n \geq 4$. ■

As the reader can check, the signed total domination number of a tree could be arbitrary large. Henning [2] characterized all trees T of order n satisfying $\gamma_{st}(T) = n$ as all trees T in which every vertex is a support vertex or is adjacent to a vertex of degree two. Now, we bound the signed total domination number of a tree from above by considering the number of its support vertices and characterize all trees attaining this bound. For this purpose, we define Ω to be the family of all trees T satisfying:

- (a) For any support vertex u with $|L_u| \geq 2$, $\deg(u) \leq 4$;
- (b) Every vertex in $V(T) \setminus (S(T) \cup L(T))$ is adjacent to a support vertex or a vertex of degree two.

Theorem 2. *Let T be a tree of order n , with s support vertices and s' support vertices of degree two. Then*

$$\gamma_{st}(T) \leq n - 2(s - s')$$

with equality if and only if $T \in \Omega$.

Proof. Let S' be the set of all support vertices of degree two. We choose just one of the leaves adjacent to v , for every support vertex $v \in S \setminus S'$ and consider L' as the set of all those leaves. We define $f : V(T) \rightarrow \{-1, 1\}$ by $f(v) = -1$ if

$v \in L'$, and $f(v) = 1$ if $v \in V(T) \setminus L'$. It is easy to see that f is a STDF of G . Thus

$$\gamma_{st}(T) \leq f(V(T)) = n - 2|L'| = n - 2(s - s').$$

Suppose now that $f : V(T) \rightarrow \{-1, 1\}$ is a minimum STDF of T with weight $n - 2(s - s')$. Let u be a support vertex with degree at least three. Since f is a minimum STDF, there must be at least one vertex v in $N(u)$ with weight -1 under f . Without loss of generality we assume that v belongs to L' . We first show that T satisfies (a). Suppose that there is a support vertex u with $|L_u| \geq 2$ in which $\deg(u) \geq 5$, and let $v \in L' \cap N(u)$. Since $f(N(v)) \geq 1$, we have $f(u) = 1$. On the other hand, since f is a minimum STDF of T and $\deg(u) \geq 5$, there exists a vertex w in $N(u)$, different from v , with weight -1 under f . This is a contradiction because of $f(V(T)) \leq n - 2(s - s') - 1 = \gamma_{st}(T) - 1$.

We now show that T satisfies (b). Suppose there exists a vertex $v \in V(T) \setminus (S(T) \cup L(T))$, in which v is not adjacent to support vertices and all of its neighbors are of degree at least three. In this case, the function $f' : V(T) \rightarrow \{-1, 1\}$ that assigns to v the value -1 and to all other vertices w the value $f(v)$ would be a STDF of G with weight $f'(V(T)) \leq f(V(T)) - 2$. This is a contradiction.

We now let $T \in \Omega$ and $f : V(T) \rightarrow \{-1, 1\}$ be a minimum STDF of T in which $f(v) = -1$, for all $v \in L'$. Suppose to the contrary that $f(V(T)) < n - 2(s - s')$. Therefore, there exists a vertex $v \in V(T) \setminus (S(T) \cup L')$ such that $f(v) = -1$. Let $v \in L(T)$. Then $v \in N(u)$ for some $u \in S \setminus S'$. Therefore $|L_u| \geq 2$ and (a) implies $f(N(u)) \leq 0$ which is a contradiction. If $v \in V(T) \setminus (S(T) \cup L(T))$, then v is not adjacent to a support vertex, similarly. Therefore, v is adjacent to a vertex w of order two, by (b). Hence, $f(N(w)) \leq 0$ which yields a contradiction. The above discussion implies $\gamma_{st}(T) = n - 2(s - s')$. ■

We conclude this section by establishing an upper bound on the signed total domination number of a connected cubic graph. Henning [2] proved that for every cubic graph of order n , $\gamma_{st}(G) \leq 5n/7$. We now show that if G is a connected cubic graph different from the Heawood graph G_{14} , then $\gamma_{st}(G) \leq 2n/3$.

We need the following useful lemma.

Lemma 3 [5]. *If G is a connected cubic graph of order n different from the Heawood graph, then $\gamma_{\times 2,t}(G) \leq 5n/6$, and this bound is sharp.*

Theorem 4. *If G is a connected cubic graph of order n different from the Heawood graph, then $\gamma_{st}(G) \leq 2n/3$, and this bound is sharp.*

Proof. Let $f : V \rightarrow \{-1, 1\}$ be a minimum STDF of G . Let V_+ be the set of vertices u for which $f(u) = 1$. Since $f(N(v)) \geq 1$, it follows that $|N(v) \cap V_+| \geq 2$




Figure 1. Heawood graph.

for every vertex $v \in V(G)$. Hence, V_+ is a double total dominating set in G . Therefore,

$$(3) \quad (n + \gamma_{st}(G))/2 = |V_+| \geq \gamma_{\times 2,t}(G).$$

Now let D be a minimum double total dominating set in G . We define $f : V \rightarrow \{-1, 1\}$ by $f(v) = 1$, if $v \in D$ and $f(v) = -1$, if $v \in V \setminus D$. Then, $f(N(v)) = |N(v) \cap D| - |N(v) \cap (V \setminus D)| \geq 1$. Therefore f is a STDF of G . Thus,

$$(4) \quad \gamma_{st}(G) \leq f(V(G)) = 2|D| - n = 2\gamma_{\times 2,t}(G) - n.$$

The inequalities (3) and (4) together imply $\gamma_{st}(G) = 2\gamma_{\times 2,t}(G) - n$. Now Lemma 3 implies the desired upper bound. ■

3. LOWER BOUNDS

At this point we are going to present some sharp lower bounds on $\gamma_{st}(G)$. First, let us introduce some notation. Let $f : V \rightarrow \{-1, 1\}$ be a minimum STDF of G . We define $V_+ = \{v \in V | f(v) = 1\}$, $V_- = \{v \in V | f(v) = -1\}$, $G_+ = G[V_+]$ and $G_- = G[V_-]$, where G_+ and G_- are the subgraphs of G induced by V_+ and V_- , respectively. For convenience, let $E_+ = E(G_+)$, $E_- = E(G_-)$ and we define V_o and V_e as the set of vertices with odd degree and the set of vertices with even degree, respectively. Also $V_+^o = V_+ \cap V_o$, $V_+^e = V_+ \cap V_e$, $V_-^o = V_- \cap V_o$ and $V_-^e = V_- \cap V_e$. Finally, $\deg_{G_+}(v) = |N(v) \cap V_+|$ and $\deg_{G_-}(v) = |N(v) \cap V_-|$.

We begin with the following useful lemma.

Lemma 5. *Considering the above notation, the following statements hold.*

(a) $(\lfloor \frac{\delta}{2} \rfloor + 1) |V_-| \leq |[V_+, V_-]| \leq (\lceil \frac{\Delta}{2} \rceil - 1) |V_+|,$

(b) $n + |V_-| + 4|E_-| + |V_e| \leq 2|E_+| + |[V_+, V_-]|.$

Proof. (a) Let f be a minimum STDF of G . Let $v \in V_-$. Since $f(N(v)) \geq 1$, we have $\deg_{G_+}(v) \geq \left\lfloor \frac{\deg(v)}{2} \right\rfloor + 1 \geq \left\lfloor \frac{\delta}{2} \right\rfloor + 1$. Therefore $|[V_+, V_-]| \geq (\left\lfloor \frac{\delta}{2} \right\rfloor + 1)|V_-|$. Now let $v \in V_+$. Then $\deg_{G_-}(v) \leq \left\lceil \frac{\deg(v)}{2} \right\rceil - 1 \leq \left\lceil \frac{\Delta}{2} \right\rceil - 1$. Therefore $|[V_+, V_-]| \leq (\left\lceil \frac{\Delta}{2} \right\rceil - 1)|V_+|$.

(b) Let f be a minimum STDF of G . First we derive a lower bound on $|[V_+, V_-]|$. Let $v \in V_-$. Since $f(N(v)) \geq 1$, we observe that $\deg_{G_-}(v) \leq |[v, V_+]| - 1$ and $\deg_{G_-}(v) \leq |[v, V_+]| - 2$ when $\deg(v)$ is even. This leads to

$$\begin{aligned} 2|E_-| &= \sum_{v \in V_-} \deg_{G_-}(v) = \sum_{v \in V_-^e} \deg_{G_-}(v) + \sum_{v \in V_-^o} \deg_{G_-}(v) \\ &\leq \sum_{v \in V_-^e} (|[v, V_+]| - 2) + \sum_{v \in V_-^o} (|[v, V_+]| - 1) = |[V_-, V_+]| - |V_-| - |V_-^e|. \end{aligned}$$

This implies

(5) $|[V_+, V_-]| \geq |V_-| + 2|E_-| + |V_-^e|.$

Now let $v \in V_+$. Since $f(N(v)) \geq 1$, we observe that $\deg_{G_+}(v) \geq |[v, V_-]| + 1$ and $\deg_{G_+}(v) \geq |[v, V_-]| + 2$ when $\deg(v)$ is even. It follows that

$$\begin{aligned} 2|E_+| &= \sum_{v \in V_+} \deg_{G_+}(v) = \sum_{v \in V_+^e} \deg_{G_+}(v) + \sum_{v \in V_+^o} \deg_{G_+}(v) \\ &\geq \sum_{v \in V_+^e} (|[v, V_-]| + 2) + \sum_{v \in V_+^o} (|[v, V_-]| + 1) = |[V_+, V_-]| + |V_+| + |V_+^e|, \end{aligned}$$

and so

(6) $|[V_+, V_-]| \leq 2|E_+| - |V_+^e| - |V_+|.$

Combining (5) and (6), we obtain $2|E_+| \geq n + 2|E_-| + |V_e|$. Using this inequality and (5), we deduce that

$$2|E_+| + |[V_-, V_+]| \geq 2|E_-| + n + |V_e| + |V_-| + 2|E_-| + |V_-^e|,$$

and this yields (b) immediately. ■

We are now in a position to present the following lower bounds.

Theorem 6. *Let G be a graph of order n and size m . Then*

(i) $\gamma_{st}(G) \geq \frac{(\left\lfloor \frac{\delta}{2} \right\rfloor - \left\lceil \frac{\Delta}{2} \right\rceil + 2)n}{\left\lfloor \frac{\delta}{2} \right\rfloor + \left\lceil \frac{\Delta}{2} \right\rceil},$

$$(ii) \quad \gamma_{st}(G) \geq \frac{(5-3\Delta-2\lceil \frac{\Delta}{2} \rceil)n+2n_e+8m}{3\Delta+2\lceil \frac{\Delta}{2} \rceil-1},$$

$$(iii) \quad \gamma_{st}(G) \geq \frac{(5+3\delta-2\lceil \frac{\Delta}{2} \rceil)n+2n_e-4m}{3\delta+2\lceil \frac{\Delta}{2} \rceil-1},$$

and these bounds are sharp, where $n_e = |V_e|$.

Proof. (i) Using (a) of Lemma 5 and $|V_+| = \frac{n+\gamma_{st}(G)}{2}$ and $|V_-| = \frac{n-\gamma_{st}(G)}{2}$, the desired bound is easy to verify.

(ii) Using $4m = 4|E_-| + 4|[V_+, V_-]| + 4|E_+|$ and Lemma 5(b), we deduce that

$$n + |V_-| + |V_e| + 4m \leq 6|E_+| + 5|[V_+, V_-]|.$$

Applying

$$2|E_+| = \sum_{v \in V_+} \deg_{G_+}(v) = \sum_{v \in V_+} (\deg(v) - |[v, V_-]|) \leq \Delta|V_+| - |[V_+, V_-]|,$$

we obtain $n + |V_-| + |V_e| + 4m \leq 3\Delta|V_+| + 2|[V_+, V_-]|$. By Lemma 5(a), we have $2|[V_+, V_-]| \leq 2(\lceil \frac{\Delta}{2} \rceil - 1)|V_+|$ and therefore

$$n + |V_-| + |V_e| + 4m \leq \left(3\Delta + 2\left\lceil \frac{\Delta}{2} \right\rceil - 2\right) |V_+|.$$

Because of $|V_+| = \frac{n+\gamma_{st}(G)}{2}$ and $|V_-| = \frac{n-\gamma_{st}(G)}{2}$, we obtain the desired bound.

(iii) Using $2m = 2|E_-| + 2|[V_+, V_-]| + 2|E_+|$ and Lemma 5(b), we conclude that

$$n + |V_-| + 6|E_-| + |V_e| + |[V_+, V_-]| \leq 2m.$$

Applying

$$2|E_-| = \sum_{v \in V_-} \deg_{G_-}(v) = \sum_{v \in V_-} (\deg(v) - |[v, V_+]|) \geq \delta|V_-| - |[V_+, V_-]|,$$

we see that $n + (1 + 3\delta)|V_-| + |V_e| - 2|[V_+, V_-]| \leq 2m$. By Lemma 5(a), we have $2|[V_+, V_-]| \leq 2(\lceil \frac{\Delta}{2} \rceil - 1)|V_+|$ and thus

$$n + (1 + 3\delta)|V_-| + |V_e| - 2\left(\left\lceil \frac{\Delta}{2} \right\rceil - 1\right) |V_+| \leq 2m.$$

This implies the last bound.

Since $\gamma_{st}(C_n) = n$, the cycle C_n attains all of the three lower bounds. ■

We note that the bound (i) in Theorem 6 can be found in the paper [2] by Henning. However, our proof is much shorter and transparent. If G is an r -regular graph of order n , then (i) leads to $\gamma_{st}(G) \geq 2n/r$ when r is even and $\gamma_{st}(G) \geq n/r$ when r is odd. This is a result by Zelinka [10].

Theorem 7. For every graph G of order n , minimum degree $\delta \geq 1$ and maximum degree Δ ,

$$\gamma_{st}(G) \geq -n + 2 \max \left\{ \left\lceil \frac{\Delta + 3}{2} \right\rceil, \left\lceil \frac{2\gamma_t(G) + \delta - 1}{2} \right\rceil \right\},$$

and this bound is sharp.

Proof. We first prove the following claims.

Claim 1. $\gamma_{st}(G) \geq -n + 2 \lceil \frac{\Delta+3}{2} \rceil$.

Proof. Let $f : V \rightarrow \{-1, 1\}$ be a minimum STDF of G . Since $f(N(v)) \geq 1$, it follows that $|N(v) \cap V_-| \leq \lfloor \frac{\Delta-1}{2} \rfloor$ for every vertex $v \in V(G)$. Therefore V_- is a $\lfloor \frac{\Delta-1}{2} \rfloor$ -total limited packing in G . Thus

$$(7) \quad (n - \gamma_{st}(G))/2 = |V_-| \leq L_{\lfloor \frac{\Delta-1}{2} \rfloor, t}(G).$$

On the other hand, similar to the proof of Theorem 1 and taking into account that $L_{\Delta+1, t}(G) = n$, we obtain

$$\begin{aligned} L_{\lfloor \frac{\Delta-1}{2} \rfloor, t}(G) &\leq L_{\lfloor \frac{\Delta-1}{2} \rfloor + 1, t}(G) - 1 \leq \dots \\ &\leq L_{\Delta+1, t}(G) - \Delta + \lfloor \frac{\Delta-3}{2} \rfloor = n - \lceil \frac{\Delta+3}{2} \rceil. \end{aligned}$$

Now inequality (7) implies $\gamma_{st}(G) \geq -n + 2 \lceil \frac{\Delta+3}{2} \rceil$, as desired. □

Claim 2. $\gamma_{st}(G) \geq -n + 2 \lceil \frac{2\gamma_t(G) + \delta - 1}{2} \rceil$.

Proof. Since $f(N(v)) \geq 1$, it follows that $|N(v) \cap V_+| \geq \lceil \frac{\delta+1}{2} \rceil$ for every vertex $v \in V(G)$. Therefore V_+ is a $\lceil \frac{\delta+1}{2} \rceil$ -tuple total dominating set in G . Thus

$$(8) \quad (n + \gamma_{st}(G))/2 = |V_+| \geq \gamma_{\times \lceil \frac{\delta+1}{2} \rceil, t}(G).$$

Now let D be a minimum $\lceil \frac{\delta+1}{2} \rceil$ -tuple total dominating set in G . Then $|N(v) \cap D| \geq \lceil \frac{\delta+1}{2} \rceil$, for every vertex $v \in V(G)$. Let $u \in D$. It is easy to see that $|N(v) \cap (D \setminus \{u\})| \geq \lceil \frac{\delta+1}{2} \rceil - 1$, for all $v \in V(G)$. Hence, $D \setminus \{u\}$ is a $(\lceil \frac{\delta+1}{2} \rceil - 1)$ -tuple total dominating set in G . Therefore, $\gamma_{\times \lceil \frac{\delta+1}{2} \rceil, t}(G) \geq \gamma_{\times (\lceil \frac{\delta+1}{2} \rceil - 1), t}(G) + 1$. By repeating this process, we obtain

$$\begin{aligned} \gamma_{\times \lceil \frac{\delta+1}{2} \rceil, t}(G) &\geq \gamma_{\times (\lceil \frac{\delta+1}{2} \rceil - 1), t}(G) + 1 \geq \dots \\ &\geq \gamma_{\times 1, t}(G) + \lceil \frac{\delta+1}{2} \rceil - 1 = \gamma_t(G) + \lceil \frac{\delta-1}{2} \rceil. \end{aligned}$$

By inequality (8),

$$(n + \gamma_{st}(G))/2 \geq \gamma_t(G) + \left\lceil \frac{\delta - 1}{2} \right\rceil.$$

This completes the proof of Claim 2. □

The result now follows by Claim 1 and Claim 2. The bound is sharp for the complete graph. ■

A graph is K_p -free if it does not contain the complete graph K_p as a subgraph. For our next lower bound, we use the following well-known theorem of Turán [8].

Theorem 8 [8]. *If G is a K_{r+1} -free graph of order n , then*

$$|E(G)| \leq \frac{r-1}{2r} \cdot n^2.$$

Theorem 9. *Let $r \geq 2$ be an integer, and let G be a K_{r+1} -free graph of order n . If $c = \lceil (\delta(G) + 1)/2 \rceil$, then*

$$\gamma_{st}(G) \geq \frac{r}{r-1} \left(-(c-1) + \sqrt{(c-1)^2 + 4\frac{r-1}{r}cn} \right) - n.$$

Proof. By Lemma 5(a), we have

$$(9) \quad |[V_+, V_-]| \geq \left\lceil \frac{\delta(G) + 1}{2} \right\rceil |V_-| = c|V_-| = c(n - |V_+|).$$

Furthermore, Theorem 8 leads to

$$\begin{aligned} |[V_+, V_-]| &= \sum_{v \in V_+} |N(v) \cap V_-| \leq \sum_{v \in V_+} (|N(v) \cap V_+| - 1) \\ &= 2|E(G[V_+])| - |V_+| \leq \frac{r-1}{r}|V_+|^2 - |V_+|. \end{aligned}$$

Combining this inequality chain with (9), we obtain

$$(n - |V_+|)c \leq \frac{r-1}{r}|V_+|^2 - |V_+|$$

and thus

$$\frac{r-1}{r}|V_+|^2 + (c-1)|V_+| - cn \geq 0.$$

It follows that

$$|V_+| \geq \frac{r}{2(r-1)} \left(-(c-1) + \sqrt{(c-1)^2 + 4\frac{r-1}{r}cn} \right),$$

and so we arrive at the desired bound

$$\gamma_{st}(G) = 2|V_+| - n \geq \frac{r}{r-1} \left(-(c-1) + \sqrt{(c-1)^2 + 4\frac{r-1}{r}cn} \right) - n. \quad \blacksquare$$

For the special case when G is an r -partite graph, Theorem 9 was proved by Shan and Cheng in [7]. In that paper the authors have constructed r -partite graphs with equality in the inequality of Theorem 9, and therefore this theorem is sharp. In addition, one can find Theorem 9 for triangle-free graphs in [7].

Acknowledgements

The authors wish to express their sincere thanks to the referees for their valuable suggestions and comments.

REFERENCES

- [1] R. Gallant, G. Gunther, B. Hartnell and D.F. Rall, *Limited packing in graphs*, Discrete Appl. Math. **158** (2010) 1357–1364.
doi:10.1016/j.dam.2009.04.014
- [2] M.A. Henning, *Signed total domination in graphs*, Discrete Math. **278** (2004) 109–125.
doi:10.1016/j.disc.2003.06.002
- [3] M.A. Henning and A.P. Kazemi, *k -tuple total domination in graphs*, Discrete Appl. Math. **158** (2010) 1006–1011.
doi:10.1016/j.dam.2010.01.009
- [4] M.A. Henning and P.J. Slater, *Open packing in graphs*, J. Combin. Math. Combin. Comput. **28** (1999) 5–18.
- [5] M.A. Henning and A. Yeo, *Strong transversals in hypergraphs and double total domination in graphs*, SIAM J. Discrete Math. **24** (2010) 1336–1355.
doi:10.1137/090777001
- [6] D.A. Mojdeh, B. Samadi and S.M. Hosseini Moghaddam, *Limited packing vs tuple domination in graphs*, Ars Combin. (to appear).
- [7] E. Shan and T.C.E. Cheng, *Remarks on the minus (signed) total domination in graphs*, Discrete Math. **308** (2008) 3373–3380.
doi:10.1016/j.disc.2007.06.015
- [8] P. Turán, *On an extremal problem in graph theory*, Math. Fiz. Lapok **48** (1941) 436–452.
- [9] D.B. West, Introduction to Graph Theory (Second Edition, Prentice Hall, USA, 2001).
- [10] B. Zelinka, *Signed total domination number of a graph*, Czechoslovak Math. J. **51** (2001) 225–229.
doi:10.1023/A:1013782511179
- [11] W. Zhao, H. Wang and G. Xu, *Total k -domination number in graphs*, Int. J. Pure Appl. Math. **35** (2007) 235–242.

Received 9 February 2015

Revised 14 August 2015

Accepted 14 August 2015