

ON THE CROSSING NUMBERS OF CARTESIAN PRODUCTS OF WHEELS AND TREES

MARIÁN KLEŠČ, JANA PETRILLOVÁ

AND

MATÚŠ VALO

Department of Mathematics and Theoretical Informatics
Faculty of Electrical Engineering and Informatics
Technical University, 042 00 Košice, Slovak Republic

e-mail: marian.klesc@tuke.sk
jana.petrilova@tuke.sk
matus.valo@tuke.sk

Abstract

Bokal developed an innovative method for finding the crossing numbers of Cartesian product of two arbitrarily large graphs. In this article, the crossing number of the join product of stars and cycles are given. Afterwards, using Bokal's zip product operation, the crossing numbers of the Cartesian products of the wheel W_n and all trees T with maximum degree at most five are established.

Keywords: graph, drawing, crossing number, join product, Cartesian product.

2010 Mathematics Subject Classification: 05C10, 05C38.

1. INTRODUCTION

Let G be a graph, whose vertex set and edge set are denoted by $V(G)$ and $E(G)$, respectively. A drawing of G is a representation of G in the plane such that its vertices are represented by distinct points and its edges by simple continuous arcs connecting the corresponding pair of points. For simplicity, we assume that in a drawing (a) no edge passes through any vertex other than its end-points, (b) no two edges touch each other (i.e., if two edges have a common interior point, then at this point they properly cross each other), and (c) no three edges cross at the

same point. The *crossing number* $cr(G)$ is the smallest number of edge crossings in any drawing of G . It is easy to see that a drawing with minimum number of crossings (an *optimal* drawing) is always a *good* drawing, meaning that no edge crosses itself, no two edges cross more than once, and no two edges incident with the same vertex cross.

The repetitive patterns in Cartesian products of graphs reflects in their drawings and makes Cartesian products one of the first graph classes for which the crossing numbers were studied (for a definition of Cartesian product see [15]). Let C_n be the cycle of length n , P_n be the path on n vertices, and S_n be the star isomorphic to $K_{1,n}$. The crossing numbers of Cartesian products of graphs have been studied since 1973, when Harary *et al.* established the crossing number of $C_3 \square C_3$ and conjectured that $cr(C_m \square C_n) = (m - 2)n$ for $3 \leq m \leq n$, [11]. This hypothesis has been proved for $m \leq 7$ by other authors, but it has not yet been proved for $3 \leq m \leq n$ in general. The best result was given by Glebsky and Salazar in [10].

Several authors were researching the crossing numbers of the Cartesian products $G \square C_n$, $G \square P_n$, $G \square S_n$ for some specific graphs G . Beineke and Ringel in [3] as well as Jendroľ and Ščerbová in [14] determined the crossing numbers of the Cartesian products of four-vertex graphs with the cycle C_n . Some other results concerning the crossing numbers of Cartesian products of special small graphs with paths, cycles and stars one can find in [15, 16, 19, 20], and [25]. Moreover, Jendroľ and Ščerbová in [14] conjectured that the crossing number of $S_n \square P_m$ is $(m - 2) \lfloor \frac{n}{2} \rfloor \lfloor \frac{m-1}{2} \rfloor$. This conjecture was proved by Bokal using zip product operation, see [4]. It was the first exact result which gives the crossing number of Cartesian product of two graphs, where both graphs are arbitrarily large. In [5], Bokal extended the properties of zip product operation and established the crossing numbers of Cartesian products of trees with stars as well as of paths with wheels. Another results concerning Cartesian products of two arbitrarily large graphs were obtained in [26] and [27]. In the paper, we extend these results by giving the crossing numbers of the Cartesian products of trees with wheels. Our methods are based on some properties of zip product operation and of known values of crossing numbers of join of stars and cycles.

The *join product* $G_i + G_j$ of graphs G_i and G_j is created from vertex-disjoint copies of G_i and G_j by adding all edges between $V(G_i)$ and $V(G_j)$. For $|V(G_i)| = m$ and $|V(G_j)| = n$, the edge set of the graph $G_i + G_j$ is the union of $E(G_i)$, $E(G_j)$ and $E(K_{m,n})$. The first results on crossing numbers of join of paths and cycles as well as of two cycles appeared in [17]. There are also some known results concerning crossing numbers of join products of discrete graphs, paths and cycles with special graphs, see for example [18, 21, 22] and [23].

We denote the number of crossings of a graph G in a drawing D by $cr_D(G)$. Let G_i and G_j be edge-disjoint subgraphs of a graph G . Then we denote by

$\text{cr}_D(G_i, G_j)$ the number of crossings between the edges of G_i and the edges of G_j in the drawing D . In the proofs of the paper, we will often use the term “region” also in nonplanar drawings. In this case, crossings are considered to be vertices of the “map”. In a good drawing D of the graph G , we say that a cycle C *separates* the vertices of a subgraph G_i not containing vertices of C if the vertices of G_i are contained in different components of $\mathbb{R}^2 \setminus C$.

In the paper, some proofs are based on Kleitman’s result on crossing numbers of complete bipartite graphs. More precisely, in [24] he proved that

$$\text{cr}(K_{p,q}) = \left\lfloor \frac{p}{2} \right\rfloor \left\lfloor \frac{p-1}{2} \right\rfloor \left\lfloor \frac{q}{2} \right\rfloor \left\lfloor \frac{q-1}{2} \right\rfloor, \quad \text{if } \min\{p, q\} \leq 6.$$

In Section 2, we give some special properties concerning drawings of join products of graphs with cycles, especially of join products of discrete graphs with cycles and of their subgraphs. Using these results, in Section 3 we find the exact values of crossing numbers for join products of stars with cycles. Let W_n be the wheel on $n + 1$ vertices and let D_m denote the discrete graph on m vertices. In Section 4, the isomorphism between $S_m + C_n$ and $W_n + D_m$, together with the Bokal’s results on zip product operation, allows us to establish the crossing number of Cartesian product of all trees with maximum degree at most five with all wheels.

2. PRELIMINARY RESULTS

Let us consider a graph G with $V(G) = \{v_1, v_2, \dots, v_m\}$ and the cycle C_n with the vertices c_1, c_2, \dots, c_n . The join product $G + C_n$ consists of one copy of the graph G , one copy of the cycle C_n , and of the edges joining every vertex of G with every vertex of C_n . The edges joining the vertices of G with the vertices of C_n form the complete bipartite graph $K_{m,n}$. For the vertices v_1, v_2, \dots, v_m of the graph G , let T^i denote the subgraph induced by n edges joining the vertex v_i with the vertices c_1, c_2, \dots, c_n of the cycle C_n . So,

$$G + C_n = G \cup K_{m,n} \cup C_n = G \cup \left(\bigcup_{i=1}^m T^i \right) \cup C_n.$$

Lemma 1. *Let G be a graph of order m , $m \geq 1$. In an optimal drawing of the join product $G + C_n$, $n \geq 3$, the edges of C_n do not cross each other.*

Proof. Assume an optimal drawing of the graph $G + C_n$ in which two edges of C_n cross. Any optimal drawing is good and in such a drawing, the edges of a 3-cycle cannot cross. Thus, assume $n \geq 4$ in the rest of the proof. Let x be the point of the plane in which two edges, say $\{c_i, c_{i+1}\}$ and $\{c_j, c_{j+1}\}$, of C_n cross. The plane is a normal space. Hence, in the plane there is an open

set A_x such that A_x contains x together with the corresponding segments of the crossed edges. All remaining edges of the drawing are disjoint with A_x , see Figure 1(a). Figure 1(b) shows that the edges $\{c_i, c_{i+1}\}$ and $\{c_j, c_{j+1}\}$ can be redrawn into new edges $\{c_i, c_j\}$ and $\{c_{i+1}, c_{j+1}\}$ which do not cross. The vertices $c_1, \dots, c_{i-1}, c_i, c_j, c_{j-1}, \dots, c_{i+2}, c_{i+1}, c_{j+1}, c_{j+2}, \dots, c_n, c_1$ form the n -cycle again. Since every vertex of the cycle C_n is adjacent to every vertex of the graph G , the new drawing of the graph $G + C_n$ with less number of crossings is obtained. This contradiction completes the proof. ■




Figure 1. Elimination of a crossing in C_n .

We note that, in Section 3, only optimal drawings of stars with cycles are considered and therefore, in the whole section it will be assumed that the cycle C_n does not cross itself.

Assume now the discrete graph D_m on m vertices. Clearly, $D_m + C_n$ is a subgraph of the graph $G + C_n$, where $|V(G)| = m$. In the proofs of the paper, several special properties of the graph $D_m + C_n$ are used. Let us present some of them. Several results of the paper are based on the next Lemma 2. Although Lemma 2 was proved in [17], we prove it again. This proof is simpler than the previous one.

Lemma 2. *Let D be a good drawing of the join product $D_m + C_n$, $m \geq 2$, $n \geq 3$, in which no edge of C_n is crossed and C_n does not separate the other vertices of the graph. Then, for all $i, j = 1, 2, \dots, m$, two different subgraphs T^i and T^j cross each other at least $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times in D .*

Proof. By hypothesis, the cycle C_n divides the plane into two regions in such a way that all m subgraphs T^1, T^2, \dots, T^m are placed in one of these regions, say in the interior region of C_n . Assume that two different subgraphs T^i and T^j cross each other less than $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times. As both are placed inside C_n , it is possible to add a new vertex w into the unbounded region of C_n together with the edges $\{w, c_1\}, \{w, c_2\}, \dots, \{w, c_n\}$ such that none of them crosses an edge of $T^i \cup T^j$. Now the edges of the new graph incident with the vertices v_i, v_j , and w form the subgraph isomorphic to $K_{3,n}$ and cross each other less than $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times. This contradiction with $cr(K_{3,n}) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ completes the proof. ■

For the case when the cycle C_n separates some of the other vertices of the graph, or when some subgraphs T^i , $i \in \{1, 2, \dots, m\}$, cross C_n , we have the next corollary.

Corollary 3. *Let D be a good drawing of the join product $D_m + C_n$, $m \geq 2$, $n \geq 3$, in which the edges of C_n do not cross each other and C_n does not separate r vertices $v_{i_1}, v_{i_2}, \dots, v_{i_r}$, $2 \leq r \leq m$. If none of the subgraphs $T^{i_1}, T^{i_2}, \dots, T^{i_r}$ induced on the edges incident with the vertices $v_{i_1}, v_{i_2}, \dots, v_{i_r}$ crosses C_n , then two different subgraphs T^{i_j} and T^{i_k} , $j, k = 1, 2, \dots, r$, cross each other at least $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times in D .*

Proof. By the assumptions, the drawing of $D_m + C_n$ contains the subdrawing of $D_r + C_n$ satisfying the conditions of Lemma 2. Thus, the result follows. ■

Corollary 4. *Let D be a good drawing of the join product $D_m + C_n$, $m \geq 2$, $n \geq 3$, in which the edges of C_n do not cross each other and C_n does not separate r vertices $v_{i_1}, v_{i_2}, \dots, v_{i_r}$, $2 \leq r \leq m$. Let $T^{i_1}, T^{i_2}, \dots, T^{i_s}$, $s < r$, be the subgraphs induced on the edges incident with the vertices $v_{i_1}, v_{i_2}, \dots, v_{i_s}$ that do not cross C_n . If k edges of some subgraph T^{i_j} induced on the edges incident with the vertex v_{i_j} , $j \in \{s + 1, s + 2, \dots, r\}$, cross the cycle C_n , then the subgraph T^{i_j} crosses each of the subgraphs $T^{i_1}, T^{i_2}, \dots, T^{i_s}$ at least $\lfloor \frac{n-k}{2} \rfloor \lfloor \frac{(n-k)-1}{2} \rfloor$ times in D .*

Proof. When $k > n - 3$, the assertion is true. For $k \leq n - 3$, consider $n - k$ vertices of the cycle C_n incident with the edges of T^{i_j} which do not cross C_n . Let us delete all edges of T^1, T^2, \dots, T^m which are not incident with these $n - k$ vertices. The resulting subgraph is homeomorphic to the graph $D_m + C_{n-k}$ and, in its subdrawing induced by D , all subgraphs $T^{i_1}, T^{i_2}, \dots, T^{i_s}$ as well as T^{i_j} of $D_m + C_{n-k}$ satisfies the conditions of Corollary 3. Hence, the subgraph T^{i_j} crosses each of the subgraphs $T^{i_1}, T^{i_2}, \dots, T^{i_s}$ at least $\lfloor \frac{n-k}{2} \rfloor \lfloor \frac{(n-k)-1}{2} \rfloor$ times. ■

3. JOIN OF STARS WITH CYCLES

Our aim in this section is to give the crossing numbers of the join products $S_m + C_n$ for $m \leq 5$ and $n \geq 3$. Clearly, the graph $K_1 + C_n$ is planar. It is shown in [17] that $cr(P_k + C_n) = \lfloor \frac{k}{2} \rfloor \lfloor \frac{k-1}{2} \rfloor \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$ for $k \geq 2$. As $S_1 + C_n$ is isomorphic to $P_2 + C_n$ and $S_2 + C_n$ is isomorphic to $P_3 + C_n$, the crossing number of $S_1 + C_n$ is one, and the crossing number of $S_2 + C_n$ is $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$. For $m \geq 3$, our proofs are based on Kleitman's result $cr(K_{p,q}) = \lfloor \frac{p}{2} \rfloor \lfloor \frac{p-1}{2} \rfloor \lfloor \frac{q}{2} \rfloor \lfloor \frac{q-1}{2} \rfloor$ for $\min\{p, q\} \leq 6$. This is the reason that, unfortunately, we are not able to give the crossing number of the graph $S_m + C_n$ for $m > 5$ and arbitrarily large n , because the graph $S_6 + C_n$ contains $K_{7,n}$ as a subgraph.

In this section, the crossing numbers of the graphs $S_m + C_n$ for $m = 3, 4, 5$, and for all $n \geq 3$ are established. The methods used in the general proofs are successful for $n \geq 5$. For the small graphs $S_3 + C_3$, $S_3 + C_4$, $S_4 + C_3$, $S_4 + C_4$, $S_5 + C_3$, and $S_5 + C_4$, the crossing numbers must be done separately. We tried to compute these values of crossing numbers using algorithm located on the website <http://crossings.uos.de/>. This algorithm can find the crossing numbers of small undirected graphs. It uses an ILP formulation, based on Kuratowski subgraphs, and solves it via branch-and-cut-and-price, see [6, 7] and [8]. The system also generates verifiable formal proofs, as described in [9]. Unfortunately, the capacity of this system is restricted and it does not establish crossing numbers for large graphs.

In the reasonable time, the algorithm found the exact values of crossing numbers for the graphs $S_3 + C_3$, $S_3 + C_4$, $S_4 + C_3$, $S_4 + C_4$, and $S_5 + C_3$. The next Lemma 5 lists these results. We remark that we can prove that all these results are correct. All these proofs are similar to the proof of Lemma 6. Moreover, all of them are much easier. So, we omit these proofs and the reader can find them on his own. Since for the graph $S_5 + C_4$ the algorithm in <http://crossings.uos.de/> found only a lower bound of 20 and an upper bound of 23, we prove in Lemma 6 that the crossing number of the graph $S_5 + C_4$ is 23.

Lemma 5. $\text{cr}(S_3 + C_3) = 5$, $\text{cr}(S_3 + C_4) = \text{cr}(S_4 + C_3) = 8$, $\text{cr}(S_4 + C_4) = 14$, and $\text{cr}(S_5 + C_3) = 13$.




Figure 2. The drawing of the graph $S_5 + C_n$.

The drawing in Figure 2 shows the graph $S_5 + C_n$. Let us denote by v_0, v_1, v_2, v_3, v_4 , and v_5 the vertices of the star S_5 , where v_0 is the central vertex, and let

c_1, c_2, \dots, c_n be the vertices of the cycle C_n . For the graph $S_5 + C_n$, as well as for its subgraphs $S_3 + C_n$ and $S_4 + C_n$, let T^{v_i} denote the subgraph induced on the n edges incident with the vertex $v_i, i = 0, 1, 2, \dots, 5$. The edges of $\bigcup_{i=0}^5 T^{v_i} \cong K_{6,n}$ cross each other $6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times. For every vertex $c_j, j = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$, the edges incident with c_j cross the edges of the star S_5 four times, and the cycle C_n crosses S_5 three times. So, in Figure 2, exactly $6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 3$ crossings appear among the edges of the graph $S_5 + C_n$ in this drawing.

Lemma 6. $cr(S_5 + C_4) = 23$.

Proof. It follows from the drawing in Figure 2 that $cr(S_5 + C_4) \leq 23$. To prove the reverse inequality assume that there is an optimal drawing D of the graph $S_5 + C_4$ with at most 22 crossings. By Lemma 1, the edges of C_4 do not cross each other in D . Moreover, the edges of C_4 are crossed at most twice in D , otherwise by deleting the edges of C_4 from D a drawing of the complete tripartite graph $K_{1,5,4}$ with less than 20 crossings is obtained. This contradicts the fact that $cr(K_{1,5,4}) = 20$, see [13]. By Lemma 5, $cr(S_4 + C_4) = 14$. This implies that in D there are at most eight crossings on the edges of every subgraph $T^{v_i} \cup \{v_0, v_i\}, i = 1, 2, 3, 4, 5$. As $(S_5 + C_4) \setminus E(C_4 \cup T^{v_0})$ is isomorphic to the complete bipartite graph $K_{5,5}$ with crossing number 16, at most six crossings appear on the edges of $C_4 \cup T^{v_0}$. We note that C_4 is crossed at least once, because otherwise all subgraphs $T^{v_i}, i = 0, 1, 2, 3, 4, 5$, are placed in the same region of C_4 and, by Lemma 2, in D there are at least $2 \binom{6}{2} > 22$ crossings.

The cycle C_4 divides the plane into two regions and, in D , the vertex v_0 is placed in one of them, say in the interior region. Our next analysis depends on the number of crossings between the edges of C_4 and the edges of the subgraph T^{v_0} in the drawing D . If $cr_D(T^{v_0}, C_4) = 0$, then at least three subgraphs $T^{v_i}, i \in \{1, 2, 3, 4, 5\}$, are placed in the same region as v_0 such that $cr(T^{v_i}, C_4) = 0$. By Corollary 3, each such subgraph T^{v_i} crosses T^{v_0} at least twice and, together with at least one crossing on C_4 , on the edges of $C_4 \cup T^{v_0}$ there are more than six crossings. Thus, $cr_D(T^{v_0}, C_4) \neq 0$.

If $cr_D(T^{v_0}, C_4) = 2$, no other crossing appears on the edges of C_4 . In this case, the star S_5 is placed in the interior region of C_4 such that none of $T^{v_i}, i = 1, 2, 3, 4, 5$, crosses C_4 . Thus, by Corollary 3, in D there are at least $2 \binom{5}{2} + 2 = 22$ crossings among the edges of $(S_5 + C_4) \setminus E(S_5)$. This forces that $cr_D(C_4 \cup T^{v_0}, T^{v_i} \cup \{v_0, v_i\}) = 0$ for all $i = 1, 2, 3, 4, 5$ and in the subdrawing induced by $C_4 \cup T^{v_0}$ there must exist a region containing all five vertices c_1, c_2, c_3, c_4 , and v_0 on its boundary. The only possibility to obtain such a drawing is that one edge of C_4 is crossed by two edges of T^{v_0} . Without loss of generality, let the edge $\{c_1, c_2\}$ of C_4 is crossed by the edges $\{v_0, c_3\}$ and $\{v_0, c_4\}$. Thus, the 5-cycle $c_1 v_0 c_2 c_3 c_4 c_1$ forms the boundary of the region containing the vertices v_1, v_2, v_3, v_4 , and v_5 . This boundary together with the edges joining the vertices v_1, v_2, v_3, v_4 , and v_5 with

the vertices c_1, c_2, c_3, c_4 , and v_0 form the graph $D_5 + C_5$. Hence, by Lemma 2, in D there are at least $4\binom{5}{2} > 22$ crossings. This confirms that $\text{cr}_D(T^{v_0}, C_4) = 1$.

Assume first that the crossing between C_4 and T^{v_0} is the only crossing on the edges of C_4 . Then the subdrawing of $C_4 \cup T^{v_0}$ induced by D is unique as shown in Figure 3(a). Clearly, every subgraph T^{v_i} , $i = 1, 2, 3, 4, 5$, crosses T^{v_0} . Moreover, none of them crosses T^{v_0} twice, otherwise the edges of $C_4 \cup T^{v_0}$ are crossed more than six times. Hence, in D , all five vertices v_i , $i \neq 0$, are placed in the region α of the subdrawing of $C_4 \cup T^{v_0}$. Now, the boundary of the region α forms the 4-cycle and, in D , all five vertices v_i inside α are adjacent with four vertices on its boundary such that the considered edges do not cross the boundary of α . Thus, applying Lemma 2 on the subgraph isomorphic to $D_5 + C_4$, the edges of $\bigcup_{i=0}^5 T^{v_i}$ cross each other at least $2\binom{5}{2}$ times and in D there are at least $2\binom{5}{2} + 5 + 1 > 22$ crossings.




Figure 3. The subdrawings of $C_4 \cup T^{v_0}$ and $C_4 \cup T^{v_0} \cup T^{v_1} \cup \{v_0, v_1\}$.

The last possibility is that $\text{cr}_D(T^{v_0}, C_4) = 1$ and the edges of C_4 are crossed twice. This implies that $\text{cr}_D(S_5, C_4 \cup T^{v_0}) = 0$, otherwise, together with the crossings between five subgraphs T^{v_i} and $C_4 \cup T^{v_0}$, more than six crossings appear on the edges of $C_4 \cup T^{v_0}$. Thus, one of the subgraphs T^{v_i} , say T^{v_1} , crosses C_4 once. The subdrawing of $C_4 \cup T^{v_1}$ induced by D is the same as the drawing in Figure 3(a) if we replace the vertex v_0 by the vertex v_1 . Since no other crossing is allowed on C_4 , every subgraph T^{v_i} , $i = 2, 3, 4, 5$, crosses both T^{v_0} and T^{v_1} . Moreover, by Corollary 3, the edges of $T^{v_2} \cup T^{v_3} \cup T^{v_4} \cup T^{v_5}$ cross each other at least $2\binom{4}{2}$ times. So, at least 22 crossings appear in D on the edges other than the edges of S_5 . Now, the unique subdrawing of the subgraph $C_4 \cup T^{v_0} \cup T^{v_1} \cup \{v_0, v_1\}$ induced by D is shown in Figure 3(b). It is easy to verify that every subgraph T^{v_i} , $i = 2, 3, 4, 5$, crosses the edges of $T^{v_0} \cup T^{v_1} \cup \{v_0, v_1\}$ at least three times and in D there are at least $4 \cdot 3 + 2\binom{4}{2} + 2 > 22$ crossings. This completes the proof. ■

Theorem 7. $\text{cr}(S_3 + C_n) = 2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + \lfloor \frac{n}{2} \rfloor + 2$ for $n \geq 3$.

Proof. By Lemma 5, the result is true for $n = 3$ and $n = 4$. The drawing of the graph $S_3 + C_n$ with $2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + \lfloor \frac{n}{2} \rfloor + 2 = \frac{1}{2}n(n-1) + 2$ crossings can be obtained by deleting the vertices v_4 and v_5 from the drawing in Figure 2. So, $\text{cr}(S_3 + C_n) \leq \frac{1}{2}n(n-1) + 2$. To prove the reverse inequality suppose that there is an optimal drawing D with less than $\frac{1}{2}n(n-1) + 2$ crossings for $n \geq 5$. The subgraph $(S_3 + C_n) \setminus E(C_n)$ is isomorphic to the complete tripartite graph $K_{1,3,n}$. Asano in [2] proved that $\text{cr}(K_{1,3,n}) = \frac{1}{2}n(n-1)$. This forces that the edges of C_n are crossed at most once in D .

By Lemma 1, the edges of C_n do not cross each other in D and the subdrawing of C_n induced by D divides the plane into two regions. If the edges of C_n are not crossed, then all four vertices v_0, v_1, v_2 , and v_3 are placed in one of these two regions and, by Lemma 2, in D there are at least $\binom{4}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor > \frac{1}{2}n(n-1) + 1$ crossings, a contradiction. So, if all vertices v_0, v_1, v_2 , and v_3 are placed in D in the same region of C_n , exactly one subgraph T^{v_i} , $i \in \{0, 1, 2, 3\}$, crosses C_n once. But, for $n \geq 4$, it crosses also all three subgraphs T^{v_j} , $j \in \{0, 1, 2, 3\}$, $j \neq i$, and Corollaries 3 and 4 imply that in D there are at least $\binom{3}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 3 \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor + 1 > \frac{1}{2}n(n-1) + 1$ crossings. This contradicts the assumption of D . Hence, the cycle C_n separates the vertices of S_3 . As C_n does not have more than one crossing, one vertex, say v_3 , of degree one of S_3 is separated from the other three. In this case, by Corollary 3, the edges of $T^{v_0} \cup T^{v_1} \cup T^{v_2}$ cross each other at least $\binom{3}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times and in D there are at least $3 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$ crossings. This number exceeds the considered number of crossings in D for $n \geq 5$ and the proof is done. ■

Theorem 8. $\text{cr}(S_4 + C_n) = 4 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2 \lfloor \frac{n}{2} \rfloor + 2$ for $n \geq 3$.

Proof. By Lemma 5, the result is true for $n = 3$ and $n = 4$. The drawing of the graph $S_4 + C_n$ with $4 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2 \lfloor \frac{n}{2} \rfloor + 2 = n(n-1) + 2$ crossings can be obtained by deleting the vertex v_5 from the drawing in Figure 2. Hence, $\text{cr}(S_4 + C_n) \leq n(n-1) + 2$. To prove the reverse inequality assume that, for $n \geq 5$, there is an optimal drawing D of the graph $S_4 + C_n$ with fewer than $n(n-1) + 2$ crossings.

By Lemma 1, the edges of C_n do not cross each other in D . The subgraph $(S_4 + C_n) \setminus E(C_n)$ of the graph $S_4 + C_n$ is isomorphic to the complete tripartite graph $K_{1,4,n}$. It was shown in [12] that $\text{cr}(K_{1,4,n}) = n(n-1)$. This implies that the edges of C_n are crossed at most once in D . On the other hand, if no edge of C_n is crossed, then all five vertices of S_4 are placed in the same region in the view of the subdrawing of C_n induced by D and, by Lemma 2, in D there are at least $\binom{5}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ crossings. This number is greater than the considered number of crossings in the drawing D . Thus, the edges of C_n are crossed exactly once in D .

Regardless of C_n is crossed by S_4 or by some subgraph T^{v_i} , $i \in \{0, 1, 2, 3, 4\}$, at least four subgraphs T^{v_i} are placed in the same region of the subdrawing of C_n and none of them crosses C_n . Thus, by Corollary 3, in D there are at least $\binom{4}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$ crossings. This exceeds the considered number of crossings in D for $n \geq 5$, and the proof is done. ■

Theorem 9. $\text{cr}(S_5 + C_n) = 6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 3$ for $n \geq 3$.

Proof. By Lemmas 5 and 6, Theorem 9 is true when $n = 3$ and $n = 4$. The drawing in Figure 2 shows that $\text{cr}(S_5 + C_n) \leq 6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 3$. For $n \geq 5$, assume that there is an optimal drawing D of the graph $S_5 + C_n$ with at most $6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 2$ crossings. By Lemma 1, the edges of C_n do not cross each other in D . The subgraph $(S_5 + C_n) \setminus E(C_n)$ is isomorphic to the complete tripartite graph $K_{1,5,n}$. It was proved in [13] that $\text{cr}(K_{1,5,n}) = 6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor$. Thus, C_n is crossed at most twice in D . At most four vertices v_i , $i \in \{0, 1, 2, 3, 4, 5\}$, are placed in the same region of C_n such that the corresponding subgraphs T^{v_i} do not cross C_n . Otherwise, by Corollary 3, in D there are at least $\binom{5}{2} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor > 6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 2$ crossings. This forces that the edges of C_n are crossed exactly twice and that every subgraph T^{v_i} crosses C_n at most once. In addition, as the deleting of all edges of the subgraph $C_n \cup T^{v_0}$ results in the graph $K_{5,n+1}$ with crossing number $4 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n+1}{2} \rfloor$, at most $2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2$ crossings can appear on the edges of $C_n \cup T^{v_0}$ in D .

We show first that T^{v_0} crosses C_n in D . If $\text{cr}_D(T^{v_0}, C_n) = 0$, at least three subgraphs T^{v_i} , $i \in \{1, 2, 3, 4, 5\}$, are placed in the same region of C_n as the vertex v_0 with $\text{cr}_D(T^{v_i}, C_n) = 0$. By Corollary 3, each of these three considered subgraphs crosses T^{v_0} at least $\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ times. Hence, the edges of $C_n \cup T^{v_0}$ are crossed at least $3 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$ times. This is greater than $2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2$ for $n \geq 5$. Thus, $\text{cr}_D(T^{v_0}, C_n) = 1$ and there is exactly one another crossing on the edges of C_n . For the case $\text{cr}(S_5, C_n) = 1$, one subgraph T^{v_i} , say T^{v_1} , is placed in the other region of C_n as v_0 with $\text{cr}_D(T^{v_1}, C_n) = 0$. The drawing in Figure 3(a) can be generalized such that it confirms that, in this case, T^{v_1} crosses T^{v_0} at least once. If $\text{cr}(S_5, C_n) = 0$, one subgraph T^{v_i} , say again T^{v_1} , crosses C_n once. But it crosses also T^{v_0} . By Corollary 4, each of the remaining four subgraphs T^{v_i} , $i = 2, 3, 4, 5$, crosses T^{v_0} at least $\lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor$ times. Hence, on the edges of $C_n \cup T^{v_0}$ there are at least $4 \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor + 3$ crossings. This contradiction with at most $2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2$ crossings on the edges of $C_n \cup T^{v_0}$ completes the proof. ■

4. CARTESIAN PRODUCTS OF WHEELS AND TREES

In this section, using the results obtained in Section 3, we give the crossing number of the Cartesian products of wheels and trees. Let W_n denote the wheel of $n + 1$

vertices. In Figure 2 it is easy to verify that the graph $S_5 + C_n$ is isomorphic to the graph $W_n + D_5$. In general, the graph $S_m + C_n$ is isomorphic to the graph $W_n + D_m$ for all integers $m \geq 1$ and $n \geq 3$. So, we know the crossing numbers of the graph $W_n + D_m$ for all $m \leq 5$. Now, using zip product operation introduced in [4], we establish the crossing number of the Cartesian product $W_n \square T$ for all trees with maximum degree five. For better reading, we repeat the related terms, notations and results introduced by Bokal in [5].

For a graph G and a discrete graph D_m , let the vertices of D_m be called *apices* of the graph $G + D_m$. Consider now two graphs G_1 and G_2 . For a multiset $L \subseteq V(G_2)$, we denote with $G_1 \square_L G_2$ the *capped Cartesian product* of graphs G_1 and G_2 , that is, the graph obtained by adding a distinct vertex v' to $G_1 \square G_2$ for each copy of a vertex $v \in L$ and joining v' to all the vertices of $G_1 \square \{v\}$. We call each v' a *cap* of v . Let $\chi_L(v)$ denote the multiplicity of v in L and $\ell(v) := \deg_{G_2}(v) + \chi_L(v)$. An edge $\{u, v\} \in E(G_2)$ is *unbalanced* if $\ell(u) \neq \ell(v)$. Let $\beta(G_2)$ be the number of unbalanced edges of G_2 .

A drawing D of $G + D_m$ is *apex-homogeneous* if there exists a permutation ρ of the vertices of G such that the vertex rotation around every apex in D is ρ or ρ^{-1} . Two drawings $D^{(i)}$ of $G + D_i$ and $D^{(j)}$ of $G + D_j$ are *pairwise apex-homogeneous*, if they are apex-homogeneous with respect to the same permutation ρ . A graph G has *all apex-homogeneous drawings* if there exist drawings $D^{(m)}$ of $G + D_m$ for all $m \geq 1$, such that every two of them are pairwise apex-homogeneous. The next result given by Bokal enables us to establish the crossing numbers of $W_n \square T$.

Theorem 10 ([5]). *Let G be a graph of order n , let T be a tree, and let $L \subseteq V(T)$ be a multiset with either $\ell(v) \geq 3$ or, if G has a dominating vertex, $\ell(v) \geq 2$ for every $v \in V(T)$. Define*

$$B = \sum_{v \in V(T)} \text{cr}(G + D_{l(v)}).$$

Then, $B \leq \text{cr}(G \square_L T) \leq B + \frac{\beta(T)}{2} \binom{n}{2}$. Also, $\text{cr}(G \square_L T) = B$ whenever G has all apex-homogeneous drawings such that each of them is optimal.

Let $v \in V(G)$ be a vertex of degree d in G . A *bundle* B_v of the vertex v is a set of d edge disjoint paths from v to some other vertex $u \in V(G)$. Let $F \subset E(G)$ be a subset of edges of G and π a permutation of F . A π -subdivision G^π of G is the graph obtained from G by subdividing every edge $e \in F$ with the vertex v_e and adding the edges $\{\{v_e, v_{\pi(e)}\} | e \in F\}$.

Theorem 11 ([5]). *Let v be a vertex that has a bundle B_v in a graph G and let π be cyclic permutation of a subset of F of all but one of the edges incident with v , $|F| \geq 3$. Then*

$$\text{cr}(G^\pi) \geq \text{cr}(G) + 1,$$

with equality if π respects the edge rotation around v in some optimal drawing of G .

In the rest of the section, we give the crossing numbers of the Cartesian products of the wheel W_n and all trees T with maximum degree at most five.

Theorem 12. *Let T be a tree with maximum degree $\Delta(T) \leq 5$. Let d_i be the number of vertices of degree i in T . Then, for $n \geq 3$,*

$$\begin{aligned} \text{cr}(W_n \square T) = & d_1 + d_2 + 2d_3 + 2d_4 + 3d_5 + \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor (d_2 + 2d_3 + 4d_4 + 6d_5) \\ & + \left\lfloor \frac{n}{2} \right\rfloor (d_3 + 2d_4 + 4d_5). \end{aligned}$$

Proof. Let T' be the tree obtained from T by removing all vertices of degree one in T . For a vertex v of T' , let r_v be the number of T -leaves adjacent to v in T , and let L be the set of vertices in T' , each with multiplicity r_v . Thus, $\ell(v) = d_{T'}(v) + r_v = d_T(v) \geq 2$ for all $v \in V(T')$. Note that the central vertex of W_n is a dominating vertex, i.e., a vertex adjacent to all other vertices of the graph. Let us restrict on the graph $W_n + D_m$ for $m \leq 5$. The drawing in Figure 2 shows that, by this restriction, the wheel W_n has all apex-homogeneous drawings such that each of them is optimal. Thus, by Theorem 10, $\text{cr}(W_n \square_L T') = \sum_{i=2}^{\Delta(T)} (d_i \cdot \text{cr}(W_n + D_i))$.

The graph $W_n \square_L T'$ is obtained from the Cartesian product $W_n \square T'$ by adding r_v caps to $W_n \square \{v\}$ for every vertex v of T' with r_v T -leaves adjacent to v . This consistency in combination with Theorem 11 also implies that a properly chosen π -subdivision of edges connecting a cap of $W_n \square_L T'$ with the corresponding rim increases the crossing number by precisely one. Thus, after such π -subdivision for each of d_1 leaves of T , the graph $W_n \square T$ is obtained. This, together with the fact that $\text{cr}(W_n + D_1) = 1$, proves that $\text{cr}(W_n \square T) = \sum_{i=1}^{\Delta(T)} (d_i \cdot \text{cr}(W_n + D_i))$.

The values of crossing numbers of $W_n + D_m$ are known for all $m \leq 5$. In [5] Bokal proved that $\text{cr}(W_n + D_2) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 1$. As the graph $W_n + D_m$ is isomorphic to the graph $S_m + C_n$, by Theorems 7, 8, and 9 we have that $\text{cr}(W_n + D_3) = 2 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + \lfloor \frac{n}{2} \rfloor + 2$, $\text{cr}(W_n + D_4) = 4 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 2 \lfloor \frac{n}{2} \rfloor + 2$, and $\text{cr}(W_n + D_5) = 6 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor + 4 \lfloor \frac{n}{2} \rfloor + 3$ for $n \geq 3$. Applying this, together with the formula $\text{cr}(W_n \square T) = \sum_{i=1}^{\Delta(T)} (d_i \cdot \text{cr}(W_n + D_i))$ for $\Delta(T) \leq 5$, the result is done. ■

At present, the crossing numbers of very few join products of graphs with discrete graphs are known. In [21] there are collected the crossing numbers of $G + D_m$ for all graphs G of order at most four. Only some few results on crossing numbers of $G + D_m$ are known for graphs G on five and six vertices. Note that Theorem 10 cannot be simply used for estimating the crossing number of the Cartesian product $G \square T$ for all graphs G , for which the crossing number of the

join product $G + D_m$ is known. The reason is that not every graph contains a dominating vertex and, in this case, the vertices of degree two in T are not treatable using zip product.

Acknowledgement

The research was supported by the Slovak VEGA grant No. 1/0389/15.

REFERENCES

- [1] D. Archdeacon and R.B. Richter, *On the parity of crossing numbers*, J. Graph Theory **12** (1988) 307–310.
doi:10.1002/jgt.3190120302
- [2] K. Asano, *The crossing number of $K_{1,3,n}$ and $K_{2,3,n}$* , J. Graph Theory **10** (1986) 1–8.
doi:10.1002/jgt.3190100102
- [3] L.W. Beineke and R.D. Ringeisen, *On the crossing numbers of products of cycles and graphs of order four*, J. Graph Theory **4** (1980) 145–155.
doi:10.1002/jgt.3190040203
- [4] D. Bokal, *On the crossing numbers of Cartesian products with paths*, J. Combin. Theory Ser. B **97** (2007) 381–384.
doi:10.1016/j.jctb.2006.06.003
- [5] D. Bokal, *On the crossing numbers of Cartesian products with trees*, J. Graph Theory **56** (2007) 287–300.
doi:10.1002/jgt.20258
- [6] C. Buchheim, M. Chimani, D. Ebner, C. Gutwenger, M. Jnger, G.W. Klau, P. Mutzel and R. Weiskircher, *A branch-and-cut approach to the crossing number problem*, Discrete Optimization **5** (2008) 373–388.
doi:10.1016/j.disopt.2007.05.006
- [7] M. Chimani, P. Mutzel and I. Bomze, *A new approach to exact crossing minimization*, 16th Annual European Symposium on Algorithms 2008, Karlsruhe (ESA 08), Springer, Lecture Notes in Comput. Sci. **5193** (2008) 284–296.
doi:10.1007/978-3-540-87744-8.24
- [8] M. Chimani, C. Gutwenger and P. Mutzel, *Experiments on exact crossing minimization using column generation*, ACM J. Exp. Algorithmics **14** (2009) 4.1–4.18.
- [9] M. Chimani and T. Wiedera, *An ILP-based proof system for the crossing number problem*, 24th Annual European Symposium of Algorithms (ESA 2016), Aarhus, Denmark, Leibniz. Int. Proc. Inform. **57** (2016) 29.1–29.13.
doi:10.4230/LIPIcs.ESA.2016.29
- [10] L.Y. Glebsky and G. Salazar, *The crossing number of $C_m \times C_n$ is as conjectured for $n \geq m(m+1)$* , J. Graph Theory **47** (2004) 53–72.
doi:10.1002/jgt.20016

- [11] F. Harary, P.C. Kainen and A.J. Schwenk, *Toroidal graphs with arbitrarily high crossing numbers*, Nanta Math. **6** (1973) 58–67.
- [12] Y. Huang and T. Zhao, *The crossing number of $K_{1,4,n}$* , Discrete Math. **308** (2008) 1634–1638.
doi:10.1016/j.disc.2006.12.002
- [13] H. Mei and Y. Huang, *The crossing number of $K_{1,5,n}$* , Internat. J. Math. Combin. **1** (2007) 33–44.
- [14] S. Jendroľ and M. Ščerbová, *On the crossing numbers of $S_m \times P_n$ and $S_m \times C_n$* , Čas. Pěstování Mat. **107** (1982) 225–230.
- [15] M. Klešč, *The crossing numbers of products of paths and stars with 4-vertex graphs*, J. Graph Theory **18** (1994) 605–614.
doi:10.1002/jgt.3190180608
- [16] M. Klešč, *The crossing numbers of Cartesian products of paths with 5-vertex graphs*, Discrete Math. **233** (2001) 353–359.
doi:10.1016/S0012-365X(00)00251-X
- [17] M. Klešč, *The join of graphs and crossing numbers*, Electron. Notes Discrete Math. **28** (2007) 349–355.
doi:10.1016/j.endm.2007.01.049
- [18] M. Klešč, *The crossing numbers of join of the special graph on six vertices with path and cycle*, Discrete Mat. **310** (2010) 1475–1481.
doi:10.1016/j.disc.2009.08.018
- [19] M. Klešč and D. Kravecová, *The crossing numbers of $P_n^2 \square C_3$* , Discrete Mat. **312** (2012) 2096–2101.
doi:10.1016/j.disc.2011.03.028
- [20] M. Klešč and J. Petrillová, *On the optimal drawings of products of paths with graphs*, Acta Electrotechnica et Informatica **13** (2013) 56–61.
doi:10.2478/aei-2013-0041
- [21] M. Klešč and Š. Schrötter, *The crossing numbers of join products of paths with graphs of order four*, Discuss. Math. Graph Theory **31** (2011) 312–331.
doi:10.7151/dmgt.1548
- [22] M. Klešč and Š. Schrötter, *The crossing numbers of join of paths and cycles with two graphs of order five*, Mathematical Modeling and Computational Science, Springer, Lecture Notes in Comput. Sci. **7125** (2012) 160–167.
doi:10.1007/978-3-642-28212-6_15
- [23] M. Klešč and M. Valo, *Minimum crossings in join of graphs with paths and cycles*, Acta Electrotechnica et Informatica **12** (2012) 32–37.
doi:10.2478/v10198-012-0028-0
- [24] D.J. Kleitman, *The crossing number of $K_{5,n}$* , J. Combin. Theory Ser. B **9** (1971) 315–323.
doi:10.1016/S0021-9800(70)80087-4

- [25] D. Kravecová and J. Petrillová, *The crossing number of $P_n^2 \square C_4$* , Acta Electrotechnica et Informatica **12** (2012) 42–46.
doi:10.2478/v10198-012-0030-6
- [26] Z. Ouyang, W. Jing and Y. Huang, *On the crossing numbers of the Cartesian product path with complete graph*, Discrete Math. **328** (2014) 71–78.
doi:10.1016/j.disc.2014.03.011
- [27] W. Zheng, X. Lin, Y. Yang and C. Deng, *On the crossing numbers of $K_m \square C_n$ and $K_{m,t} \square P_n$* , Discrete Appl. Math. **156** (2008) 1899–1907.
doi:10.1016/j.dam.2007.09.007

Received 30 November 2015

Revised 9 November 2016

Accepted 9 November 2016