

DESCRIBING NEIGHBORHOODS OF 5-VERTICES
IN 3-POLYTOPES WITH MINIMUM DEGREE 5
AND WITHOUT VERTICES OF
DEGREES FROM 7 TO 11¹

OLEG V. BORODIN, ANNA O. IVANOVA

AND

OLESYA N. KAZAK

Institute of Mathematics Siberian Branch
Russian Academy of Sciences
Novosibirsk, 630090, Russia

e-mail: brdnoleg@math.nsc.ru
shmgnanna@mail.ru
agazandjelos@gmail.com

Abstract

In 1940, Lebesgue proved that every 3-polytope contains a 5-vertex for which the set of degrees of its neighbors is majorized by one of the following sequences:

(6, 6, 7, 7, 7), (6, 6, 6, 7, 9), (6, 6, 6, 6, 11),
(5, 6, 7, 7, 8), (5, 6, 6, 7, 12), (5, 6, 6, 8, 10), (5, 6, 6, 6, 17),
(5, 5, 7, 7, 13), (5, 5, 7, 8, 10), (5, 5, 6, 7, 27),
(5, 5, 6, 6, ∞), (5, 5, 6, 8, 15), (5, 5, 6, 9, 11),
(5, 5, 5, 7, 41), (5, 5, 5, 8, 23), (5, 5, 5, 9, 17),
(5, 5, 5, 10, 14), (5, 5, 5, 11, 13).

In this paper we prove that every 3-polytope without vertices of degree from 7 to 11 contains a 5-vertex for which the set of degrees of its neighbors is majorized by one of the following sequences: (5, 5, 6, 6, ∞), (5, 6, 6, 6, 15), (6, 6, 6, 6, 6), where all parameters are tight.

Keywords: planar graph, structure properties, 3-polytope, neighborhood.

2010 Mathematics Subject Classification: 05C15.

¹The work was funded by the Russian Science Foundation, grant 16–11–10054.

1. INTRODUCTION

By a *3-polytope* we mean a finite 3-dimensional convex polytope. As proved by Steinitz [31], the 3-polytopes are in one to one correspondence with the 3-connected planar graphs.

The *degree* $d(v)$ of a vertex v ($r(f)$ of a face f) in a 3-polytope P is the number of edges incident with it. By Δ and δ we denote the maximum and minimum vertex degrees of P , respectively. A k -*vertex* (k -*face*) is a vertex (face) with degree k ; a k^+ -*vertex* has degree at least k , etc.

The *weight* of a face in P is the degree sum of its boundary vertices, and $w(P)$, or simply w , denotes the minimum weight of 5^- -faces in P .

In 1904, Wernicke [32] proved that every 3-polytope with $\delta = 5$ has a 5-vertex adjacent with a 6^- -vertex, which was strengthened by Franklin [15] in 1922, who proved that every 3-polytope with $\delta = 5$ has a 5-vertex adjacent with two 6^- -vertices. Recently, Borodin and Ivanova [11] proved that every such 3-polytope has also a vertex of degree at most 6 adjacent to a 5-vertex and another vertex of degree at most 6, which is tight.

We say that v is a *vertex of type* (k_1, k_2, \dots) or simply a (k_1, k_2, \dots) -*vertex* if the set of degrees of the vertices adjacent to v is majorized by the vector (k_1, k_2, \dots) . If the order of neighbors in the type is not important, then we put a line over the corresponding degrees. The following description of the neighborhoods of 5-vertices in a 3-polytope with $\delta = 5$ was given by Lebesgue [28, p. 36] in 1940, which includes the results of Wernicke [32] and Franklin [15].

Theorem 1 (Lebesgue [28]). *Every triangulated 3-polytope with minimum degree 5 contains a 5-vertex of one of the following types:*

$$\begin{aligned} & (\overline{6, 6, 7, 7, 7}), (\overline{6, 6, 6, 7, 9}), (\overline{6, 6, 6, 6, 11}), \\ & (\overline{5, 6, 7, 7, 8}), (\overline{5, 6, 6, 7, 11}), (\overline{5, 6, 6, 8, 8}), \\ & (5, 6, \overline{6, 9, 7}), (5, 7, 6, 6, 12), (5, 8, 6, 6, 10), (5, 6, 6, 6, 17), \\ & (5, 5, \overline{7, 7, 8}), (5, 13, 5, 7, 7), (5, 10, 5, 7, 8), \\ & (5, 8, 5, 7, 9), (5, 7, 5, 7, 10), (5, 7, 5, 8, 8), \\ & (5, 5, 7, 6, 12), (5, 5, 8, 6, 10), (5, 6, 5, 7, 12), \\ & (5, 6, 5, 8, 10), (5, 17, 5, 6, 7), (5, 11, 5, 6, 8), \\ & (5, 11, 5, 6, 9), (5, 7, 5, 6, 13), (5, 8, 5, 6, 11), (5, 9, 5, 6, 10), (5, 6, 6, 5, \infty), \\ & (5, 5, 7, 5, 41), (5, 5, 8, 5, 23), (5, 5, 9, 5, 17), (5, 5, 10, 5, 14), (5, 5, 11, 5, 13). \end{aligned}$$

Theorem 1, along with other ideas in Lebesgue [28], has many applications to plane graph coloring problems (first examples of such applications and a recent survey can be found in [7, 30]). Some parameters of Lebesgue's Theorem were improved for narrow classes of plane graphs. For example, in 1963, Kotzig [27] proved that every plane triangulation with $\delta = 5$ satisfies $w \leq 18$ and conjectured

that $w \leq 17$. In 1989, Kotzig's conjecture was confirmed by Borodin [3] in a more general form.

Theorem 2 (Borodin [3]). *Every 3-polytope with $\delta = 5$ has a $(5, 5, 7)$ -face or a $(5, 6, 6)$ -face, where all parameters are tight.*

By a *minor k -star* $S_k^{(m)}$ we mean a star with k rays centered at a 5^- -vertex. The Lebesgue's description [28, p.36] of the neighborhoods of 5-vertices in 3-polytopes with minimum degree 5, \mathbf{P}_5 , shows that there is a 5-vertex with three 8^- -neighbors. Another corollary of Lebesgue's description [28] is that $w(S_3^{(m)}) \leq 24$, which was improved in 1996 by Jendrol' and Madaras [23] to the sharp bound $w(S_3^{(m)}) \leq 23$. Furthermore, Jendrol' and Madaras [23] gave a precise description of minor 3-stars in \mathbf{P}_5 : there is a $(6, 6, 6)$ - or $(5, 6, 7)$ -star.

Also, Lebesgue [28] proved that $w(S_4^{(m)}) \leq 31$, which was strengthened by Borodin and Woodall [13] to the sharp bound $w(S_4^{(m)}) \leq 30$. Note that $w(S_3^{(m)}) \leq 23$ easily implies $w(S_2^{(m)}) \leq 17$ and immediately follows from $w(S_4^{(m)}) \leq 30$ (in both cases, it suffices to delete a vertex of maximum degree from a minor star of minimum weight). In [9], Borodin and Ivanova obtained a tight description of minor 4-stars in \mathbf{P}_5 .

As for minor 5-stars in \mathbf{P}_5 , it follows from Lebesgue [28, p. 36] that if there are no minor $(5, 5, 6, 6)$ -stars, then $w(S_5^{(m)}) \leq 68$ and $h(S_5^{(m)}) \leq 41$. Borodin, Ivanova, and Jensen [10] showed that the presence of minor $(5, 5, 6, 6)$ -stars can make $w(S_5^{(m)})$ arbitrarily large and otherwise lowered Lebesgue's bounds to $w(S_5^{(m)}) \leq 55$ and $h(S_5^{(m)}) \leq 28$. On the other hand, a construction in [10] shows that $w(S_5^{(m)}) \geq 48$ and $h(S_5^{(m)}) \geq 20$. Recently, Borodin and Ivanova [12] proved that $w(S_5^{(m)}) \leq 51$ and $h(S_5^{(m)}) \leq 23$.

More results on the structure of edges and higher stars in various classes of 3-polytopes can be found in [1, 2, 4–6, 8, 9, 14, 16, 19–22, 24–26], with a detailed summary in [12].

In [28] Lebesgue did not give a proof of Theorem 1 and only gave its idea. In 2013, Ivanova and Nikiforov [17] gave a full proof of Theorem 1 and corrected the following imprecisions in its statement:

- (1) in the type $(5, 11, 5, 6, 8)$ there should be 15 instead of 11;
- (2) in the type $(5, 17, 5, 6, 7)$ there should be 27 instead of 17;
- (3) in the type $(\overline{6, 6, 6, 6, 11})$ the line is not needed;
- (4) instead of type $(\overline{5, 6, 7, 7, 8})$ there should be $(5, 8, \overline{6, 7, 7})$ and $(5, 7, 6, 8, 7)$;

- (5) the type $(5, 6, \overline{6, 9}, 7)$ is redundant;
 (6) instead of $(5, 5, \overline{7, 7}, 8)$ it suffices to write $(5, 5, 7, \overline{7, 8})$.

Later on, Ivanova and Nikiforov [18, 29] improved the corrected version of Theorem 1 by replacing 41 and 23 in the types $(5, 5, 7, 5, 41)$ and $(5, 5, 8, 5, 23)$ to 31 and 22, respectively.

Theorem 3 (Ivanova, Nikiforov [17, 18, 29]). *Every 3-polytope with minimum degree 5 contains a 5-vertex of one of the following types:*

$$\begin{aligned} & (\overline{6, 6, 7, 7, 7}), (\overline{6, 6, 6, 7, 9}), (6, 6, 6, 6, 11), \\ & (5, 8, \overline{6, 7, 7}), (5, 7, 6, 8, 7), (5, 6, \overline{6, 7}, 11), (5, 6, \overline{6, 8}, 8), \\ & (5, 7, 6, 6, 12), (5, 8, 6, 6, 10), (5, 6, 6, 6, 17), \\ & (5, 5, 7, \overline{7, 8}), (5, 13, 5, 7, 7), (5, 10, 5, 7, 8), (5, 8, 5, 7, 9), \\ & (5, 7, 5, 7, 10), (5, 7, 5, 8, 8), (5, 5, 7, 6, 12), (5, 5, 8, 6, 10), \\ & (5, 6, 5, 7, 12), (5, 6, 5, 8, 10), (5, 27, 5, 6, 7), (5, 15, 5, 6, 8), \\ & (5, 11, 5, 6, 9), (5, 7, 5, 6, 13), (5, 8, 5, 6, 11), (5, 9, 5, 6, 10), \\ & (5, 6, 6, 5, \infty), \\ & (5, 5, 7, 5, 31), (5, 5, 8, 5, 22), (5, 5, 9, 5, 17), (5, 5, 10, 5, 14), (5, 5, 11, 5, 13). \end{aligned}$$

Theorem 1 subject to the corrections (1)–(6) implies the following fact.

Corollary 4. *Every 3-polytope with minimum degree 5 contains a 5-vertex of one of the following types:*

$$\begin{aligned} & (\overline{6, 6, 7, 7, 7}), (\overline{6, 6, 6, 7, 9}), (\overline{6, 6, 6, 6, 11}), \\ & (\overline{5, 6, 7, 7, 8}), (\overline{5, 6, 6, 7, 12}), (\overline{5, 6, 6, 8, 10}), (\overline{5, 6, 6, 6, 17}), \\ & (\overline{5, 5, 7, 7, 13}), (\overline{5, 5, 7, 8, 10}), (\overline{5, 5, 6, 7, 27}), \\ & (\overline{5, 5, 6, 6, \infty}), (\overline{5, 5, 6, 8, 15}), (\overline{5, 5, 6, 9, 11}), \\ & (\overline{5, 5, 5, 7, 41}), (\overline{5, 5, 5, 8, 23}), (\overline{5, 5, 5, 9, 17}), (\overline{5, 5, 5, 10, 14}), (\overline{5, 5, 5, 11, 13}). \end{aligned}$$

We can see already from Theorem 1 that if vertices of degree from 7 to 11 are forbidden, then there is a 5-vertex of one of the following types: $(\overline{5, 5, 6, 6, \infty})$, $(\overline{5, 6, 6, 6, 17})$, $(6, 6, 6, 6, 6)$.

The purpose of this note is to obtain a precise description of 5-stars in this subclass of \mathbf{P}_5 .

Theorem 5. *Every 3-polytope with minimum degree 5 and without vertices of degree from 7 to 11 contains a 5-vertex of one of the following types: $(\overline{5, 5, 6, 6, \infty})$, $(\overline{5, 6, 6, 6, 15})$, $(6, 6, 6, 6, 6)$, where all parameters are tight.*

2. PROVING THEOREM 5

All parameters in Theorem 5 are best possible. Indeed, the following construction confirming the tightness of the type $(\overline{5, 5, 6, 6, \infty})$ appears in [10]. Take three

concentric n -cycles $C^i = v_1^i \cdots v_n^i$, where n is not limited and $1 \leq i \leq 3$, and join C^2 with C^1 by edges $v_j^2 v_j^1$ and $v_j^2 v_{j+1}^1$, where $1 \leq j \leq n$ (addition modulo n). Then do the same with C^2 and C^3 . Finally, join all vertices of C^1 with a new n -vertex, and do the same for C^3 .

The tightness of $(6, 6, 6, 6, 6)$ is confirmed by putting a 5-vertex in each face of the dodecahedron.

To confirm the tightness of $(5, 6, 6, 6, 15)$, we take the dodecahedron and insert the fragment shown in Figure 1 into each face. As a result, we have a 3-polytope with only $(5, 6, 6, 6, 15)$ -vertices.




Figure 1. The insert in each face of the dodecahedron to produce a 3-polytope with 5-vertices only of type $(5, 6, 6, 6, 15)$.

Now suppose a 3-polytope P' is a counterexample to Theorem 5. Let P be a counterexample on the same number of vertices with maximum possible number of edges.

Remark 6. In P , each 4^+ -face $f = v_1 \cdots v_{d(f)}$ with $d(v_1) = 5$ or $d(v_1) \geq 15$ satisfies $d(v_i) \geq 6$ whenever $3 \leq i \leq d(f) - 1$. Otherwise, we could put a diagonal $v_1 v_i$, which contradicts the maximality of P .

Corollary 7. In P , each 4^+ -face has at most two vertices with degree 5 and/or at least 15. Moreover, if there are precisely two such vertices, then they are adjacent to each other.

2.1. Discharging

The sets of vertices, edges, and faces of P are denoted by V , E , and F , respectively. Euler's formula $|V| - |E| + |F| = 2$ for P implies

$$(1) \quad \sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2r(f) - 6) = -12.$$

We assign an *initial charge* $\mu(v) = d(v) - 6$ to every vertex v and $\mu(f) = 2d(f) - 6$ to every face f , so that only 5^- -vertices have negative charge. Using the properties of P as a counterexample, we define a local redistribution of charges, preserving their sum, such that the *new charge* $\mu'(x)$ is non-negative whenever $x \in V \cup F$. This will contradict the fact that the sum of the new charges is, by (1), equal to -12 . The technique of discharging is often used in solving structural and coloring problems on plane graphs.

Let $v_1, \dots, v_{d(v)}$ denote the neighbors of a vertex v in a cyclic order round v , and let $f_1, \dots, f_{d(v)}$ be the faces incident with v in the same order.

We use the following rules of discharging (see Figure 2).




Figure 2. Rules of discharging.

R1. Every 4^+ -face gives 1 to every incident 5 -vertex.

R2. Every 12^+ -vertex v gives a simplicial 5 -vertex v_2 the following charge through a face $f = v_2vv_3$:

- (a) $\frac{1}{4}$ if $d(v_3) = 5$,

- (b) $\frac{1}{2}$ if $d(v_3) \geq 6$,
with the following exception.
- (e) If $d(v) \geq 16$, $d(v_1) = 5$, $d(v_3) = d(x) = d(y) = 6$, where v_2 is incident to face v_2xy , then v gives $\frac{2}{3}$ to v_2 through face v_2vv_3 and $\frac{1}{3}$ through face v_1vv_2 .

R3. Suppose a simplicial 5-vertex v is adjacent to a 16-vertex v_1 , simplicial 5-vertices v_2 and v_5 , and v_2 is surrounded by v_1, v, v_3, x, y , where $d(v_3) = d(x) = d(y) = 6$, (consequently $d(v_4) \geq 12$), while v_5 is surrounded by v_1, v, v_4, w, z , where $d(z) \geq 6$. Then v gives $\frac{1}{4}$ to v_1 .

2.2. Proving $\mu'(x) \geq 0$ whenever $x \in V \cup F$

First consider a face f in P . If $d(f) = 3$, then f does not participate in discharging, and so $\mu'(v) = \mu(f) = 2 \times 3 - 6 = 0$. Note that every 4^+ -face is incident with at most two 5-vertices due to Corollary 7, which implies that $\mu'(v) = 2d(f) - 6 - 2 \times 1 \geq 0$ by R1.

Now let v be a vertex in P .

Case 1. $d(v) = 5$. If v is incident with a 4^+ -face, then $\mu'(v) \geq 5 - 6 + 1 = 0$ due to R1. In what follows we can assume that v is simplicial.

Subcase 1.1. v is incident only with 6^+ -vertices. Then there is at least one v_i with $d(v_i) \geq 12$ due to the absence of $(6, 6, 6, 6, 6)$ -vertices in P . Hence, $\mu'(v) \geq -1 + 2 \times \frac{1}{2} = 0$ by R2(b).

Subcase 1.2. v is incident with precisely one 5-vertex. Since there is no $(5, 6, 6, 6, 15)$ -vertex in P , we can assume that v has either at least two 12^+ -neighbors, or precisely one 16^+ -neighbor. So we have either $\mu'(v) \geq -1 + 2 \times \frac{1}{2} + 2 \times \frac{1}{4} > 0$ by R2(a),(b), or $\mu'(v) = -1 + \frac{2}{3} + \frac{1}{3} = 0$ by R2(e), respectively.

Subcase 1.3. v is incident with at least two 5-vertices. Note that now R2(e) is not applicable to v . Also note that v cannot be incident with more than three 5-vertices due to the absence of $(5, 5, 6, 6, \infty)$ -vertices in P , which implies that v has at least two 12^+ -neighbors. If v is incident with precisely three 5-vertices, then we have $\mu'(v) \geq -1 + 4 \times \frac{1}{4} = 0$ by R2(a),(b).

Suppose v is incident with precisely two 5-vertices. If v does not participate in R3, then $\mu'(v) \geq -1 + 3 \times \frac{1}{4} + \frac{1}{2} > 0$ by R2(a),(b). Note that if v participates in R3, then it gives $\frac{1}{4}$ only to one 16-neighbor, hence $\mu'(v) \geq -1 + 3 \times \frac{1}{4} + \frac{1}{2} - \frac{1}{4} = 0$.

Case 2. $d(v) = 6$. Since v does not participate in discharging, we have $\mu'(v) = \mu'(v) = 6 - 6 = 0$.

Case 3. $12 \leq d(v) \leq 15$. Now R2(e) is not applicable to v , so v sends at most $\frac{1}{2}$ through each face by R2(a),(b), which implies that $\mu'(v) \geq d(v) - 6 - d(v) \times \frac{1}{2} = \frac{d(v)-12}{2} \geq 0$.

Case 4. $16 \leq d(v) \leq 17$. Note that v gives at most $\frac{2}{3}$ through each 3-face and only to a simplicial 5-vertex. If v gives nothing through at least one incident face, then $\mu'(v) \geq 16 - 6 - 15 \times \frac{2}{3} = 0$ by R1, R2. Further, we can assume that v is simplicial and each face takes away some positive charge from v , which implies that each face at v is incident with a 5-vertex, and all 5-vertices adjacent to v are simplicial. Thus, $\mu'(v) \geq d(v) - 6 - d(v) \times \frac{2}{3} = \frac{d(v)-18}{3}$, and we have the deficiency $\frac{1}{3}$ for a 17-vertex and $\frac{2}{3}$ for a 16-vertex with respect to donating $\frac{2}{3}$ per face.

Suppose $S_k = v_1, \dots, v_k$ is a sequence of neighbors of v with $d(v_1) \geq 6$, $d(v_k) \geq 6$, while $d(v_i) = 5$ whenever $2 \leq i \leq k-1$ and $k \geq 3$, and f_1, \dots, f_{k-1} are the corresponding faces. (It is not excluded that $S_k = S_{d(v)}$, which happens when v has precisely one 6^+ -neighbor.) We say that the sequence of faces f_1, \dots, f_{k-1} *saves* ε with respect to the level of $\frac{2}{3}$ if these faces take away the total of $(k-1) \times \frac{2}{3} - \varepsilon$ from v .

Remark 8. Only v_2 and v_{k-1} in S_k can receive the charge $\frac{2}{3}$ from v by R2(e), while each of the other 5-vertices v_i receives precisely $\frac{1}{4}$ from v through each incident face. So, if $k \geq 5$, then v_2 receives at most 1, and v_3 receives $\frac{1}{2}$ from v through incident faces.

Remark 9. If v is completely surrounded by 5-vertices, then $\mu'(v) \geq d(v) - 6 - \frac{d(v)}{2} = \frac{d(v)-12}{2} > 0$, and hence we can assume from now on that the neighborhood of v is partitioned into S_k s.

(P1) If $k = 3$, then $\varepsilon = \frac{1}{3}$. Indeed, here v_2 receives $\frac{1}{2}$ through each of the faces $v_1v_2v_3$ and $v_2v_3v_1$ by R2(b), whence $\varepsilon = 2 \times \frac{2}{3} - 2 \times \frac{1}{2} = \frac{1}{3}$.

(P2) If $k = 4$, then $\varepsilon = 0$. Now each of v_2 and v_3 receives at most 1 from v by Remark 8, so $\varepsilon = 3 \times \frac{2}{3} - 2 = 0$.

(P3) If $k = 5$, then $\varepsilon = \frac{2}{3}$. Suppose w_1, \dots, w_4 are the neighbors of v_1, \dots, v_5 such that there are the faces $v_iw_iv_{i+1}$, where $1 \leq i \leq 4$.

If v_2 receives 1 by R2(e), then $d(w_1) = d(w_2) = 6$. Hence, $d(w_3) \geq 12$ due to the absence of a $(5, 5, 6, 6, \infty)$ -vertex in P , which implies that w_4 is adjacent to two 12^+ -vertices, whence it receives $\frac{1}{2}$ from v through f_4 and $\frac{1}{4}$ through f_3 . Moreover, v_3 gives $\frac{1}{4}$ to v by R3. Hence, $\varepsilon = 4 \times \frac{2}{3} - 1 - \frac{1}{2} - \frac{3}{4} + \frac{1}{4} = \frac{2}{3}$.

If R2(e) is not applicable to v , then $\varepsilon = 4 \times \frac{2}{3} - 4 \times \frac{1}{2} = \frac{2}{3}$.

(P4) If $k = 6$, then $\varepsilon = \frac{1}{3}$. Here, each of v_2 and v_5 receives at most 1, while each of v_3 and v_4 receives $\frac{1}{2}$ from v by Remark 8, so $\varepsilon = 5 \times \frac{2}{3} - 2 \times 1 - 2 \times \frac{1}{2} = \frac{1}{3}$.

(P5) If $k = 7$, then $\varepsilon = \frac{1}{2}$. Now we have $\varepsilon = 6 \times \frac{2}{3} - 2 \times 1 - 3 \times \frac{1}{2} = \frac{1}{2}$ by Remark 8.

(P6) If $k \geq 8$, then $\varepsilon \geq \frac{2}{3}$. Now we have $\varepsilon = (k-1) \times \frac{2}{3} - 2 \times 1 - (k-4) \times \frac{1}{2} = \frac{k-4}{6} \geq \frac{2}{3}$.

If $d(v) = 17$, then it suffices to assume that the neighborhood of v consists of pairs of 5-vertices separated from each other by 6^+ -vertices by (P1)–(P6) (since otherwise we pay off the deficiency), which is impossible due to the fact that 17 is not divisible by 3.

Suppose that $d(v) = 16$ and $\mu'(v) < 0$. As follows from (P1)–(P6), the neighborhood of v can have at most one of the paths S_{t+2} of t vertices of degree 5, where $t \in \{1, 4, 5\}$, while all other vertices are partitioned into pairs of 5-vertices separated from each other by 6-vertices. Indeed, if there are either two paths with $t \in \{1, 4, 5\}$, or at least one path with $t = 3$ or $t \geq 6$, then we can pay off the deficiency $\frac{2}{3}$, a contradiction. But none of these cases is possible due to the divisibility by 3. Namely, if $t = 1$ we have $16 - 2 = 14$ faces to be divided into triplets of faces with a sequence S_4 of neighbors of v as in (P2), or $16 - 5 = 11$ and $16 - 6 = 10$ faces for $t = 4$ and $t = 5$, respectively; a contradiction.

Case 6. $d(v) \geq 18$. Now $\mu'(v) \geq d(v) - 6 - d(v) \times \frac{2}{3} = \frac{d(v)-18}{3} \geq 0$ by R2.

Thus we have proved $\mu'(x) \geq 0$ for every $x \in V \cup F$, which contradicts (1) and completes the proof of Theorem 5.

REFERENCES

- [1] V.A. Aksenov, O.V. Borodin and A.O. Ivanova, *Weight of 3-paths in sparse plane graphs*, Electron. J. Combin. **22** (2015) #P3.28.
- [2] K. Ando, S. Iwasaki and A. Kaneko, *Every 3-connected planar graph has a connected subgraph with small degree sum*, Annual Meeting of Mathematical Society of Japan (1993).
- [3] O.V. Borodin, *Solution of Kotzig's and Grünbaum's problems on the separability of a cycle in a planar graph*, Mat. Zametki **46** (1989) 9–12, in Russian.
- [4] O.V. Borodin, *Joint generalization of the theorems of Lebesgue and Kotzig on the combinatorics of planar maps*, Diskret. Mat. **3** (1991) 24–27, in Russian.
- [5] O.V. Borodin, *Joint extension of two Kotzig's theorems on 3-polytopes*, Combinatorica **13** (1993) 121–125.
doi:10.1007/BF01202794
- [6] O.V. Borodin, *Minimal vertex degree sum of a 3-path in plane maps*, Discuss. Math. Graph Theory **17** (1997) 279–284.
doi:10.7151/dmgt.1055
- [7] O.V. Borodin, *Colorings of plane graphs: a survey*, Discrete Math. **313** (2013) 517–539.
doi:10.1016/j.disc.2012.11.011

- [8] O.V. Borodin and A.O. Ivanova, *Describing $(d - 2)$ -stars at d -vertices, $d \leq 5$, in normal plane maps*, Discrete Math. **313** (2013) 1700–1709.
doi:10.1016/j.disc.2013.04.026
- [9] O.V. Borodin and A.O. Ivanova, *Describing 4-stars at 5-vertices in normal plane maps with minimum degree 5*, Discrete Math. **313** (2013) 1710–1714.
doi:10.1016/j.disc.2013.04.025
- [10] O.V. Borodin, A.O. Ivanova and T.R. Jensen, *5-stars of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **34** (2014) 539–546.
doi:10.7151/dmgt.1748
- [11] O.V. Borodin and A.O. Ivanova, *An analogue of Franklin's Theorem*, Discrete Math. **339** (2016) 2553–2556.
doi:10.1016/j.disc.2016.04.019
- [12] O.V. Borodin and A.O. Ivanova, *Light and low 5-stars in normal plane maps with minimum degree 5*, Sibirsk. Mat. Zh. **57** (2016) 596–602, in Russian.
doi:10.1134/S0037446616030071
- [13] O.V. Borodin and D.R. Woodall, *Short cycles of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **18** (1998) 159–164.
doi:10.7151/dmgt.1071
- [14] B. Ferencová and T. Madaras, *Light graph in families of polyhedral graphs with prescribed minimum degree, face size, edge and dual edge weight*, Discrete Math. **310** (2010) 1661–1675.
doi:10.1016/j.disc.2009.11.027
- [15] Ph. Franklin, *The four colour problem*, Amer. J. Math. **44** (1922) 225–236.
doi:10.2307/2370527
- [16] J. Harant and S. Jendrol', *On the existence of specific stars in planar graphs*, Graphs Combin. **23** (2007) 529–543.
doi:10.1007/s00373-007-0747-7
- [17] A.O. Ivanova and D.V. Nikiforov, *The structure of neighborhoods of 5-vertices in plane triangulation with minimum degree 5*, Mat. Zam. YaSU **20(2)** (2013) 66–78, in Russian.
- [18] A.O. Ivanova and D.V. Nikiforov, *Combinatorial structure of triangulated 3-polytopes with minimum degree 5*, in: Proceedings of the Scientific Conference of students, graduate students, and young researchers. XVII and XVIII Lavrent'ev's reading, Yakutsk; Kirov: International Center for Research Project (2015) 22–27, in Russian.
- [19] S. Jendrol', *A structural property of convex 3-polytopes*, Geom. Dedicata **68** (1997) 91–99.
doi:10.1023/A:1004993723280
- [20] S. Jendrol', *Paths with restricted degrees of their vertices in planar graphs*, Czechoslovak Math. J. **49** (1999) 481–490.
doi:10.1023/A:1022411100562

- [21] S. Jendrol' and M. Maceková, *Describing short paths in plane graphs of girth at least 5*, Discrete Math. **338** (2015) 149–158.
doi:10.1016/j.disc.2014.09.014
- [22] S. Jendrol', M. Maceková and R. Soták, *Note on 3-paths in plane graphs of girth 4*, Discrete Math. **338** (2015) 1643–1648.
doi:10.1016/j.disc.2015.04.011
- [23] S. Jendrol' and T. Madaras, *On light subgraphs in plane graphs of minimal degree five*, Discuss. Math. Graph Theory **16** (1996) 207–217.
doi:10.7151/dmgt.1035
- [24] S. Jendrol' and T. Madaras, *Note on an existence of small degree vertices with at most one big degree neighbour in planar graphs*, Tatra Mt. Math. Publ. **30** (2005) 149–153.
- [25] S. Jendrol' and H.-J. Voss, *Light subgraphs of graphs embedded in the plane—a survey*, Discrete Math. **313** (2013) 406–421.
doi:10.1016/j.disc.2012.11.007
- [26] A. Kotzig, *Contribution to the theory of Eulerian polyhedra*, Mat. Eas. SAV (Math. Slovaca) **5** (1955) 101–113.
- [27] A. Kotzig, *From the theory of Eulerian polyhedra*, Mat. Eas. SAV (Math. Slovaca) **13** (1963) 20–34, in Russian.
- [28] H. Lebesgue, *Quelques conséquences simples de la formule d'Euler*, J. Math. Pures Appl. **19** (1940) 27–43.
- [29] D.V. Nikiforov, *The structure of neighborhoods of 5-vertices in normal plane maps with minimum degree 5*, Mat. Zam. YaSU, **23** (2016) 56–66, in Russian.
- [30] O. Ore and M.D. Plummer, *Cyclic coloration of plane graphs*, in: Recent Progress in Combinatorics, W.T. Tutte, (Ed.), (Academic Press, New York, 1969) 287–293.
- [31] E. Steinitz, *Polyeder und Raumeinteilungen*, Enzykl. Math. Wiss. (Geometrie), 3AB **12** (1922) 1–139.
- [32] P. Wernicke, *Über den kartographischen Vierfarbensatz*, Math. Ann. **58** (1904) 413–426.
doi:10.1007/BF01444968

Received 12 July 2016

Revised 13 January 2017

Accepted 13 January 2017