

Discussiones Mathematicae
Graph Theory 38 (2018) 717–741
doi:10.7151/dmgt.2046

TURÁN FUNCTION AND H -DECOMPOSITION PROBLEM FOR GEM GRAPHS

HENRY LIU

School of Mathematics and Statistics
Central South University
Changsha 410083, China

e-mail: henry-liu@csu.edu.cn

AND

TERESA SOUSA

Escola Naval and Centro de Investigação Naval
Escola Naval - Alfeite
2810-001 Almada, Portugal

and

Centro de Matemática e Aplicações
Faculdade de Ciências e Tecnologia
Universidade Nova de Lisboa
Campus de Caparica
2829-516 Caparica, Portugal

e-mail: teresa.maria.sousa@marinha.pt

Abstract

Given a graph H , the *Turán function* $\text{ex}(n, H)$ is the maximum number of edges in a graph on n vertices not containing H as a subgraph. For two graphs G and H , an H -*decomposition* of G is a partition of the edge set of G such that each part is either a single edge or forms a graph isomorphic to H . Let $\phi(n, H)$ be the smallest number ϕ such that any graph G of order n admits an H -decomposition with at most ϕ parts. Pikhurko and Sousa conjectured that $\phi(n, H) = \text{ex}(n, H)$ for $\chi(H) \geq 3$ and all sufficiently large n . Their conjecture has been verified by Özkahya and Person for all edge-critical graphs H . In this article, we consider the *gem graphs* gem_4 and gem_5 . The graph gem_4 consists of the path P_4 with four vertices a, b, c, d and edges ab, bc, cd plus a universal vertex u adjacent to a, b, c, d , and the graph gem_5 is similarly defined with the path P_5 on five vertices. We determine

the Turán functions $\text{ex}(n, \text{gem}_4)$ and $\text{ex}(n, \text{gem}_5)$, and verify the conjecture of Pikhurko and Sousa when H is the graph gem_4 and gem_5 .

Keywords: gem graph, Turán function, extremal graph, graph decomposition.

2010 Mathematics Subject Classification: 05C35, 05C70.

1. INTRODUCTION

Given a graph H , the *Turán function* $\text{ex}(n, H)$ is the maximum number of edges in a graph on n vertices, and not containing a copy of H as a subgraph. The important result of Turán [13] states that when $H = K_r$ is the complete graph on $r \geq 3$ vertices, we have $\text{ex}(n, K_r) = t_{r-1}(n)$. Here $t_{r-1}(n)$ denotes the number of edges in the *Turán graph* of order n , $T_{r-1}(n)$, which is the unique complete $(r-1)$ -partite graph on n vertices where every partition class has either $\lfloor \frac{n}{r-1} \rfloor$ or $\lceil \frac{n}{r-1} \rceil$ vertices. Moreover, $T_{r-1}(n)$ is the unique extremal graph on n vertices that has the maximum number of edges not containing K_r as a subgraph. For general graphs H , the Turán function $\text{ex}(n, H)$ has been well studied by numerous researchers, which led to many important results and open problems in extremal graph theory. For example, when $H = C_{2k}$ is the even cycle of length $2k$, where $k \geq 2$, the exact determination of the function $\text{ex}(n, C_{2k})$ is still a wide open problem. It has been conjectured that $\text{ex}(n, C_{2k}) = (c_k + o(1))n^{1+1/k}$ for some constant $c_k > 0$, and this conjecture is only known to be true for $k = 2, 3, 5$. See for example [8] and the references therein. When $H = P_k$ is the path of order $k \geq 3$, Faudree and Schelp [5] have determined the function $\text{ex}(n, P_k)$ exactly. In order to obtain $\text{ex}(n, P_k)$, we can take the graph on n vertices containing as many disjoint copies of K_{k-1} as possible, and a smaller complete graph on the remaining vertices. For odd k , this graph is the unique P_k -free extremal graph attaining $\text{ex}(n, P_k)$, and for even k and certain values of n , there are other such extremal graphs. Here we state the result of Faudree and Schelp as follows, which will be useful in this paper.

Theorem 1.1 [5]. *Let $k \geq 3$ and $n = a(k-1) + b$, where $a \geq 0$ and $0 \leq b < k-1$. Then $\text{ex}(n, P_k) = a \binom{k-1}{2} + \binom{b}{2}$. Moreover, a P_k -free graph on n vertices attaining $\text{ex}(n, P_k)$ is $aK_{k-1} \cup K_b$, the disjoint union of a copies of K_{k-1} and one copy of K_b .*

For two graphs G and H , an H -decomposition of G is a partition of the edge set of G such that each part is either a single edge or forms a graph isomorphic to H . Let $\phi(G, H)$ be the smallest possible number of parts in an H -decomposition of G . It is easy to see that, for non-empty H , we have $\phi(G, H) = e(G) -$

$p_H(G)(e(H) - 1)$, where $p_H(G)$ is the maximum number of pairwise edge-disjoint copies of H that can be packed into G and $e(G)$ denotes the number of edges in G . Dor and Tarsi [3] showed that if H has a component with at least three edges, then the problem of checking whether a graph G admits a partition into H -subgraphs is NP-complete. Thus, it is NP-hard to compute the function $\phi(G, H)$ for such H . Here we study the function

$$\phi(n, H) = \max\{\phi(G, H) \mid v(G) = n\},$$

which is the smallest number ϕ such that any graph G of order n admits an H -decomposition with at most ϕ parts.

This function was first studied, in 1966, by Erdős, Goodman and Pósa [4], who were motivated by the problem of representing graphs by set intersections. They proved that $\phi(n, K_3) = t_2(n)$. A decade later, this result was extended by Bollobás [2], who proved that $\phi(n, K_r) = t_{r-1}(n)$, for all $n \geq r \geq 3$.

General graphs H were only considered recently by Pikhurko and Sousa [9]. They proved the following result.

Theorem 1.2 (See Theorem 1.1 from [9]). *Let H be any fixed graph of chromatic number $r \geq 3$. Then,*

$$\phi(n, H) = \text{ex}(n, H) + o(n^2).$$

Pikhurko and Sousa also made the following conjecture.

Conjecture 1.3 [9]. *For any graph H of chromatic number $r \geq 3$, there exists $n_0 = n_0(H)$ such that $\phi(n, H) = \text{ex}(n, H)$ for all $n \geq n_0$.*

A graph H is *edge-critical* if there exists an edge $e \in E(H)$ such that $\chi(H) > \chi(H - e)$, where $\chi(H)$ denotes the *chromatic number* of H . For $r \geq 4$ a *clique-extension of order r* is a connected graph that consists of a K_{r-1} plus another vertex, say v , adjacent to at most $r - 2$ vertices of K_{r-1} . Conjecture 1.3 has been verified by Sousa for some edge-critical graphs, namely, clique-extensions of order $r \geq 4$ ($n \geq r$) [11] and the cycles of length 5 ($n \geq 6$) and 7 ($n \geq 10$) [10, 12]. Later, Özkahya and Person [7] verified the conjecture for all edge-critical graphs with chromatic number $r \geq 3$. Their result is the following.

Theorem 1.4 (See Theorem 3 from [7]). *For any edge-critical graph H with chromatic number $r \geq 3$, there exists $n_0 = n_0(H)$ such that $\phi(n, H) = \text{ex}(n, H)$, for all $n \geq n_0$. Moreover, the only graph attaining $\text{ex}(n, H)$ is the Turán graph $T_{r-1}(n)$.*

Recently, as an extension of Özkahya and Person's work, Allen, Böttcher, and Person [1] improved the error term obtained by Pikhurko and Sousa in Theorem 1.2. In fact, they proved that the error term $o(n^2)$ can be replaced by $O(n^{2-\alpha})$

for some $\alpha > 0$. Furthermore, they also showed that this error term has the correct order of magnitude. Their result is indeed an extension of Theorem 1.4 since the error term $O(n^{2-\alpha})$ that they obtained vanishes for every edge-critical graph H .

Conjecture 1.3 has also been verified by Liu and Sousa [6] for the k -fan graph F_k , which is the graph on $2k + 1$ vertices consisting of k triangles intersecting in exactly one common vertex. Observe that $\chi(F_k) = 3$ and for $k \geq 2$ the graph F_k is not edge-critical. Thus, the result of Liu and Sousa is not a particular case of Theorem 1.4 by Özkahya and Person.

In this article, we consider the *gem graphs* gem_4 and gem_5 , defined as follows. For the graph gem_4 , we take the path P_4 with vertices a, b, c, d and edges ab, bc, cd and add a universal vertex u adjacent to a, b, c, d . Similarly for the graph gem_5 , we take the path P_5 with vertices a, b, c, d, e and edges ab, bc, cd, de and add a universal vertex u adjacent to a, b, c, d, e . See Figure 1 below. For convenience, we write $abcd + u$ and $abcde + u$ for these two graphs.




Figure 1. The graphs gem_4 and gem_5 .

In Section 2, we will determine the Turán functions $\text{ex}(n, \text{gem}_4)$ for $n \geq 6$, and $\text{ex}(n, \text{gem}_5)$ for $n \geq 8$. Then, in Section 3, we will prove Pikhurko and Sousa conjecture for these two gem graphs. That is, we will show that $\phi(n, \text{gem}_4) = \text{ex}(n, \text{gem}_4)$ for $n \geq 6$, and $\phi(n, \text{gem}_5) = \text{ex}(n, \text{gem}_5)$ for $n \geq 8$. Note that $\chi(\text{gem}_4) = \chi(\text{gem}_5) = 3$, and that gem_4 and gem_5 are not edge-critical graphs. Thus, our results are again not implied by Theorem 1.4.

Our notations throughout the paper are fairly standard. For a vertex v in a graph G , the *neighbourhood* of v , denoted by $N(v)$, is the set of vertices in G that are adjacent to v . The *degree* of v is $\deg(v) = |N(v)|$, and the *minimum degree* and *maximum degree* of G are $\delta(G)$ and $\Delta(G)$, respectively. For a set $U \subset V(G)$, let $\deg(v, U)$ denote the number of vertices in U that are adjacent to v , and let $G[U]$ denote the subgraph of G induced by U .

2. TURÁN FUNCTION FOR THE GEM GRAPHS

In this section, we will determine the Turán functions $\text{ex}(n, \text{gem}_4)$ for $n \geq 6$, and $\text{ex}(n, \text{gem}_5)$ for $n \geq 8$. Furthermore, we will determine the extremal graphs in

each case. That is, we will determine all gem_4 -free graphs on $n \geq 6$ vertices with $\text{ex}(n, \text{gem}_4)$ edges, and all gem_5 -free graphs on $n \geq 8$ vertices with $\text{ex}(n, \text{gem}_5)$ edges.

2.1. Turán function for gem_4

We will now determine the function $\text{ex}(n, \text{gem}_4)$. In order to state our result, we first define the family of graphs $\mathcal{F}_{n,4}$, which will consist of all the extremal graphs. Let $n \geq 6$ and $\mathcal{F}_{n,4}$ be the family of graphs on n vertices as follows. For $n \equiv 0 \pmod{4}$, let G_n^0 be the graph obtained by taking the Turán graph $T_2(n)$ and embedding a maximum matching into a class of $T_2(n)$. For $n \equiv 1 \pmod{4}$, let G_n^{11} and G_n^{12} be the graphs obtained by embedding a maximum matching into the smaller class and the larger class of $T_2(n)$, respectively. For $n \equiv 2 \pmod{4}$, let G_n^{21} and G_n^{22} be the graphs obtained by embedding a maximum matching into a class of $T_2(n)$, and into the larger class of the complete bipartite graph $K_{n/2-1, n/2+1}$, respectively. For $n \equiv 3 \pmod{4}$, let G_n^3 be the graph obtained by embedding a maximum matching into the larger class of $T_2(n)$. Let the vertex classes of G_n^0 be A_n^0 and B_n^0 , with similar notations for the other graphs. Let $\mathcal{F}_{n,4} = \{G_n^0\}$, $\mathcal{F}_{n,4} = \{G_n^{11}, G_n^{12}\}$, $\mathcal{F}_{n,4} = \{G_n^{21}, G_n^{22}\}$ and $\mathcal{F}_{n,4} = \{G_n^3\}$ for $n \equiv 0, 1, 2, 3 \pmod{4}$, respectively. Figure 2 below shows the graphs of $\mathcal{F}_{n,4}$. Note that in G_n^{12} , we have an unmatched vertex in the class B_n^{12} , and similarly for G_n^{21} with the class B_n^{21} .




Figure 2. The graphs of $\mathcal{F}_{n,4}$.

It is easy to see that every graph of $\mathcal{F}_{n,4}$ is gem_4 -free. Let $G \in \mathcal{F}_{n,4}$, and suppose that there exists a copy of gem_4 in G , say $abcd + u$. We may consider in turn whether u is in the independent class of G , or in the class containing the maximum matching. In each case, we can easily verify that no four neighbours of u form a path P_4 in G , which is a contradiction. Also, for any graph of $\mathcal{F}_{n,4}$, by adding an edge, we obtain a graph that contains a copy of gem_4 . Indeed, let $G \in \mathcal{F}_{n,4}$. Since $n \geq 6$, if an edge cu is added to the independent class of G , then we may find an edge ab and another vertex d in the other class. If an edge bu is added to the class of G containing the maximum matching, then we may assume that du is an edge in the matching, and choose vertices a, c in the other class. In both cases, we have $abcd + u$ is a copy of gem_4 .

We can easily check that for $n \geq 6$, all graphs of $\mathcal{F}_{n,4}$ have the same number of edges. Thus for $G \in \mathcal{F}_{n,4}$, we let e_n denote the number of edges in the graph G . Then, we can easily check that the number of edges of G is

$$(1) \quad e(G) = e_n = \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n}{4} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Moreover, for $n \geq 7$, $G \in \mathcal{F}_{n,4}$ and $G' \in \mathcal{F}_{n-1,4}$, we have

$$(2) \quad e(G) - e(G') = e_n - e_{n-1} = \left\lfloor \frac{n}{2} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

We have the following result for the Turán function $\text{ex}(n, \text{gem}_4)$.

Theorem 2.1. *For $n \geq 6$, we have*

$$\text{ex}(n, \text{gem}_4) = e_n = \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n}{4} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Moreover, the only gem_4 -free graphs with n vertices and $\text{ex}(n, \text{gem}_4)$ edges are the members of $\mathcal{F}_{n,4}$.

We will prove Theorem 2.1 by induction on n . We first prove the base case as follows.

Lemma 2.2. *$\text{ex}(6, \text{gem}_4) = e_6 = 10$ and the only gem_4 -free graphs with six vertices and 10 edges are G_6^{21} and G_6^{22} .*

Proof. It suffices to prove that, for any graph G with six vertices and $e_6 = 10$ edges, either G contains a copy of the graph gem_4 , or $G \in \mathcal{F}_{6,4} = \{G_6^{21}, G_6^{22}\}$. Then for any graph G' with six vertices and $e(G') \geq 11$, we can take a spanning subgraph $G \subset G'$ with $e(G) = e_6 = 10$, so that either G contains a copy of gem_4 , or $G \in \mathcal{F}_{6,4}$. In either case, G' contains a copy of gem_4 and we are done.

Let G be a graph with six vertices and $e_6 = 10$ edges. Note that G has either a vertex of degree 5, or two vertices of degree 4. Otherwise, we have $e(G) \leq \lfloor \frac{1}{2}(4 + 5 \cdot 3) \rfloor = 9 < 10 = e_6$, a contradiction.

Suppose first that G has a vertex u with $\deg(u) = 5$. By Theorem 1.1, we have $\text{ex}(5, P_4) = \binom{3}{2} + \binom{2}{2} = 4$. We have $e(G - u) = 10 - 5 = 5 > 4 = \text{ex}(5, P_4)$, and thus $G - u$ contains a copy of the path P_4 , which together with u , form a copy of gem_4 in G .

Now, suppose that G has two vertices of degree 4, say u and v . Let x_1, x_2, x_3, x_4 be the remaining four vertices, and assume that G does not contain a copy of gem_4 . Suppose first that $uv \in E(G)$. If u and v have three common neighbours, say x_1, x_2, x_3 , then we must have $x_i x_4 \in E(G)$ for $i = 1, 2, 3$, so that $G = G_6^{21}$. If u and v have two common neighbours, say x_1, x_2 , then let $u x_3, v x_4 \in E(G)$ and $u x_4, v x_3 \notin E(G)$. We see that only the edges $x_1 x_2, x_3 x_4$ can be added to avoid creating a copy of gem_4 , so that G can only have at most nine edges, a contradiction. Now, suppose that $uv \notin E(G)$. Then G contains all edges between $\{u, v\}$ and $\{x_1, x_2, x_3, x_4\}$. If G does not contain a copy of gem_4 , then the remaining two edges must be independent within $\{x_1, x_2, x_3, x_4\}$, so that $G = G_6^{22}$.

We conclude that either G contains a copy of gem_4 , or $G \in \mathcal{F}_{6,4}$, as required. ■

We are now able to prove Theorem 2.1.

Proof of Theorem 2.1. Let $n \geq 6$. The lower bound $\text{ex}(n, \text{gem}_4) \geq e_n$ follows instantly by considering any graph of $\mathcal{F}_{n,4}$. We prove the upper bound $\text{ex}(n, \text{gem}_4) \leq e_n$ by induction on n . Lemma 2.2 proves the result for $n = 6$. Now suppose that $n \geq 7$, and the theorem holds for $n - 1$. We will prove that if G is a graph on n vertices and $e(G) = e_n$, then either G contains a copy of gem_4 , or G is one of the graphs of $\mathcal{F}_{n,4}$. This clearly implies the upper bound $\text{ex}(n, \text{gem}_4) \leq e_n$, and thus the theorem for n . Indeed, if we have a graph G' with n vertices and $e(G') > e_n$, then by taking a spanning subgraph $G \subset G'$ with $e(G) = e_n$, we see that either G contains a copy of gem_4 , or $G \in \mathcal{F}_{n,4}$. In either case, G' contains a copy of gem_4 .

First, suppose that $\delta(G) \leq \lfloor \frac{n}{2} \rfloor$ and let $v \in V(G)$ be a vertex of minimum degree. Then by (2), we have

$$(3) \quad e(G - v) = e(G) - \deg(v) \geq e_n - \lfloor \frac{n}{2} \rfloor \geq e_{n-1}.$$

If $e(G - v) > e_{n-1}$, then by induction, $G - v$, and thus G , contains a copy of gem_4 . Next, $e(G - v) = e_{n-1}$ holds if and only if $\deg(v) = \lfloor \frac{n}{2} \rfloor$ and $e_n - e_{n-1} = \lfloor \frac{n}{2} \rfloor$. The latter condition holds for $n \not\equiv 3 \pmod{4}$. By induction, either $G - v$, and thus G , contains a copy of gem_4 and we are done, or $G - v \in \mathcal{F}_{n-1,4}$, and we must consider the following cases.

Case 1. $n \equiv 0 \pmod{4}$. We have $G - v = G_{n-1}^3$ with classes A_{n-1}^3 and B_{n-1}^3 , where $|A_{n-1}^3| = \frac{n}{2} - 1$ and $|B_{n-1}^3| = \frac{n}{2}$, and B_{n-1}^3 containing a perfect matching. Since $\deg(v) = \frac{n}{2}$, if $N(v) = B_{n-1}^3$, then $G = G_n^0$. Otherwise, if v has neighbours $c \in A_{n-1}^3$ and $u \in B_{n-1}^3$, then $abcv + u$ is a copy of gem_4 in G , where $a \in A_{n-1}^3 \setminus \{c\}$ and $b \in B_{n-1}^3$ is the vertex adjacent to u .

Case 2. $n \equiv 1 \pmod{4}$. We have $G - v = G_{n-1}^0$ with classes A_{n-1}^0 and B_{n-1}^0 , where $|A_{n-1}^0| = |B_{n-1}^0| = \frac{n-1}{2}$, with B_{n-1}^0 containing a perfect matching. Since $\deg(v) = \frac{n-1}{2}$, it follows that if $N(v) = B_{n-1}^0$ then $G = G_n^{11}$, and if $N(v) = A_{n-1}^0$ then $G = G_n^{12}$. Otherwise, v has a neighbour in both A_{n-1}^0 and B_{n-1}^0 , so that as in Case 1, G contains a copy of gem_4 .

Case 3. $n \equiv 2 \pmod{4}$. We have $G - v \in \{G_{n-1}^{11}, G_{n-1}^{12}\}$. Suppose first that $G - v = G_{n-1}^{11}$. Then the classes of $G - v$ are A_{n-1}^{11} and B_{n-1}^{11} , where $|A_{n-1}^{11}| = \frac{n}{2} - 1$ and $|B_{n-1}^{11}| = \frac{n}{2}$, with A_{n-1}^{11} containing a perfect matching. Since $\deg(v) = \frac{n}{2}$, it follows that if $N(v) = B_{n-1}^{11}$, then $G = G_n^{21}$. Otherwise, v has a neighbour in both A_{n-1}^{11} and B_{n-1}^{11} , and G contains a copy of gem_4 as in Case 1. Now suppose that $G - v = G_{n-1}^{12}$. Then the classes are A_{n-1}^{12} and B_{n-1}^{12} , where $|A_{n-1}^{12}| = \frac{n}{2} - 1$ and $|B_{n-1}^{12}| = \frac{n}{2}$, with B_{n-1}^{12} containing a maximum matching with one unmatched vertex, say w . Since $\deg(v) = \frac{n}{2}$, it follows that if $N(v) = B_{n-1}^{12}$ then again $G = G_n^{21}$, and if $N(v) = A_{n-1}^{12} \cup \{w\}$ then $G = G_n^{22}$. Otherwise, v has a neighbour in both A_{n-1}^{12} and $B_{n-1}^{12} \setminus \{w\}$, and again as in Case 1, G contains a copy of gem_4 .

Next, suppose that $\delta(G) \geq \lfloor \frac{n}{2} \rfloor + 1$. In view of (1), if n is even, then we have $e(G) \geq \frac{n}{2}(\frac{n}{2} + 1) > e_n$. If $n \equiv 1 \pmod{4}$, then $e(G) \geq \lceil \frac{n}{2} (\lfloor \frac{n}{2} \rfloor + 1) \rceil = \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n}{4} \rfloor + 1 > e_n$. We have a contradiction in these cases. Now let $n \equiv 3 \pmod{4}$. We have $e(G) \geq \lceil \frac{n}{2} (\lfloor \frac{n}{2} \rfloor + 1) \rceil = \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n}{4} \rfloor + 1 = e_n$. We must have equality, and thus G is a $(\lfloor \frac{n}{2} \rfloor + 1)$ -regular graph. Let $v \in V(G)$, so that by (2)

$$(4) \quad e(G - v) = e(G) - \deg(v) = e_n - \left(\lfloor \frac{n}{2} \rfloor + 1 \right) = e_{n-1}.$$

By induction, either $G - v$, and thus G , contains a copy of gem_4 , or $G - v \in \mathcal{F}_{n-1,4}$. If the latter holds, then $G - v \in \{G_{n-1}^{21}, G_{n-1}^{22}\}$. Suppose first that $G - v = G_{n-1}^{21}$. The classes are A_{n-1}^{21} and B_{n-1}^{21} , where $|A_{n-1}^{21}| = |B_{n-1}^{21}| = \frac{n-1}{2}$, with B_{n-1}^{21} containing a maximum matching with one unmatched vertex, say w . Since $\deg(v) = \frac{n-1}{2} + 1$, in order for G to be $(\lfloor \frac{n}{2} \rfloor + 1)$ -regular, we must have $N(v) = A_{n-1}^{21} \cup \{w\}$. This gives $G = G_n^3$. Now, suppose that $G - v = G_{n-1}^{22}$. The classes are A_{n-1}^{22} and B_{n-1}^{22} , where $|A_{n-1}^{22}| = \frac{n-1}{2} - 1$ and $|B_{n-1}^{22}| = \frac{n-1}{2} + 1$, with B_{n-1}^{22} containing a perfect matching. Again since G is $(\lfloor \frac{n}{2} \rfloor + 1)$ -regular, we must have $N(v) = B_{n-1}^{22}$, and this also implies $G = G_n^3$.

This completes the proof of Theorem 2.1. ■

2.2. Turán function for gem_5

We will next determine the function $\text{ex}(n, \text{gem}_5)$. Analogously, we first define the family of graphs $\mathcal{F}_{n,5}$, which will consist of all the extremal graphs. Let $n \geq 8$ and $\mathcal{F}_{n,5}$ be the family of graphs on n vertices as follows. For $n \geq 11$, we let $\mathcal{F}_{n,5} = \mathcal{F}_{n,4}$. For $n = 8, 9, 10$, the family $\mathcal{F}_{n,5}$ will consist of all graphs of $\mathcal{F}_{n,4}$ and some additional graphs. Let G'_n be the graph obtained by adding one edge into each class of $T_2(n)$. Also for $n = 8$, let G''_8 be the graph obtained by embedding two vertex-disjoint triangles into the larger class of the complete bipartite graph $K_{2,6}$. For $n = 9$, let G''_9 be the graph obtained by taking G'_8 and joining another vertex to the four unmatched vertices within the classes of G'_8 . As before, let A'_8 and B'_8 be the classes of G'_8 , with similar notations for the other graphs. Figure 3 below shows these additional graphs. Let $\mathcal{F}_{8,5} = \{G'_8, G''_8, G'_9\}$, $\mathcal{F}_{9,5} = \{G''_9, G'_9, G''_9, G'_9\}$, and $\mathcal{F}_{10,5} = \{G''_{10}, G'_{10}, G'_{10}\}$.




Figure 3. The additional graphs in $\mathcal{F}_{n,5}$ for $n = 8, 9, 10$.

Note that every graph of $\mathcal{F}_{n,5}$ is gem_5 -free. Indeed, let $G \in \mathcal{F}_{n,5}$. If $G \notin \{G'_8, G''_8, G'_9, G''_9, G'_{10}\}$, then G is gem_4 -free as before, so that G is gem_5 -free. Suppose that $G \in \{G'_8, G''_8, G'_9, G''_9, G'_{10}\}$ and G contains a copy of gem_5 , say $abcde + u$. It is easy to check that in each choice for G , whichever vertex of G is chosen for u , we have that u does not have five neighbours that form a path P_5 in G . This is a contradiction.

Also, by adding an edge to any graph of $\mathcal{F}_{n,5}$, we obtain a graph that contains a copy of gem_5 . To see this, let $G \in \mathcal{F}_{n,5}$. Suppose first that $G \notin \{G'_8, G''_8, G'_9, G''_9, G'_{10}\}$. Then similar to before, since $n \geq 8$, it follows that if an edge cu is added to the independent class of G , then we can find two independent edges ab, de in the other class. If an edge bu is added to the class of G containing the

maximum matching, then we may assume that du is an edge in the matching, and choose vertices a, c, e in the other class. In both cases, we have $abcde + u$ is a copy of gem_5 . Next, the case $G \in \{G'_8, G'_9, G'_{10}\}$ can be considered similarly, according to whether or not the added edge is incident with an edge within a class of G . Now, consider $G = G''_8$. If the edge bu is added into A''_8 , then let cde be a triangle and a be another vertex in B''_8 . If an edge is added into B''_8 , then there exists a path $abcde$ of order 5 in B''_8 , and we let $u \in A''_8$. In both cases, $abcde + u$ is a copy of gem_5 . Finally, consider $G = G''_9$. Since G''_9 contains G''_8 as a subgraph on $A''_9 \cup B''_9$, it follows that if an edge is added into A''_9 or B''_9 , then we have a copy of gem_5 . Thus, we may assume that the edge au is added to G''_9 , where a is an end-vertex of the edge in A''_9 , and u is the vertex outside of $A''_9 \cup B''_9$. Then if $c, e \in A''_9$ and $b, d \in B''_9$ are the neighbours of u in G''_9 , we have $abcde + u$ is a copy of gem_5 .

We can easily check that for $n \geq 8$, all graphs of $\mathcal{F}_{n,5}$ have the same number of edges, which is also the same as the number of edges in any graph of $\mathcal{F}_{n,4}$. Thus, we may also let e_n denote the number of edges in any graph of $\mathcal{F}_{n,5}$. Then, equations (1) and (2) remain true. That is, for $G \in \mathcal{F}_{n,5}$, we have

$$(5) \quad e(G) = e_n = \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n}{4} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}, \end{cases}$$

and for $n \geq 9$, $G \in \mathcal{F}_{n,5}$ and $G' \in \mathcal{F}_{n-1,5}$, we have

$$(6) \quad e(G) - e(G') = e_n - e_{n-1} = \left\lfloor \frac{n}{2} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

We have the following result for the Turán function $\text{ex}(n, \text{gem}_5)$.

Theorem 2.3. *For $n \geq 8$, we have*

$$\text{ex}(n, \text{gem}_5) = e_n = \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n}{4} \right\rfloor + \begin{cases} 0 & \text{if } n \equiv 0, 1, 2 \pmod{4}, \\ 1 & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Moreover, the only gem_5 -free graphs with n vertices and $\text{ex}(n, \text{gem}_5)$ edges are the members of $\mathcal{F}_{n,5}$.

As before, Theorem 2.3 will be proved by induction on n . We first prove the base case, which will involve a bit more of case analysis than in Lemma 2.2.

Lemma 2.4. *$\text{ex}(8, \text{gem}_5) = e_8 = 18$ and the only gem_5 -free graphs with eight vertices and 18 edges are G_8^0, G'_8 and G''_8 .*

To prove Lemma 2.4, the following lemma will be useful.

Lemma 2.5. *Let H be a graph with vertex set $A \cup B$, where $A = \{x, y\}$ and $B = \{z_1, z_2, z_3, z_4\}$. Suppose that $xy, xz_4 \in E(H)$, and H also contains all edges between $\{x, y\}$ and $\{z_1, z_2, z_3\}$. Suppose that $H[B]$ contains two edges f_1, f_2 , and either z_4 belongs to at least one of f_1, f_2 , or $yz_4 \in E(H)$. Then H contains a copy of gem_5 .*

Proof. First, if z_4 belongs to one of f_1, f_2 , then we may assume that either $f_1 = z_1z_2, f_2 = z_3z_4$ or $f_1 = z_1z_2, f_2 = z_2z_4$ or $f_1 = z_1z_4, f_2 = z_2z_4$. Then $z_1z_2yz_3z_4+x$ or $z_3yz_1z_2z_4+x$ or $z_3yz_1z_4z_2+x$ is a copy of gem_5 in H , respectively.

Next, if $yz_4 \in E(H)$ and z_4 does not belong to f_1 and f_2 , then we may assume that $f_1 = z_1z_2$ and $f_2 = z_2z_3$. Then $z_1z_2z_3yz_4+x$ is a copy of gem_5 in H . ■

Proof of Lemma 2.4. Let G be a graph with eight vertices and $e_8 = 18$ edges. As in Lemma 2.2, it suffices to prove that either G contains a copy of gem_5 , or $G \in \mathcal{F}_{8,5} = \{G_8^0, G_8', G_8''\}$. Let $\Delta = \Delta(G)$ be the maximum degree of G . Note that $5 \leq \Delta \leq 7$, otherwise if $\Delta \leq 4$, then $e(G) \leq \lfloor \frac{1}{2} \cdot 8 \cdot 4 \rfloor = 16 < 18 = e_8$, a contradiction. Let $d_1 \geq d_2 \geq \dots \geq d_8$ be the degree sequence of G . Let $u \in V(G)$ be a vertex of maximum degree, so that $\deg(u) = \Delta = d_1$. We consider three cases according to the value of Δ .

Case 1. $\Delta = 7$. By Theorem 1.1, we have $\text{ex}(7, P_5) = \binom{4}{2} + \binom{3}{2} = 9$. Thus $e(G - u) = 18 - 7 = 11 > 9 = \text{ex}(7, P_5)$, and there exists a copy of the path P_5 in $G - u$, which together with u , form a copy of gem_5 in G .

Case 2. $\Delta = 6$. Let $v \in V(G) \setminus \{u\}$ be a vertex with $\deg(v) = d_2$. Note that $\deg(v) = 6$ or $\deg(v) = 5$, otherwise $e(G) \leq \lfloor \frac{1}{2}(6 + 7 \cdot 4) \rfloor = 17 < 18 = e_8$, a contradiction.

Subcase 2.1. $\deg(v) = 6$. Suppose first that $uv \notin E(G)$. We have $e(G - \{u, v\}) = 18 - 2 \cdot 6 = 6$. If there exists $x \in V(G) \setminus \{u, v\}$ with at least three neighbours in $V(G) \setminus \{u, v, x\}$, say x_1, x_2, x_3 , then $x_1ux_2vx_3+x$ is a copy of gem_5 in G . Otherwise, since $e(G - \{u, v\}) = 6$, we see that every vertex of $V(G) \setminus \{u, v\}$ must have exactly two neighbours in $V(G) \setminus \{u, v\}$, and thus, the subgraph $G - \{u, v\}$ must be either C_6 or two vertex-disjoint copies of C_3 . If the former, then there is a copy of P_5 in $G - \{u, v\}$, which together with u , form a copy of gem_5 . If the latter, then $G = G_8''$.

Now, suppose that $uv \in E(G)$. Observe first that u and v have at least four common neighbours in $V(G) \setminus \{u, v\}$. If $G[N(u) \setminus \{v\}]$ contains two edges, then Lemma 2.5 implies that G contains a copy of gem_5 . Otherwise, we may assume that $G[N(u) \setminus \{v\}]$ contains at most one edge. If y is the vertex not adjacent to u in G , then y has at most five neighbours in $N(u) \setminus \{v\}$. Therefore, we have $e(G - \{u, v\}) \leq 1 + 5 = 6$. This is a contradiction, since we have $e(G - \{u, v\}) = 18 - 1 - 2 \cdot 5 = 7$.

Subcase 2.2. $\deg(v) = 5$. Let $w \in V(G) \setminus \{u, v\}$ be a vertex with $\deg(w) = d_3$. Note that $\deg(w) = 5$, otherwise, $e(G) \leq \lfloor \frac{1}{2}(6 + 5 + 6 \cdot 4) \rfloor = 17 < 18 = e_8$. Thus, without loss of generality, we may assume $uw \in E(G)$, so that $e(G - \{u, v\}) = 18 - 1 - 5 - 4 = 8$. Let y be the vertex not adjacent to u . Suppose that G does not contain a copy of gem_5 .

Let $vy \notin E(G)$. Then v has exactly four neighbours in $N(u) \setminus \{v\}$, and by Lemma 2.5, $G[N(u) \setminus \{v\}]$ contains at most one edge, so that $e(G - \{u, v\}) \leq 6$, a contradiction.

Now let $vy \in E(G)$. Let x_1, x_2, x_3 be the common neighbours of u and v , and z_1, z_2 be the remaining two vertices, so that $uz_1, uz_2 \in E(G)$ and $vz_1, vz_2 \notin E(G)$. Again by Lemma 2.5, each of y, z_1, z_2 has at most one neighbour in $\{x_1, x_2, x_3\}$. If there are no edges between $\{y, z_1, z_2\}$ and $\{x_1, x_2, x_3\}$, then $e(G - \{u, v\}) \leq 6$, a contradiction. Otherwise, if there exists an edge between $\{y, z_1, z_2\}$ and $\{x_1, x_2, x_3\}$, then by Lemma 2.5, there are no edges in $G[\{x_1, x_2, x_3\}]$. Since there are at most three edges in $G[\{y, z_1, z_2\}]$ and at most three edges between $\{y, z_1, z_2\}$ and $\{x_1, x_2, x_3\}$, we have $e(G - \{u, v\}) \leq 6$, another contradiction.

Case 3. $\Delta = 5$. We have $d_1 = d_2 = d_3 = d_4 = \Delta = 5$, otherwise, $e(G) \leq \lfloor \frac{1}{2}(3 \cdot 5 + 5 \cdot 4) \rfloor = 17 < 18 = e_8$. This means that, we may assume there exists $v \in V(G) \setminus \{u\}$ with $\deg(v) = 5$ and $uv \in E(G)$, so that $e(G - \{u, v\}) = 18 - 1 - 2 \cdot 4 = 9$. If G contains a copy of gem_5 , then we are done, so assume otherwise.

Suppose first that u and v have four common neighbours, say x_1, x_2, x_3, x_4 . Let y_1, y_2 be the remaining two vertices. By Lemma 2.5, $G[\{x_1, x_2, x_3, x_4\}]$ contains at most one edge. If there is exactly one edge, say $x_1x_2 \in E(G)$, then there are 10 edges already in G . The edges between $\{y_1, y_2\}$ and $\{x_1, x_2, x_3, x_4\}$, as well as y_1y_2 , may possibly be present, and since $e(G) = 18$, exactly one of these nine edges is not present. Suppose first that $y_1y_2 \in E(G)$. We may assume that $y_1x_1, y_1x_2, y_2x_1 \in E(G)$, but then $uvx_2y_1y_2 + x_1$ is a copy of gem_5 . Otherwise, if $y_1y_2 \notin E(G)$, then we have $G = G'_8$. Finally, if there does not exist an edge in $G[\{x_1, x_2, x_3, x_4\}]$, then a similar edge count shows that G contains all edges between $\{y_1, y_2\}$ and $\{x_1, x_2, x_3, x_4\}$, as well as y_1y_2 . This gives $G = G_8^0$.

Next, suppose that u and v have three common neighbours, say x_1, x_2, x_3 . Let y, z_1, z_2 be the remaining vertices, where $uz_1, vz_2 \in E(G)$ and $uy, vy, uz_2, vz_1 \notin E(G)$. By Lemma 2.5, each of z_1, z_2 has at most one neighbour in $\{x_1, x_2, x_3\}$. If there exists an edge between $\{z_1, z_2\}$ and $\{x_1, x_2, x_3\}$, then again by Lemma 2.5, there are no edges in $G[\{x_1, x_2, x_3\}]$. Since there are at most three edges in $G[\{y, z_1, z_2\}]$, and at most five edges between $\{y, z_1, z_2\}$ and $\{x_1, x_2, x_3\}$, we have $e(G - \{u, v\}) \leq 8$, a contradiction. Otherwise, suppose that there are no edges between $\{z_1, z_2\}$ and $\{x_1, x_2, x_3\}$. Then we have $\deg(z_i) \leq 3$ for $i = 1, 2$. This implies that the remaining six vertices must each have degree 5, otherwise $e(G) \leq \lfloor \frac{1}{2}(5 \cdot 5 + 4 + 2 \cdot 3) \rfloor = 17 < 18 = e_8$. In particular, we have $x_ix_j \in E(G)$

for $1 \leq i \neq j \leq 3$ and $yx_i \in E(G)$ for $i = 1, 2, 3$. But then $uvx_2x_3y + x_1$ is a copy of gem_5 .

Finally, suppose that u and v have two common neighbours, say x_1, x_2 . Let y_1, y_2, z_1, z_2 be the remaining vertices, where $uy_1, uy_2, vz_1, vz_2 \in E(G)$ and $uz_1, uz_2, vy_1, vy_2 \notin E(G)$. Suppose first that there are at most two edges in $G[\{x_1, x_2, y_1, y_2\}]$, and at most two edges in $G[\{x_1, x_2, z_1, z_2\}]$. Since there are at most four edges between $\{y_1, y_2\}$ and $\{z_1, z_2\}$, we have $e(G - \{u, v\}) \leq 2 \cdot 2 + 4 = 8$, a contradiction. Now, suppose that there are at least three edges in $G[\{x_1, x_2, y_1, y_2\}]$. If $x_1y_1, y_1y_2 \in E(G)$ or $x_1y_1, x_2y_2 \in E(G)$, then $x_2vx_1y_1y_2 + u$ or $y_1x_1vx_2y_2 + u$ is a copy of gem_5 . Thus, we may assume that $x_1x_2, x_1y_1, x_2y_1 \in E(G)$ and $x_1y_2, x_2y_2, y_1y_2 \notin E(G)$. If there are at most two edges in $G[\{x_1, x_2, z_1, z_2\}]$, including x_1x_2 , then since there are at most four edges between $\{y_1, y_2\}$ and $\{z_1, z_2\}$, we have $e(G - \{u, v\}) \leq 3 + 1 + 4 = 8$, a contradiction. Thus, there are at least three edges in $G[\{x_1, x_2, z_1, z_2\}]$, and by similarly considering the edges in $G[\{x_1, x_2, z_1, z_2\}]$, we may assume that $x_1z_1, x_2z_1 \in E(G)$ and $x_1z_2, x_2z_2, z_1z_2 \notin E(G)$. But now, $y_1ux_2vz_1 + x_1$ is a copy of gem_5 .

Therefore, we conclude that either G contains a copy of gem_5 , or $G \in \mathcal{F}_{8,5}$. This completes the proof of Lemma 2.4. ■

We are now able to prove Theorem 2.3. The proof is generally similar to that of Theorem 2.1 but with a little more case analysis.

Proof of Theorem 2.3. Let $n \geq 8$. Again, the lower bound $\text{ex}(n, \text{gem}_5) \geq e_n$ follows by considering any graph of $\mathcal{F}_{n,5}$. We prove the upper bound $\text{ex}(n, \text{gem}_5) \leq e_n$ by induction on n . Lemma 2.4 proves the result for $n = 8$. Now suppose that $n \geq 9$, and the theorem holds for $n - 1$. As before, it suffices to prove that if G is a graph on n vertices and $e(G) = e_n$, then either G contains a copy of gem_5 , or $G \in \mathcal{F}_{n,5}$.

First, suppose that $\delta(G) \leq \lfloor \frac{n}{2} \rfloor$ and let $v \in V(G)$ be a vertex of minimum degree. Then exactly as in (3), we have $e(G - v) \geq e_{n-1}$. Again we are done unless $e(G - v) = e_{n-1}$, whence $\text{deg}(v) = \lfloor \frac{n}{2} \rfloor$ and $e_n - e_{n-1} = \lfloor \frac{n}{2} \rfloor$, and $n \not\equiv 3 \pmod{4}$. By induction, either $G - v$, and thus G , contains a copy of gem_5 and we are done, or $G - v \in \mathcal{F}_{n-1,5}$, and we must consider the following cases.

Case 1. $n \equiv 0 \pmod{4}$. We have $G - v = G_{n-1}^3$ with classes A_{n-1}^3 and B_{n-1}^3 , where $|A_{n-1}^3| = \frac{n}{2} - 1$ and $|B_{n-1}^3| = \frac{n}{2}$, and B_{n-1}^3 containing a perfect matching. We have $\text{deg}(v) = \frac{n}{2}$. If $N(v) = B_{n-1}^3$, then $G = G_n^0$. Otherwise, if v has neighbours $c, d \in A_{n-1}^3$ and $u \in B_{n-1}^3$, then $abcvd + u$ is a copy of gem_5 in G , where $a \in A_{n-1}^3 \setminus \{c, d\}$ and $b \in B_{n-1}^3$ is the vertex adjacent to u . If v has exactly one neighbour $u \in A_{n-1}^3$, then since $|B_{n-1}^3| = \frac{n}{2} > 4$, we can find $a, b, c, d \in B_{n-1}^3$ such that $ab, cd, bv, cv \in E(G)$. We have $abcvd + u$ is a copy of gem_5 in G .

Case 2. $n \equiv 1 \pmod{4}$. If $n \geq 13$, we have $G - v = G_{n-1}^0$. If $n = 9$, we have $G - v \in \{G_8^0, G_8', G_8''\}$.

Subcase 2.1. $n \geq 9$ and $G - v = G_{n-1}^0$. The classes of $G - v$ are A_{n-1}^0 and B_{n-1}^0 . Since $|B_{n-1}^0| = \frac{n-1}{2} \geq 4$, this subcase can be considered by combining the arguments used in Case 2 of Theorem 2.1 and in Case 1 above. We find that either G contains a copy of gem_5 , or $G \in \{G_n^{11}, G_n^{12}\}$.

Subcase 2.2. $n = 9$ and $G - v \in \{G_8', G_8''\}$. Suppose first that $G - v = G_8'$, so that the classes of $G - v$ are A_8' and B_8' with $|A_8'| = |B_8'| = 4$, and each class containing one edge, say cu and ab are the edges in A_8' and B_8' . We have $\deg(v) = 4$. If $N(v) = A_8'$ or $N(v) = B_8'$, then $G = G_9'$, and if $N(v) = (A_8' \cup B_8') \setminus \{a, b, c, u\}$, then $G = G_9''$. Otherwise, let $d \in B_8' \setminus \{a, b\}$. We may assume that $uv \in E(G)$, and either $av \in E(G)$ or $dv \in E(G)$. Then $vabcd + u$ or $abcdv + u$ is a copy of gem_5 .

Now, suppose that $G - v = G_8''$. The classes of $G - v$ are A_8'' and B_8'' with $|A_8''| = 2$, $|B_8''| = 6$, and there are two vertex-disjoint triangles embedded into B_8'' . Let $A_8'' = \{b, d\}$ and acu be one of the triangles in B_8'' . We have $\deg(v) = 4$. If $bv, dv \in E(G)$, then we may assume that $uv \in E(G)$. We have $abcdv + u$ is a copy of gem_5 . Otherwise, v has at least three neighbours in B_8'' , and we may assume that $av, uv \in E(G)$. Then $vabcd + u$ is a copy of gem_5 .

Case 3. $n \equiv 2 \pmod{4}$. If $n \geq 14$, then we have $G - v \in \{G_{n-1}^{11}, G_{n-1}^{12}\}$. If $n = 10$, then we have $G - v \in \{G_9^{11}, G_9^{12}, G_9', G_9''\}$.

Subcase 3.1. $n \geq 10$ and $G - v \in \{G_{n-1}^{11}, G_{n-1}^{12}\}$. If $G - v = G_{n-1}^{11}$, then $|A_{n-1}^{11}| = \frac{n}{2} - 1 \geq 4$. If $G - v = G_{n-1}^{12}$, then $G - v$ has the class B_{n-1}^{12} which contains a maximum matching with an unmatched vertex, say w . We have $|B_{n-1}^{12} \setminus \{w\}| = \frac{n}{2} - 1 \geq 4$. Since $\deg(v) = \frac{n}{2}$, this subcase can be considered by combining the arguments used in Case 3 of Theorem 2.1 and in Case 1 above. We find that either G contains a copy of gem_5 , or $G \in \{G_n^{21}, G_n^{22}\}$.

Subcase 3.2. $n = 10$ and $G - v \in \{G_9', G_9''\}$. Suppose first that $G - v = G_9'$, so that the classes of $G - v$ are A_9' and B_9' with $|A_9'| = 4$, $|B_9'| = 5$, and each class containing one edge. We have $\deg(v) = 5$. If $N(v) = B_9'$, then $G = G_{10}'$. If v has a neighbour which is incident with the edge in A_9' or the edge in B_9' , then as in the argument in the first part of Subcase 2.2, G contains a copy of gem_5 . Otherwise, $N(v)$ consists of the five vertices not incident with the two edges within A_9' and B_9' . Therefore, if $b, d \in A_9'$ and $a, c, e \in B_9'$ are these five neighbours of v , then $abcde + v$ is a copy of gem_5 .

Now, suppose that $G - v = G_9''$. The graph $G - v$ consists of two sets A_9'' and B_9'' where $|A_9''| = |B_9''| = 4$, with one edge in each set, say f_1 in A_9'' and f_2 in B_9'' , and another vertex, say z , joined to the four vertices not incident with f_1, f_2 . Let $b, d \in A_9''$ and $a, c \in B_9''$ be the neighbours of z in $G - v$. We have $\deg(v) = 5$.

Again, if v has a neighbour in each of A''_9 and B''_9 where at least one is incident with f_1 or f_2 , then by the argument in Subcase 2.2, G contains a copy of gem_5 . Otherwise, we may assume that $N(v) = A''_9 \cup \{z\}$ or $N(v) = \{a, b, c, d, z\}$, and $abcdv + z$ is a copy of gem_5 .

This concludes the case when $\delta(G) \leq \lfloor \frac{n}{2} \rfloor$.

Next, suppose that $\delta(G) \geq \lfloor \frac{n}{2} \rfloor + 1$. Then exactly as in the proof of Theorem 2.1, we must have $n \equiv 3 \pmod{4}$, and that G is a $(\lfloor \frac{n}{2} \rfloor + 1)$ -regular graph. Again for $v \in V(G)$, we have $e(G-v) = e_{n-1}$, using exactly the same argument as in (4). By induction, either $G-v$, and thus G , contains a copy of gem_5 , or $G-v \in \mathcal{F}_{n-1,5}$. If the latter holds, then for $n \geq 15$ we have $G-v \in \{G_{n-1}^{21}, G_{n-1}^{22}\}$, and for $n = 11$ we have $G-v \in \{G_{10}^{21}, G_{10}^{22}, G'_{10}\}$. If $n \geq 11$ and $G-v \in \{G_{n-1}^{21}, G_{n-1}^{22}\}$, then as in the proof of Theorem 2.1, the fact that G is a $(\lfloor \frac{n}{2} \rfloor + 1)$ -regular graph implies that $G = G_n^3$. Otherwise, we have $n = 11$ and $G-v = G'_{10}$. Then G is a 6-regular graph, which means that $N(v)$ consists of the six vertices not incident with the two edges within A'_{10} and B'_{10} . Therefore, if $a, c, e \in A'_{10}$ and $b, d \in B'_{10}$ are neighbours of v , then $abcde + v$ is a copy of gem_5 .

This completes the proof Theorem 2.3. ■

3. DECOMPOSITIONS OF GRAPHS INTO GEM GRAPHS AND SINGLE EDGES

Recall that for a fixed graph H , $\phi(n, H)$ denotes the smallest integer ϕ such that any graph on n vertices admits an H -decomposition with at most ϕ parts. In this section we will verify Pikhurko and Sousa conjecture (Conjecture 1.3) for the gem graphs gem_4 and gem_5 . That is, we will show that $\phi(n, \text{gem}_4) = \text{ex}(n, \text{gem}_4)$ for $n \geq 6$, and $\phi(n, \text{gem}_5) = \text{ex}(n, \text{gem}_5)$ for $n \geq 8$.

3.1. gem_4 -decompositions

We begin by considering gem_4 -decompositions, and prove the following result.

Theorem 3.1. *For $n \geq 6$ we have*

$$\phi(n, \text{gem}_4) = \text{ex}(n, \text{gem}_4).$$

Moreover, the only graphs attaining $\text{ex}(n, \text{gem}_4)$ are the members of $\mathcal{F}_{n,4}$.

Proof. Let $n \geq 6$. The lower bound $\phi(n, \text{gem}_4) \geq \text{ex}(n, \text{gem}_4)$ holds by considering any graph of $\mathcal{F}_{n,4}$. We prove the matching upper bound. By Theorem 2.1, we know that $\text{ex}(n, \text{gem}_4) = e_n$ for $n \geq 6$. Let G be a graph on $n \geq 6$ vertices. We must prove that $\phi(G, \text{gem}_4) \leq \text{ex}(n, \text{gem}_4) = e_n$, with equality if and only if $G \in \mathcal{F}_{n,4}$.

We proceed by induction on n . For $n = 6$, if $e(G) < e_6 = 10$, then we can simply decompose G into single edges to obtain $\phi(G, \text{gem}_4) < e_6$. Otherwise, let

$10 = e_6 \leq e(G) \leq 15$. By Theorem 2.1, we either have $G \in \mathcal{F}_{6,4}$, or G contains a copy of gem_4 . If $G \in \mathcal{F}_{6,4}$, then $e(G) = e_6 = 10$ and we must decompose G into single edges, thus, $\phi(G, \text{gem}_4) = e_6$ as required. If G contains a copy of gem_4 , then $\phi(G, \text{gem}_4) \leq 1 + e(G) - e(\text{gem}_4) \leq 9 < 10 = e_6$. Thus, the theorem holds for $n = 6$.

Now, let $n \geq 7$, and suppose that the theorem holds for $n - 1$. Let G be a graph on n vertices. As before, if $e(G) < e_n$, then $\phi(G, \text{gem}_4) < e_n$, simply by decomposing G into single edges. If $e(G) = e_n$, then by Theorem 2.1, either G contains a copy of gem_4 , in which case $\phi(G, \text{gem}_4) \leq 1 + e(G) - e(\text{gem}_4) = e_n - 6 < e_n$, or $G \in \mathcal{F}_{n,4}$, in which case we can only decompose G into e_n single edges for a gem_4 -decomposition, and $\phi(G, \text{gem}_4) = e_n$ as required.

Now, suppose that $e(G) > e_n$, and let $v \in V(G)$ be a vertex of minimum degree. If $\deg(v) \leq \lfloor \frac{n}{2} \rfloor$, then by equation (2) we have $e(G - v) = e(G) - \deg(v) > e_n - \lfloor \frac{n}{2} \rfloor \geq e_{n-1}$, that is, $G - v \notin \mathcal{F}_{n-1,4}$ and by the induction hypothesis we have

$$\phi(G - v, \text{gem}_4) < \text{ex}(n - 1, \text{gem}_4) = e_{n-1}.$$

Therefore, when going from $G - v$ to G we only need to use the edges joining v to the other vertices of G , and there are at most $\lfloor \frac{n}{2} \rfloor$ of these edges at v . We have

$$\phi(G, \text{gem}_4) \leq \phi(G - v, \text{gem}_4) + \deg(v) < e_{n-1} + \lfloor \frac{n}{2} \rfloor \leq e_n,$$

as required.

Therefore, we may assume that $\deg(v) \geq \lfloor \frac{n}{2} \rfloor + 1$ and let $\deg(v) = \lfloor \frac{n}{2} \rfloor + m$ for some integer $m \geq 1$. For every $x \in N(v)$, we have

$$\begin{aligned} \deg(x, N(v)) &\geq \lfloor \frac{n}{2} \rfloor + m - \left(n - \lfloor \frac{n}{2} \rfloor - m \right) \\ (7) \qquad \qquad &= 2 \lfloor \frac{n}{2} \rfloor + 2m - n \geq 2m - 1. \end{aligned}$$

This means that $G[N(v)]$ must contain a path P_{2m} on $2m$ vertices. Otherwise, if the longest path in $G[N(v)]$ has at most $2m - 1$ vertices, say with an end-vertex y , then all neighbours of y in $N(v)$ must lie in the path, so that $\deg(y, N(v)) \leq 2m - 2$, contradicting (7).

If $m \geq 2$, then the path P_{2m} contains $\lfloor \frac{2m}{4} \rfloor = \lfloor \frac{m}{2} \rfloor$ vertex-disjoint paths of order 4. Thus, we have $\lfloor \frac{m}{2} \rfloor$ edge-disjoint copies of gem_4 , where each copy is formed by a path of order 4, together with v . Let $F \subset G - v$ be the subgraph of order $n - 1$ obtained by deleting the edges of the paths of order 4 from $G - v$. By induction and (2), and since $m \geq 2$, we have

$$\begin{aligned} \phi(G, \text{gem}_4) &\leq \phi(F, \text{gem}_4) + \lfloor \frac{m}{2} \rfloor + \deg(v) - 4 \lfloor \frac{m}{2} \rfloor \\ &\leq e_{n-1} + \lfloor \frac{n}{2} \rfloor + m - 3 \lfloor \frac{m}{2} \rfloor < e_{n-1} + \lfloor \frac{n}{2} \rfloor \leq e_n. \end{aligned}$$

To complete the proof it remains to consider the case $m = 1$. For this case, we will repeatedly use the following claim.

Claim 3.2. *Suppose that there exists a vertex $z \in V(G)$ with $\deg(z) = \lfloor \frac{n}{2} \rfloor + 1$, and G has a copy of gem_4 with at least three edges incident to z . Then $\phi(G, \text{gem}_4) < e_n$.*

Proof. Let $F \subset G - z$ be the subgraph on $n - 1$ vertices obtained from $G - z$ by deleting the edges of the copy of gem_4 . By induction and (2), we have

$$\phi(G, \text{gem}_4) \leq \phi(F, \text{gem}_4) + 1 + \deg(z) - 3 \leq e_{n-1} + \lfloor \frac{n}{2} \rfloor - 1 < e_n. \quad \square$$

We now consider three cases. Let $\overline{N}(v) = V(G) \setminus (N(v) \cup \{v\})$, and note that

$$|N(v)| = \lfloor \frac{n}{2} \rfloor + 1 \geq 4 \quad \text{and} \quad |\overline{N}(v)| = \lfloor \frac{n}{2} \rfloor - 2 \geq 2.$$

Case 1. $G[N(v)]$ contains a path P of order 4. Then P and v form a copy of gem_4 , and we have $\phi(G, \text{gem}_4) < e_n$ by Claim 3.2.

Case 2. The order of the longest path in $G[N(v)]$ is 3. Let x_1x_2 be a path of order 3 in $G[N(v)]$.

Subcase 2.1. $x_1x_2 \in E(G)$. We have $\deg(x, N(v)) = 2$, for otherwise $G[N(v)]$ would contain a P_4 . We must have $\deg(x, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 3 \geq |\overline{N}(v)| - 1$. Similarly for x_1, x_2 . This implies that two of x, x_1, x_2 have a common neighbour in $\overline{N}(v)$, say $y \in \overline{N}(v)$ is a common neighbour of x, x_1 . Then $x_2vx_1y + x$ is a copy of gem_4 , and by Claim 3.2 with $z = v$, we have $\phi(G, \text{gem}_4) < e_n$.

Subcase 2.2. $x_1x_2 \notin E(G)$. Let $N(v) = \{x, x_1, x_2, \dots, x_{\lfloor n/2 \rfloor}\}$. For $i = 1, 2$, we have $\deg(x_i, N(v)) = 1$, and

$$(8) \quad \deg(x_i, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 2 \geq \lceil \frac{n}{2} \rceil - 2 = |\overline{N}(v)|.$$

We must have equality to hold throughout, whence n is odd, $\deg(x_1) = \deg(x_2) = \lfloor \frac{n}{2} \rfloor + 1$, and both x_1, x_2 are adjacent to all vertices of $\overline{N}(v)$. If x has a neighbour $y \in \overline{N}(v)$, then $x_1vx_2y + x$ is a copy of gem_4 , and again $\phi(G, \text{gem}) < e_n$ by Claim 3.2 with $z = v$.

Otherwise, suppose that x does not have a neighbour in $\overline{N}(v)$. Then $\deg(x) \leq |N(v) \cup \{v\}| - 1 = \lfloor \frac{n}{2} \rfloor + 1$, so that $\deg(x) = \lfloor \frac{n}{2} \rfloor + 1$ and $xx_i \in E(G)$ for all $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$. Moreover, we have $x_ix_j \notin E(G)$ for all $i \neq j$, otherwise there would exist a copy of P_4 in $G[N(v)]$. By a similar argument as in (8), we have $\deg(x_i) = \lfloor \frac{n}{2} \rfloor + 1$, and x_i is adjacent to all vertices of $\overline{N}(v)$ for all $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$. In order to get a contradiction, suppose that there does not exist a path of order 3

in $G[\overline{N}(v)]$. Then the maximum number of edges in $G[\overline{N}(v)]$ is $\lfloor \frac{1}{2}|\overline{N}(v)| \rfloor$. Recall that n is odd. We have

$$\begin{aligned} e(G) &\leq 2|N(v)| - 1 + (|N(v)| - 1)|\overline{N}(v)| + \left\lfloor \frac{1}{2}|\overline{N}(v)| \right\rfloor \\ &= 2\left\lfloor \frac{n}{2} \right\rfloor + 1 + \left\lfloor \frac{n}{2} \right\rfloor \left(\left\lceil \frac{n}{2} \right\rceil - 2 \right) + \left\lfloor \frac{1}{2} \left(\left\lceil \frac{n}{2} \right\rceil - 2 \right) \right\rfloor \\ &= \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n+1}{4} \right\rfloor = e_n, \end{aligned}$$

by (1), which contradicts the assumption $e(G) > e_n$. Therefore, $G[\overline{N}(v)]$ must have a path of order 3, say $y_1y_2y_3$. Note that $|\overline{N}(v)| = \lceil \frac{n}{2} \rceil - 2 \geq 3$ and thus we must have n odd and $n \geq 9$. Then, $x_1y_1x_2y_3 + y_2$ is a copy of gem_4 , and by Claim 3.2 with $z = x_1$, we have $\phi(G, \text{gem}) < e_n$.

Case 3. The longest path in $G[N(v)]$ has order 2. Note that this is indeed the remaining case, since $\deg(x, N(v)) \geq 2m - 1 = 1$ for all $x \in N(v)$ by (7). Moreover, $N(v)$ induces a perfect matching in G . Now by a similar argument as in (8), we must have n odd, and for every $x \in N(v)$, we have $\deg(x) = \lfloor \frac{n}{2} \rfloor + 1$ and x is adjacent to all vertices of $\overline{N}(v)$. Thus, we can find an edge x_1x_2 in $G[N(v)]$ and a common neighbour $y \in \overline{N}(v)$ of x_1, x_2 . Now, since vx_2y is a path of order 3 in $G[N(x_1)]$, we are done by applying Case 1 or Case 2 with x_1 in place of v .

The induction step is complete, and this completes the proof of Theorem 3.1. ■

3.2. gem_5 -decompositions

By using the same ideas as in the proof of Theorem 3.1, but with more case analysis, we will be able to prove a similar result for gem_5 -decompositions. That is, we will prove the following theorem.

Theorem 3.3. *For $n \geq 8$ we have*

$$\phi(n, \text{gem}_5) = \text{ex}(n, \text{gem}_5).$$

Moreover, the only graphs attaining $\text{ex}(n, \text{gem}_5)$ are the members of $\mathcal{F}_{n,5}$.

Proof. Let $n \geq 8$. As before, we have $\phi(n, \text{gem}_5) \geq \text{ex}(n, \text{gem}_5)$ by considering any graph of $\mathcal{F}_{n,5}$. By Theorem 2.3, to prove the matching upper bound, we must prove that if G is a graph on $n \geq 8$ vertices, then $\phi(G, \text{gem}_5) \leq \text{ex}(n, \text{gem}_5) = e_n$, with equality if and only if $G \in \mathcal{F}_{n,5}$.

We proceed by induction on n . For $n = 8$, if $e(G) < e_8 = 18$, then we can simply decompose G into single edges to obtain $\phi(G, \text{gem}_4) < e_8$. Next,

suppose that $18 = e_8 \leq e(G) \leq 25$. By Theorem 2.3, we either have $G \in \mathcal{F}_{8,5}$, or G contains a copy of gem_5 . If $G \in \mathcal{F}_{8,5}$, then $e(G) = e_8 = 18$ and we must decompose G into single edges, and $\phi(G, \text{gem}_5) = e_8$. If G contains a copy of gem_5 , then $\phi(G, \text{gem}_5) \leq 1 + e(G) - e(\text{gem}_5) \leq 17 < 18 = e_8$. Finally, suppose that $26 \leq e(G) \leq 28$. Clearly, there exist two vertices $x, y \in V(G)$ of degree 7, so that $e(G - \{x, y\}) \geq 26 - 1 - 2 \cdot 6 = 13$. Since $\text{ex}(6, P_5) = \binom{4}{2} + \binom{2}{2} = 7$ by Theorem 1.1, this means that we can find two edge-disjoint copies of P_5 in $G - \{x, y\}$. These two copies of P_5 , together with x and y , form two edge-disjoint copies of gem_5 in G . Thus, $\phi(G, \text{gem}_5) \leq 2 + e(G) - 2e(\text{gem}_5) \leq 12 < 18 = e_8$. The theorem holds for $n = 8$.

Now, let $n \geq 9$, and suppose that the theorem holds for $n - 1$. Let G be a graph on n vertices. As before, if $e(G) < e_n$, then $\phi(G, \text{gem}_5) < e_n$, simply by decomposing G into single edges. If $e(G) = e_n$, then by Theorem 2.3, either G contains a copy of gem_5 , in which case $\phi(G, \text{gem}_5) \leq 1 + e(G) - e(\text{gem}_5) = e_n - 8 < e_n$, or $G \in \mathcal{F}_{n,5}$, in which case we can only decompose G into e_n single edges for a gem_5 -decomposition, and $\phi(G, \text{gem}_5) = e_n$ as required.

Now, suppose that $e(G) > e_n$, and let $v \in V(G)$ be a vertex of minimum degree. If $\deg(v) \leq \lfloor \frac{n}{2} \rfloor$, then by equation (6), we have $e(G - v) = e(G) - \deg(v) > e_n - \lfloor \frac{n}{2} \rfloor \geq e_{n-1}$, that is, $G - v \notin \mathcal{F}_{n-1,5}$. By induction, we have $\phi(G - v, \text{gem}_5) < \text{ex}(n - 1, \text{gem}_5) = e_{n-1}$. Thus, when going from $G - v$ to G we only need to use the edges joining v to the other vertices of G . We have

$$\phi(G, \text{gem}_5) \leq \phi(G - v, \text{gem}_5) + \deg(v) < e_{n-1} + \left\lfloor \frac{n}{2} \right\rfloor \leq e_n.$$

Therefore, we may assume that $\deg(v) \geq \lfloor \frac{n}{2} \rfloor + 1$ and let $\deg(v) = \lfloor \frac{n}{2} \rfloor + m$ for some integer $m \geq 1$. As in (7), for every $x \in N(v)$, we have $\deg(x, N(v)) \geq 2m - 1$, and that $G[N(v)]$ must contain a path P_{2m} on $2m$ vertices.

If $m \geq 3$, then the path P_{2m} contains $\lfloor \frac{2m}{5} \rfloor$ vertex-disjoint paths of order 5. Thus, we have $\lfloor \frac{2m}{5} \rfloor$ edge-disjoint copies of gem_5 , where each copy is formed by a path of order 5, together with v . Let $F \subset G - v$ be the subgraph of order $n - 1$ obtained by deleting the edges of the paths of order 5 from $G - v$. By induction and (6), and since $m \geq 3$, we have

$$\begin{aligned} \phi(G, \text{gem}_5) &\leq \phi(F, \text{gem}_5) + \left\lfloor \frac{2m}{5} \right\rfloor + \deg(v) - 5 \left\lfloor \frac{2m}{5} \right\rfloor \\ &\leq e_{n-1} + \left\lfloor \frac{n}{2} \right\rfloor + m - 4 \left\lfloor \frac{2m}{5} \right\rfloor < e_{n-1} + \left\lfloor \frac{n}{2} \right\rfloor \leq e_n. \end{aligned}$$

For the rest of the proof, let $\overline{N}(v) = V(G) \setminus (N(v) \cup \{v\})$. Next, suppose that $m = 2$, so that $|N(v)| = \lfloor \frac{n}{2} \rfloor + 2 \geq 6$ and $|\overline{N}(v)| = \lceil \frac{n}{2} \rceil - 3 \geq 2$. If $G[N(v)]$ contains a path P_5 of order 5, then this path together with v form a copy of gem_5 .

Let $F \subset G - v$ be the subgraph of order $n - 1$, obtained by deleting the edges of the P_5 . Then,

$$\phi(G, \text{gem}_5) \leq \phi(F, \text{gem}_5) + 1 + \deg(v) - 5 \leq e_{n-1} + \lfloor \frac{n}{2} \rfloor + 2 - 4 < e_n.$$

Therefore, we may assume that the longest path in $G[N(v)]$ has order 4. Let $x_1x_2x_3x_4$ be such a path in $G[N(v)]$. Since $\deg(x_1, N(v)) \geq 2 \cdot 2 - 1 = 3$, we must have $x_1x_3, x_1x_4 \in E(G)$. Moreover, the only neighbours of x_1 in $N(v)$ are x_2, x_3, x_4 , so that

$$\deg(x_1, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 2 - 4 \geq \lceil \frac{n}{2} \rceil - 3 = |\overline{N}(v)|.$$

We must have equality, so that n is odd, $\deg(x_1) = \lfloor \frac{n}{2} \rfloor + 2$, and x_1 is adjacent to every vertex of $\overline{N}(v)$. The same argument holds for x_4 , so that x_1, x_4 have a common neighbour $y \in \overline{N}(v)$. Now, since $vx_2x_3x_4y$ is a path of order 5 in $G[N(x_1)]$, we are done by applying the previous argument with x_1 in place of v .

To complete the proof it remains to consider the case $m = 1$. As before, we will repeatedly use the following claim which is analogous to Claim 3.2.

Claim 3.4. *Suppose that there exists a vertex $z \in V(G)$ with $\deg(z) = \lfloor \frac{n}{2} \rfloor + 1$, and G has a copy of gem_5 with at least three edges incident to z . Then $\phi(G, \text{gem}_5) < e_n$.*

Proof. Exactly the same as the proof of Claim 3.2. □

We now consider four cases. Note that we have

$$|N(v)| = \lfloor \frac{n}{2} \rfloor + 1 \geq 5 \quad \text{and} \quad |\overline{N}(v)| = \lceil \frac{n}{2} \rceil - 2 \geq 3.$$

Case 1. $G[N(v)]$ contains a path P of order 5. Then P and v form a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4.

Case 2. The order of the longest path in $G[N(v)]$ is 4. Let $x_1x_2x_3x_4$ be such a path in $G[N(v)]$. It suffices to consider the following subcases.

Subcase 2.1. $x_1x_3, x_1x_4 \in E(G)$. For $i = 1, 2, 3, 4$, x_i does not have a neighbour in $N(v) \setminus \{x_1, x_2, x_3, x_4\}$, so that $\deg(x_i, N(v)) \leq 3$. Thus,

$$(9) \quad \deg(x_i, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 4 \geq \lceil \frac{n}{2} \rceil - 4 = |\overline{N}(v)| - 2.$$

If $x_2x_4 \notin E(G)$, then we have $\deg(x_j, N(v)) = 2$, and $\deg(x_j, \overline{N}(v)) \geq |\overline{N}(v)| - 1$ for $j = 2, 4$. With (9), this implies that either x_1, x_2 or x_2, x_3 or x_1, x_3 , have a common neighbour $y \in \overline{N}(v)$. Then, either $x_4vx_3x_2y + x_1$; or $x_4vx_1x_2y + x_3$; or $x_4vx_2x_3y + x_1$, is a copy of gem_5 , respectively. By Claim 3.4 with $z = v$, we

have $\phi(G, \text{gem}_5) < e_n$. Now, if $x_2x_4 \in E(G)$, then by (9), two of x_1, x_2, x_3, x_4 have a common neighbour in $\overline{N}(v)$. We may assume that x_1, x_2 have a common neighbour $y \in \overline{N}(v)$. Then we have $\phi(G, \text{gem}_5) < e_n$ by the same argument.

Subcase 2.2. $x_1x_3 \in E(G)$ and $x_1x_4, x_2x_4 \notin E(G)$. We see that x_3 is the only neighbour of x_4 in $N(v)$, so that

$$\deg(x_4, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 2 \geq \lceil \frac{n}{2} \rceil - 2 = |\overline{N}(v)|.$$

We must have equality throughout, so that $\deg(x_4) = \lfloor \frac{n}{2} \rfloor + 1$ and n is odd. Moreover, x_4 is adjacent to every vertex of $\overline{N}(v)$. If x_3 has a neighbour $y \in \overline{N}(v)$, then $x_1x_2vx_4y + x_3$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = v$. Now suppose that x_3 does not have a neighbour in $\overline{N}(v)$. Let $x_5, x_6, \dots, x_{\lfloor n/2 \rfloor + 1}$ be the remaining vertices of $N(v)$. Then $\deg(x_3) \geq \lfloor \frac{n}{2} \rfloor + 1$ implies that $x_3x_i \in E(G)$ for every $i \geq 5$. Moreover, we have $x_1x_i, x_2x_i \notin E(G)$ for all $i \geq 5$, otherwise we are in Subcase 2.1. This means that $\deg(x_i) = \lfloor \frac{n}{2} \rfloor + 1$ and x_i is adjacent to every vertex of $\overline{N}(v)$ for all $i \geq 4$. Also, note that for $i = 1, 2$,

$$\deg(x_i, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 3 = \lceil \frac{n}{2} \rceil - 3 = |\overline{N}(v)| - 1.$$

Suppose first that $G[\overline{N}(v)]$ contains a path of order 3, say $y_1y_2y_3$. If $n \geq 11$ so that $|N(v)| = \lfloor \frac{n}{2} \rfloor + 1 \geq 6$, then $x_4y_1x_5y_3x_6 + y_2$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = x_5$. Now let $n = 9$, and suppose that $x_1y_1, x_1y_2 \in E(G)$. Then $x_1y_1x_4y_3x_5 + y_2$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = x_4$. Thus, we may assume that $x_1y_1, x_1y_3, x_2y_1, x_2y_3 \in E(G)$ and $x_1y_2, x_2y_2 \notin E(G)$. It is easy to check that G is the graph G_9'' with $A_9'' = \{x_1, x_2, x_4, x_5\}$, $B_9'' = \{v, x_3, y_1, y_3\}$, and y_2 is the remaining vertex, so that $\phi(G, \text{gem}_5) = e_9 = \text{ex}(9, \text{gem}_5)$.

Now, suppose that $G[\overline{N}(v)]$ contains an edge, say y_1y_2 . If x_1 is adjacent to every vertex in $\overline{N}(v)$, then we may assume that $x_2y_1 \in E(G)$. Then $x_3vx_2y_1y_2 + x_1$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = v$. Thus we may assume that x_1 and x_2 are not adjacent to exactly one vertex in $\overline{N}(v)$. Since there are at most $|N(v)|$ edges in $G[N(v)]$ and at most $\lfloor \frac{1}{2}|\overline{N}(v)| \rfloor$ edges in $G[\overline{N}(v)]$, we have

$$\begin{aligned} e(G) &\leq 2|N(v)| + 2(|\overline{N}(v)| - 1) + (|N(v)| - 3)|\overline{N}(v)| + \left\lfloor \frac{1}{2}|\overline{N}(v)| \right\rfloor \\ &= 2n - 4 + \left(\lfloor \frac{n}{2} \rfloor - 2 \right) \left(\lceil \frac{n}{2} \rceil - 2 \right) + \left\lfloor \frac{1}{2} \left(\lceil \frac{n}{2} \rceil - 2 \right) \right\rfloor \\ &= \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n+1}{4} \right\rfloor = e_n, \end{aligned}$$

by (5) and since n is odd, which contradicts the assumption $e(G) > e_n$. Finally, if $G[\overline{N}(v)]$ does not contain an edge, then

$$\begin{aligned} e(G) &\leq 2|N(v)| + (|N(v)| - 1)|\overline{N}(v)| \\ &= 2\left(\left\lfloor \frac{n}{2} \right\rfloor + 1\right) + \left\lfloor \frac{n}{2} \right\rfloor \left(\left\lceil \frac{n}{2} \right\rceil - 2\right) = \left\lfloor \frac{n^2}{4} \right\rfloor + 2 \leq e_n, \end{aligned}$$

another contradiction.

Subcase 2.3. $x_1x_4 \in E(G)$ and $x_1x_3, x_2x_4 \notin E(G)$. For $i = 1, 2, 3, 4$, x_i does not have a neighbour in $N(v) \setminus \{x_1, x_2, x_3, x_4\}$, so that $\deg(x_i, N(v)) = 2$. Thus,

$$(10) \quad \deg(x_i, \overline{N}(v)) \geq \left\lfloor \frac{n}{2} \right\rfloor + 1 - 3 \geq \left\lceil \frac{n}{2} \right\rceil - 3 = |\overline{N}(v)| - 1.$$

If $\deg(x_1, \overline{N}(v)) = |\overline{N}(v)|$, then we can find $y_1, y_2 \in \overline{N}(v)$ such that y_1 is a common neighbour of x_1, x_2 , and y_2 is a common neighbour of x_2, x_3 . Then $y_1x_1vx_3y_2 + x_2$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = v$. Otherwise, we must have equality in (10) for $i = 1, 2, 3, 4$, so that n is odd, and for $i = 1, 2, 3, 4$, we have $\deg(x_i) = \left\lfloor \frac{n}{2} \right\rfloor + 1$, and x_i is not adjacent to exactly one vertex in $\overline{N}(v)$. If $n \geq 11$ so that $|\overline{N}(v)| = \left\lceil \frac{n}{2} \right\rceil - 2 \geq 4$, then we can again find the vertices $y_1, y_2 \in \overline{N}(v)$ and we are done as before. Now let $n = 9$, so that $|N(v)| = 5$, $|\overline{N}(v)| = 3$, and each x_i has exactly two neighbours in $\overline{N}(v)$. If x_1 and x_2 have two common neighbours in $\overline{N}(v)$, then we can again find $y_1, y_2 \in \overline{N}(v)$ as before and we are done. Otherwise, we may assume that $\overline{N}(v) = \{z_1, z_2, z_3\}$ with $x_1z_1, x_1z_2, x_2z_1, x_2z_3 \in E(G)$. If $z_1z_2 \in E(G)$, then $x_4vx_2z_1z_2 + x_1$ is a copy of gem_5 , and again $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = v$. A similar argument holds if $z_1z_3 \in E(G)$. Otherwise, we have at most one edge in $G[\overline{N}(v)]$, and since there are exactly nine edges in $G[N(v) \cup \{v\}]$ and at most $4 \cdot 2 + 3 = 11$ edges between $N(v)$ and $\overline{N}(v)$, we have $e(G) \leq 1 + 9 + 11 = 21 < 22 = e_9$, which is a contradiction.

Subcase 2.4. $x_1x_3, x_1x_4, x_2x_4 \notin E(G)$. We first note that x_2 is the only neighbour of x_1 in $N(v)$, so that

$$\deg(x_1, \overline{N}(v)) \geq \left\lfloor \frac{n}{2} \right\rfloor + 1 - 2 \geq \left\lceil \frac{n}{2} \right\rceil - 2 = |\overline{N}(v)|.$$

We must have equality throughout, so that n is odd, $\deg(x_1) = \left\lfloor \frac{n}{2} \right\rfloor + 1$, and x_1 is adjacent to all vertices of $\overline{N}(v)$. The exact same properties hold for x_4 . Next, suppose that x_2 has p neighbours in $N(v) \setminus \{x_1, x_2, x_3, x_4\}$, where $0 \leq p \leq \left\lfloor \frac{n}{2} \right\rfloor - 3$. Let S_2 be the set of these p neighbours. We have

$$(11) \quad \deg(x_2, \overline{N}(v)) \geq \left\lfloor \frac{n}{2} \right\rfloor + 1 - 3 - p = \left\lceil \frac{n}{2} \right\rceil - 3 - p.$$

Now, x_3 does not have a neighbour in S_2 , otherwise there would exist a path of order 5 in $G[N(v)]$. Thus, x_3 has at most $|N(v)| - 4 - p = \lfloor \frac{n}{2} \rfloor - 3 - p$ neighbours in $N(v) \setminus \{x_1, x_2, x_3, x_4\}$. Let S_3 be these neighbours of x_3 , so that $S_2 \cap S_3 = \emptyset$. We have

$$(12) \quad \deg(x_3, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 3 - \left(\lfloor \frac{n}{2} \rfloor - 3 - p \right) = p + 1.$$

Suppose that x_2, x_3 have a common neighbour $y_1 \in \overline{N}(v)$. Clearly, from (11) and (12), at least one of x_2, x_3 has at least two neighbours in $\overline{N}(v)$. If x_2 has this property, then x_1, x_2 have a common neighbour $y_2 \in \overline{N}(v) \setminus \{y_1\}$. Thus, $y_1 x_3 v x_1 y_2 + x_2$ is a copy of gem_5 , and by Claim 3.4 with $z = v$, we have $\phi(G, \text{gem}_5) < e_n$. A similar argument holds if x_3 has at least two neighbours in $\overline{N}(v)$, with x_4 in place of x_1 .

Thus, if $T_2, T_3 \subset \overline{N}(v)$ are the sets of neighbours of x_2, x_3 in $\overline{N}(v)$, respectively, then we may assume that $T_2 \cap T_3 = \emptyset$. Note that from (11) and (12), we have

$$\deg(x_2, \overline{N}(v)) + \deg(x_3, \overline{N}(v)) \geq \lceil \frac{n}{2} \rceil - 2 = |\overline{N}(v)|.$$

Thus, we must have equality above, as well as in (11) and (12). This means that $\deg(x_2) = \deg(x_3) = \lfloor \frac{n}{2} \rfloor + 1$, and we have the partitions $N(v) \setminus \{x_1, x_2, x_3, x_4\} = S_2 \dot{\cup} S_3$ and $\overline{N}(v) = T_2 \dot{\cup} T_3$. Clearly, there are no edges in $G[S_2 \cup S_3]$, otherwise there would exist a path of order 5 in $G[N(v)]$. Next, suppose that there is a path of order 3 in $G[\overline{N}(v)]$, say $y_1 y_2 y_3$. Suppose that $y_2 \in T_2$. Then $x_2 x_1 y_1 x_4 y_3 + y_2$ is a copy of gem_5 , so that by Claim 3.4 with $z = x_1$, we have $\phi(G, \text{gem}_5) < e_n$. A similar argument holds if $y_2 \in T_3$. Otherwise, we have $|N(v)| - 1$ edges in $G[N(v)]$, $|\overline{N}(v)|$ edges between $\{x_2, x_3\}$ and $\overline{N}(v)$, and at most $\lfloor \frac{1}{2} |\overline{N}(v)| \rfloor$ edges in $G[\overline{N}(v)]$. By (5) and since n is odd,

$$\begin{aligned} e(G) &\leq 2|N(v)| - 1 + |\overline{N}(v)| + (|N(v)| - 2)|\overline{N}(v)| + \left\lfloor \frac{1}{2} |\overline{N}(v)| \right\rfloor \\ &= 2 \lfloor \frac{n}{2} \rfloor + \lceil \frac{n}{2} \rceil - 1 + \left(\lfloor \frac{n}{2} \rfloor - 1 \right) \left(\lceil \frac{n}{2} \rceil - 2 \right) + \left\lfloor \frac{1}{2} \left(\lceil \frac{n}{2} \rceil - 2 \right) \right\rfloor \\ &= \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n+1}{4} \right\rfloor = e_n, \end{aligned}$$

which contradicts the assumption $e(G) > e_n$.

Case 3. The order of the longest path in $G[N(v)]$ is 3. Let $x_1 x x_2$ be such a path in $G[N(v)]$. We consider the following subcases.

Subcase 3.1. $x_1 x_2 \in E(G)$. We have $\deg(x, N(v)) = 2$, for otherwise $G[N(v)]$ would contain a P_4 . Thus

$$\deg(x, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 3 \geq \lceil \frac{n}{2} \rceil - 3 = |\overline{N}(v)| - 1.$$

Similar inequalities hold for x_1, x_2 . If $\deg(x, \overline{N}(v)) = |\overline{N}(v)|$, then there exist $y_1, y_2 \in \overline{N}(v)$ such that y_i is a common neighbour of x, x_i for $i = 1, 2$. Then $y_1x_1vx_2y_2 + x$ is a copy of gem_5 , and by Claim 3.4 with $z = v$, we have $\phi(G, \text{gem}_5) < e_n$. Otherwise, we have $\deg(x, \overline{N}(v)) = |\overline{N}(v)| - 1$, whence n is odd and $\deg(x) = \lfloor \frac{n}{2} \rfloor + 1$. We may assume that x, x_1 have a common neighbour $y \in \overline{N}(v)$. Now, vx_2x_1y is a path of order 4 in $G[N(x)]$, and we are done by applying Case 1 or Case 2 with x in place of v .

Subcase 3.2. $x_1x_2 \notin E(G)$. Let $N(v) = \{x, x_1, x_2, \dots, x_{\lfloor n/2 \rfloor}\}$. For $i = 1, 2$, we have

$$(13) \quad \deg(x_i, \overline{N}(v)) \geq \lfloor \frac{n}{2} \rfloor + 1 - 2 \geq \lceil \frac{n}{2} \rceil - 2 = |\overline{N}(v)|.$$

We must have equality to hold throughout, whence n is odd, $\deg(x_1) = \deg(x_2) = \lfloor \frac{n}{2} \rfloor + 1$, and both x_1, x_2 are adjacent to all vertices of $\overline{N}(v)$. If x has neighbours $y_1, y_2 \in \overline{N}(v)$, then we are done as in Subcase 3.1. If x has exactly one neighbour $y \in \overline{N}(v)$, then we have

$$\deg(x, N(v) \setminus \{x, x_1, x_2\}) \geq \lfloor \frac{n}{2} \rfloor + 1 - 4 \geq 1,$$

and we may assume that $xx_3 \in E(G)$. Then $x_1yx_2vx_3 + x$ is a copy of gem_5 , and we have $\phi(G, \text{gem}_5) < e_n$ by Claim 3.4 with $z = v$. Otherwise, suppose that x does not have a neighbour in $\overline{N}(v)$. We may apply the exact same argument as in Subcase 2.2 of Theorem 3.1 to deduce that x_i is adjacent to all vertices of $\overline{N}(v)$ for all $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$, and $G[\overline{N}(v)]$ must contain a path of order 3, say $y_1y_2y_3$. Then $x_1y_1x_2y_3x_3 + y_2$ is a copy of gem_5 , and by Claim 3.4 with $z = x_2$, we have $\phi(G, \text{gem}_5) < e_n$.

Case 4. The longest path in $G[N(v)]$ has order 2. Note that this is indeed the remaining case, since $\deg(x, N(v)) \geq 2m - 1 = 1$ for all $x \in N(v)$. Moreover, $N(v)$ induces a perfect matching in G . By a similar argument as in (13), we must have n odd, and for every $x \in N(v)$, we have $\deg(x) = \lfloor \frac{n}{2} \rfloor + 1$ and x is adjacent to all vertices of $\overline{N}(v)$. Thus, we can find an edge x_1x_2 in $G[N(v)]$ and a common neighbour $y \in \overline{N}(v)$ of x_1, x_2 . Now, since vx_2y is a path of order 3 in $G[N(x_1)]$, we are done by applying Case 1, Case 2 or Case 3 with x_1 in place of v .

The induction step is complete, and this completes the proof of Theorem 3.3. ■

Acknowledgements

Henry Liu was supported by the International Interchange Plan of CSU, and the China Postdoctoral Science Foundation (Nos. 2015M580695 and 2016T90756). Teresa Sousa was partially supported by FCT - Fundação para a Ciência e a Tecnologia (Portuguese Science Foundation, Portugal), through the project UID/MAT/00297/2013 (Centro de Matemática e Aplicações). The authors thank the anonymous referee for the careful reading of the manuscript.

REFERENCES

- [1] P. Allen, J. Böttcher and Y. Person, *An improved error term for minimum H -decompositions of graphs*, J. Combin. Theory Ser. B **108** (2014) 92–101.
doi:10.1016/j.jctb.2014.03.001
- [2] B. Bollobás, *On complete subgraphs of different orders*, Math. Proc. Cambridge Philos. Soc. **79** (1976) 19–24.
doi:10.1017/S0305004100052063
- [3] D. Dor and M. Tarsi, *Graph decomposition is NP-complete: a complete proof of Holyer’s conjecture*, SIAM J. Comput. **26** (1997) 1166–1187.
doi:10.1137/S0097539792229507
- [4] P. Erdős, A.W. Goodman and L. Pósa, *The representation of a graph by set intersections*, Canad. J. Math. **18** (1966) 106–112.
doi:10.4153/CJM-1966-014-3
- [5] R. Faudree and R. Schelp, *Path Ramsey numbers in multicolorings*, J. Combin. Theory Ser. B **19** (1975) 150–160.
doi:10.1016/0095-8956(75)90080-5
- [6] H. Liu and T. Sousa, *Decompositions of graphs into fans and single edges*, J. Graph Theory **85** (2017) 400–411.
doi:/10.1002/jgt.22069
- [7] L. Özkahya and Y. Person, *Minimum H -decompositions of graphs: edge-critical case*, J. Combin. Theory Ser. B **102** (2012) 715–725.
doi:10.1016/j.jctb.2011.10.004
- [8] O. Pikhurko, *A note on the Turán function of even cycles*, Proc. Amer. Math. Soc. **140** (2012) 3687–3692.
doi:10.1090/S0002-9939-2012-11274-2
- [9] O. Pikhurko and T. Sousa, *Minimum H -decompositions of graphs*, J. Combin. Theory Ser. B **97** (2007) 1041–1055.
doi:10.1016/j.jctb.2007.03.002
- [10] T. Sousa, *Decompositions of graphs into 5-cycles and other small graphs*, Electron. J. Combin. **12** (2005) #R49.
- [11] T. Sousa, *Decompositions of graphs into a given clique-extension*, Ars Combin. **100** (2011) 465–472.
- [12] T. Sousa, *Decompositions of graphs into cycles of length seven and single edges*, Ars Combin. **119** (2015) 321–329.
- [13] P. Turán, *On an extremal problem in graph theory*, Mat. Fiz. Lapok **48** (1941) 436–452.

Received 10 October 2016

Revised 26 January 2017

Accepted 2 February 2017