

*Discussiones Mathematicae*  
*Graph Theory* 38 (2018) 977–989  
doi:10.7151/dmgt.2058

## TOTAL COLORINGS OF EMBEDDED GRAPHS WITH NO 3-CYCLES ADJACENT TO 4-CYCLES

BING WANG

*Department of Mathematics*  
*Zaozhuang University, Shandong, 277160, China*

**e-mail:** hellobingzi@163.com.

JIAN-LIANG WU

*School of Mathematics*  
*Shandong University, Jinan, 250100, China*

AND

LIN SUN

*School of Mathematics*  
*Shandong University, Jinan, 250100, China*

*Department of Mathematics*  
*Changji University, Changji, 831100, China*

### Abstract

A *total- $k$ -coloring* of a graph  $G$  is a coloring of  $V \cup E$  using  $k$  colors such that no two adjacent or incident elements receive the same color. The *total chromatic number*  $\chi''(G)$  of  $G$  is the smallest integer  $k$  such that  $G$  has a total- $k$ -coloring. Let  $G$  be a graph embedded in a surface of Euler characteristic  $\varepsilon \geq 0$ . If  $G$  contains no 3-cycles adjacent to 4-cycles, that is, no 3-cycle has a common edge with a 4-cycle, then  $\chi''(G) \leq \max\{8, \Delta + 1\}$ .

**Keywords:** total coloring, embedded graph, cycle.

**2010 Mathematics Subject Classification:** 05C15.

### 1. INTRODUCTION

All graphs considered in this paper are simple, finite and undirected, and we follow [2] for the terminologies and notations not defined here. Let  $G$  be a graph.

We use  $V(G)$ ,  $E(G)$ ,  $\Delta(G)$  and  $\delta(G)$  (or simply  $V$ ,  $E$ ,  $\Delta$  and  $\delta$ ) to denote the vertex set, the edge set, the maximum degree and the minimum degree of  $G$ , respectively. A *total- $k$ -coloring* of a graph  $G$  is a coloring of  $V \cup E$  using  $k$  colors such that no two adjacent or incident elements receive the same color. The *total chromatic number*  $\chi''(G)$  of  $G$  is the smallest integer  $k$  such that  $G$  has a total- $k$ -coloring. Clearly,  $\chi''(G) \geq \Delta + 1$ . Behzad [1] and Vizing [18] posed independently the following famous conjecture, which is known as the Total Coloring Conjecture (TCC).

**Conjecture.** *For any graph  $G$ ,  $\chi''(G) \leq \Delta + 2$ .*

This conjecture was confirmed for all graphs with  $\Delta \leq 3$  independently by Vijayaditya and Rosenfeld in 1971, and in [13, 14], Kostochka proved that if  $4 \leq \Delta \leq 5$ , then  $\chi''(G) \leq \Delta + 2$ . Later, Kostochka [15] renewed the proof for  $\Delta = 5$ . We summary these result to the following lemma.

**Lemma 1.** *Let  $G$  be a graph with  $\Delta(G) \leq 5$ . Then  $\chi''(G) \leq 7$ .*

But for planar graphs, the famous conjecture was first proved by Borodin [4] for  $\Delta \geq 11$  and then for  $\Delta \geq 9$  [3], which was extended to  $\Delta \geq 8$  by Jensen and Toft [9] and to  $\Delta \geq 7$  by Sanders and Zhao [17]. So the only open case is  $\Delta = 6$ .

Interestingly, planar graphs with high maximum degree allow a stronger assertion, that is, every planar graph with high maximum degree  $\Delta$  has a total- $(\Delta + 1)$ -coloring. This result was first established in [4] for  $\Delta \geq 16$ , which was extended to  $\Delta \geq 14$  [3],  $\Delta \geq 12$  [5],  $\Delta \geq 11$  [6],  $\Delta \geq 10$  [25] and finally  $\Delta \geq 9$  [10]. However, for  $\Delta \in \{4, 5, 6, 7, 8\}$ , it is not known if the assertion still holds true. Such a study has attracted a considerable amount of attention. Recently, Shen *et al.* [11] proved that if  $G$  is a planar graph with  $\Delta = 8$  and  $G$  contains no chordal 5-cycles or no chordal 6-cycles, then  $\chi''(G) = \Delta + 1$ . Wang and Wu [19] proved that if  $G$  is a planar graph with  $\Delta \geq 7$  and every vertex is incident with at most one triangle, then  $\chi''(G) = \Delta + 1$ . Wang and Wu [20] proved that if  $G$  is a planar graph with  $\Delta \geq 7$  with no 4-cycles, then  $\chi''(G) = \Delta + 1$  (later, it is extended to  $\Delta \geq 6$  by Shen and Wang [12]). Chang *et al.* [7] proved that if  $G$  is a planar graph with  $\Delta \geq 7$  and every vertex  $v$  has an integer  $k_v \in \{3, 4, 5, 6\}$ , such that  $v$  is not in any  $k_v$ -cycle, then  $\chi''(G) = \Delta + 1$ .

Let  $G$  be a graph embedded in a surface of Euler characteristic  $\varepsilon$ , where *surfaces* in this paper are compact, connected 2-dimensional manifolds without boundary. All embeddings considered in this paper are *2-cell embeddings*. Wu and Wang [24] proved that if  $\varepsilon < 0$  and  $\Delta(G) \geq \sqrt{25 - 24\varepsilon} + 10$ , then  $\chi'_{list}(G) = \Delta(G)$  and  $\chi''_{list}(G) = \Delta(G) + 1$ , which extends a result of Borodin, Kostochka and Woodall in [5]. They also proved that  $\chi''(G) = \Delta(G) + 1$  if  $\varepsilon \geq 0$ ,  $\Delta(G) \geq 9$  and no two triangles have a common edge, or if  $\varepsilon \geq 0$ ,  $\Delta(G) \geq 8$  and no two triangles have a common vertex. Wang *et al.* [22] proved that if  $\varepsilon \geq 0$  and  $\Delta(G) \geq 7$ ,

then  $\chi''(G) \leq \Delta + 2$ . Wang *et al.* [23] proved that if  $\varepsilon \geq 0$  and  $\Delta \geq 9$ , then  $\chi''(G) = \Delta + 1$ . In this paper, we shall prove the following result.

**Theorem 2.** *Let  $G$  be a graph embedded in a surface of Euler characteristic  $\varepsilon \geq 0$ . If  $G$  contains no 3-cycles adjacent to 4-cycles, then  $\chi''(G) \leq \max\{8, \Delta(G)+1\}$ .*

The theorem shows that if a graph  $G$  can be embedded in a surface of Euler characteristic  $\varepsilon \geq 0$ , and contains no 3-cycles adjacent to 4-cycles, and  $\Delta \geq 7$ , then  $\chi''(G) = \Delta + 1$ .

## 2. PROOF OF THEOREM 2

We will introduce some more notations and definitions here for convenience. Let  $G = (V, E, F)$  be an embedded graph, where  $F$  is the face set of  $G$ . For a vertex  $v \in V$ , let  $N(v)$  denote the set of vertices adjacent to  $v$ , and let  $d(v) = |N(v)|$  denote the degree of  $v$ , and for a face  $f$ , the degree of a face  $f$ , denoted by  $d(f)$ , is the number of edges incident with it, where each cut-edge is counted twice. A  $k$ -vertex, a  $k^+$ -vertex or a  $k^-$ -vertex is a vertex of degree  $k$ , at least  $k$  or at most  $k$ , respectively. Similarly, A  $k$ -face, a  $k^+$ -face is a face of degree  $k$  or at least  $k$ , respectively. Let  $n_t(v)$  be the number of  $t$ -vertices adjacent to a vertex  $v$ , and  $f_k(v)$  the number of  $k$ -faces incident with  $v$ . Especially, let  $f_3(v) = t$ . Let  $v_1, v_2, \dots, v_d$  be neighbors of  $v$  in an anticlockwise order. Let  $f_i$  be face incident with  $v$ ,  $v_i$  and  $v_{i+1}$ , for all  $i$  such that  $i \in \{1, 2, \dots, d\}$ . Note that all the subscripts in the paper are taken modulo  $d$ . For convenience,  $(d_1, d_2, \dots, d_n)$  denotes a cycle (or a face) whose boundary vertices are of degree  $d_1, d_2, \dots, d_n$  in the anticlockwise order. Specially,  $(i, j^+, k^+)$ -face is a 3-face  $uvw$  such that  $d(u) = i \leq j \leq d(v) \leq k \leq d(w)$ .

**Proof of Theorem 2.** Let  $m = \max\{7, \Delta\}$  and  $G = (V, E, F)$  be a minimal counterexample to Theorem 2 with  $|V| + |E|$  as small as possible. Then every proper subgraph of  $G$  has a total- $(m + 1)$ -coloring, but  $G$  itself does not. First we show some known properties of  $G$ .

- (a) Every 3-cycle is not adjacent to a  $4^-$ -face. It follows that  $f_3(v) \leq \lfloor \frac{d(v)}{2} \rfloor$  for any  $v \in V(G)$ .
- (b) For any edge  $uv \in E(G)$ , if  $\min\{d(u), d(v)\} \leq \lfloor \frac{m}{2} \rfloor$ , then  $d(u) + d(v) \geq m + 2$ . So all neighbors of any 2-vertex are  $7^+$ -vertices and all neighbors of any 3-vertex are  $6^+$ -vertices (see [20]).
- (c) The subgraph  $G_2$  of  $G$  induced by all edges incident with 2-vertices is a forest. So for any component of  $G_2$ , we root it at a  $7^+$ -vertex. Then every 2-vertex has exactly one *parent* and exactly one *child* (see [3, 6]).

- (d) Each 3-face of  $G$  is not incident with two  $4^-$ -vertices (see [16]).
- (e) If  $v$  is a vertex of  $G$  with  $n_2(v) \geq 1$ , then  $n_{4^+}(v) \geq 1$  (see [7]).

**Lemma 3** [21]. *Suppose  $v$  is a  $d$ -vertex of  $G$  with  $d \geq 5$ . Let  $v_1, \dots, v_d$  be the neighbors of  $v$  and  $f_1, \dots, f_d$  be the faces incident with  $v$  in clockwise order, where  $f_i$  is incident with  $v_i$  and  $v_{i+1}$ ,  $i = 1, 2, \dots, d$ . Note that eventually  $v_1$  and  $v_{d+1}$  is the same vertex. Then there does not exist an integer  $i$  ( $2 \leq i \leq d$ ) such that  $d(v_1) = d(v_i) = 2$ ,  $d(v_k) = 3$  ( $2 \leq k \leq i - 1$ ) and  $d(f_t) = 4$  ( $1 \leq t \leq i - 1$ ).*

**Lemma 4.**  *$G$  contains no subgraph isomorphic to one of the configurations in Figure 1, where the vertices marked by  $\bullet$  have no other neighbors in  $G$ .*

**Proof.** The proof that  $G$  contains no subgraph isomorphic to one of the configurations in Figure 1(1)–(4) can be found in [8]. It remains to prove that  $G$  has no configurations depicted in Figure 1(5)–(13).

By the minimality of  $G$ , every proper subgraph of  $G$  has a total- $(m + 1)$ -coloring  $\varphi$  with the color set  $C = \{1, 2, \dots, m + 1\}$ . Erase the colors on all  $3^-$ -vertices. Let  $C(v) = \{\varphi(uv) : u \in N(v)\} \cup \{\varphi(v)\}$ .

Suppose that  $G$  contains a configuration depicted in Figure 1(5). Then  $G' = G - vv_6$  has a total-8-coloring  $\varphi$ . If  $\varphi(v_6x_5) \in C(v)$  or  $\varphi(v_6x_6) \in C(v)$ , then the forbidden colors for  $vv_6$  is at most 7, so  $vv_6$  can be properly colored. By recoloring the erased vertices, we obtain a total-8-coloring of  $G$ , a contradiction. So we can assume that  $\varphi(v_6x_5) \notin C(v)$  and  $\varphi(v_6x_6) \notin C(v)$ . Without loss of generality, assume that  $\varphi(v) = 6$ ,  $\varphi(v_6x_5) = 7$ ,  $\varphi(v_6x_6) = 8$ , and  $\varphi(vv_j) = j$  for  $j \in \{1, 2, \dots, 5\}$ . Then we recolor  $v$  with 7 or 8, and color  $vv_6$  with 6. By recoloring the erased vertices, we obtain a total-8-coloring of  $G$ , a contradiction.

Suppose that  $G$  contains a configuration depicted in Figure 1(6)–(13). Then  $G' = G - vv_7$  has a total-8-coloring  $\varphi$ . If  $\varphi(v_7x_7) \in C(v)$ , then the forbidden colors for  $vv_7$  is at most 7, so  $vv_7$  can be properly colored. By recoloring the erased vertices, we obtain a total-8-coloring of  $G$ , a contradiction. So we can assume that  $\varphi(vv_7) \notin C(v)$ . Without loss of generality, assume that  $\varphi(v) = 8$ ,  $\varphi(v_7x_7) = 7$ , and  $\varphi(vv_j) = j$  for  $j \in \{1, 2, \dots, 6\}$ . Thus, for each  $3^-$ -vertex  $v_k$  ( $1 \leq k \leq 7$ ), there is an edge incident with  $v_k$  colored 7, otherwise we can recolor  $vv_k$  with 7, and color  $vv_7$  with  $k$  to obtain a total-8-coloring of  $G$ , a contradiction.

For each 4-vertex  $v_i$  ( $1 \leq i \leq 6$ ), suppose its adjacent vertices are  $v, x_{i-1}, x_i, x_j$ . If  $\varphi(v_i) \neq 7$  ( $1 \leq i \leq 6$ ), then recolor  $v$  with 7, and color  $vv_7$  with 8. By recoloring the erased vertices, we obtain a total-8-coloring of  $G$ , a contradiction. Otherwise, there is at least one 4-vertex colored with 7. Suppose  $v$  is adjacent to only one 4-vertex  $v_i$  colored with 7. If  $|C(v_i)| < 8$ , then we recolor  $v_i$  with a color in  $C \setminus C(v_i)$ , recolor  $v$  with 7, and color  $vv_7$  with 8. Otherwise,  $|C(v_i)| = 8$ . If  $i \notin \{\varphi(x_{i-1}), \varphi(x_i), \varphi(x_j)\}$ , then we recolor  $v_i$  with  $i$ , recolor  $vv_i$  with 7, and color




Figure 1. Reducible configurations.

$vv_7$  with  $i$ . Otherwise,  $i \in \{\varphi(x_{i-1}), \varphi(x_i), \varphi(x_j)\}$ . Without loss of generality,  $\varphi(x_i) = i$ , then  $8 \notin \{C(v_i) \setminus \varphi(v)\}$ . Therefore, we recolor  $v_i$  with 8, recolor  $v$  with 7, and color  $vv_7$  with 8. Finally, we recolor the erased vertices, we obtain a total-8-coloring of  $G$ , a contradiction. Otherwise,  $v$  is adjacent to two or three 4-vertices colored with 7, then we take the same operations as above, respectively. Thus we can also obtain a total-8-coloring of  $G$ , a contradiction. ■

Let  $G = (V, E, F)$  be a graph which is embedded in a surface of nonnegative Euler characteristic. By Euler's formula  $|V| - |E| + |F| = \varepsilon$ , we have

$$\sum_{v \in V} (2d(v) - 6) + \sum_{f \in F} (d(f) - 6) = -6(|V| - |E| + |F|) = -6\varepsilon \leq 0.$$

Now we define the initial charge function  $ch(x)$  of  $x \in V \cup F$  to be  $ch(v) = 2d(v) - 6$  if  $v \in V$  and  $ch(f) = d(f) - 6$  if  $f \in F$ . It follows that  $\sum_{x \in V \cup F} ch(x) \leq 0$ . Now we design appropriate discharging rules and redistribute weights accordingly. Note that any discharging procedure preserves the total charge of  $G$ . If we can define suitable discharging rules to charge the initial charge function  $ch$  to the final charge function  $ch'$  on  $V \cup F$  such that  $\sum_{x \in V \cup F} ch'(x) > 0$ , then we get an obvious contradiction.

Our discharging rules are defined as follows.

- R1.** Every 2-vertex receives  $\frac{3}{2}$  from its child and  $\frac{1}{2}$  from its parent.
- R2.** Let  $f$  be a 3-face. If  $f$  is incident with a  $3^-$ -vertex, then it gets  $\frac{3}{2}$  from each of its incident  $6^+$ -vertices. If  $f$  is incident with a 4-vertex, then it gets  $\frac{1}{2}$  from the 4-vertex and gets  $\frac{5}{4}$  from each of its incident  $5^+$ -vertices. If  $f$  is not incident with any  $4^-$ -vertex, then it gets 1 from each of its incident  $5^+$ -vertices.
- R3.** Let  $f$  be a 4-face. If  $f$  is incident with two  $3^-$ -vertices, then it gets 1 from each of its two incident  $6^+$ -vertices. If  $f$  is incident with only one  $3^-$ -vertex and one 4-vertex, then it gets  $\frac{1}{2}$  from the incident 4-vertex and gets  $\frac{3}{4}$  from each of its two incident  $6^+$ -vertices. If  $f$  is incident with only one  $3^-$ -vertex and no 4-vertex, then it gets  $\frac{2}{3}$  from each of its incident  $5^+$ -vertices. If  $f$  is not incident with any  $3^-$ -vertex, then it gets  $\frac{1}{2}$  from each of its incident vertices.
- R4.** Every 5-face gets  $\frac{1}{3}$  from each of its incident  $4^+$ -vertices.

First, we begin to check  $ch'(x) \geq 0$  for all  $x \in V \cup F$ . By our discharging rules, it is easy to check that  $ch'(f) \geq 0$  for all  $f \in F$  and  $ch'(v) \geq 0$  for all 2-vertices  $v \in V$ . If  $d(v) = 3$ , then  $ch'(v) = ch(v) = 0$ . So it suffices to check that  $ch'(v) \geq 0$  for all  $4^+$ -vertices  $G$ .

Let  $v$  be a  $4^+$ -vertex of  $G$ . If  $d(v) = 4$ , then  $v$  sends at most  $\frac{1}{2}$  to each of its incident faces by R2 and R3, and it follows that  $ch'(v) \geq ch(v) - \frac{1}{2} \times 4 = 0$ . Suppose  $d(v) = 5$ . Then  $v$  sends at most  $\frac{5}{4}$  to each of its incident 3-faces by

R2, at most  $\frac{2}{3}$  to each of its incident  $4^+$ -faces by R3, at most  $\frac{1}{3}$  to each of its incident 5-faces by R4. By (a),  $f_3(v) \leq 2$ . If  $f_3(v) = 2$ , then  $v$  is incident with at least three  $5^+$ -faces, that is,  $f_{5^+}(v) \geq 3$ , and it follows that  $ch'(v) \geq ch(v) - \frac{5}{4} \times 2 - \frac{1}{3} \times 3 = \frac{1}{2} > 0$ . If  $f_3(v) = 1$ , then  $f_{5^+}(v) \geq 2$  and  $f_{4^+}(v) \leq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{5}{4} - \frac{1}{3} \times 2 - \frac{2}{3} \times 2 = \frac{3}{4} > 0$ . If  $f_3(v) = 0$ , then  $ch'(v) \geq ch(v) - \frac{2}{3} \times 5 = \frac{2}{3} > 0$ . Suppose  $d(v) = 6$ . Then  $f_3(v) \leq 3$  and  $v$  sends at most  $\frac{3}{2}$  to each of its incident 3-faces by R2, at most 1 to each of its incident  $4^+$ -faces by R3, at most  $\frac{1}{3}$  to each of its incident 5-faces by R4. Thus, if  $1 \leq f_3(v) \leq 3$ , then by the similar argument as above, we have  $ch'(v) \geq ch(v) - \max\{\frac{3}{2} \times 3 + \frac{1}{3} \times 3, \frac{3}{2} \times 2 + \frac{1}{3} \times 3 + 1, \frac{3}{2} + \frac{1}{3} \times 2 + 1 \times 3\} = \frac{1}{2} > 0$ . Otherwise,  $v$  is adjacent to at least one  $4^+$ -vertex or incident with at least one  $5^+$ -face by Lemma 4, configuration (5). If  $v$  is adjacent to at least one  $4^+$ -vertex, then  $ch'(v) \geq ch(v) - 2 \times \frac{3}{4} - 4 = \frac{1}{2} > 0$  by R3. If  $v$  is incident with at least one  $5^+$ -face, then  $ch'(v) \geq ch(v) - \frac{1}{3} - 5 = \frac{2}{3} > 0$  by R4.

Suppose  $d(v) = d \geq 7$ . Let  $N(v) = \{v_1, v_2, \dots, v_d\}$  and  $f_1, f_2, \dots, f_d$  be faces incident with  $v$  in the clockwise order, where  $f_i$  is incident with  $v_i$  and  $v_{i+1}$ , for  $i \in \{1, 2, \dots, d\}$ , where all the subscripts here are taken modulo  $d$ . If  $n_2(v) \geq 1$ , then  $v$  sends at most  $\frac{n_2(v)+2}{2}$  to all its adjacent 2-vertices by R1, at most  $\frac{3}{2}$  to each of its incident 3-faces by R2, at most 1 to each of its incident  $4^+$ -faces by R3, at most  $\frac{1}{3}$  to each of its incident 5-faces by R4.

**Lemma 5.** *Suppose that  $d(v_i) = d(v_k) = 2$  and  $d(v_j) \geq 3$  for all  $j = i + 1, \dots, k - 1$ . If  $f_i, f_{i+1}, \dots, f_{k-1}$  are  $4^+$ -faces, then  $v$  sends at most  $\frac{3}{2} + (k - 3)$  (in total) to  $f_i, f_{i+1}, \dots, f_{k-1}$ .*

**Proof.** By Lemma 3,  $\max\{d(v_{i+1}), \dots, d(v_{k-1})\} \geq 4$  or  $\max\{d(f_1), \dots, d(f_{k-1})\} \geq 5$ . If  $\max\{d(v_{i+1}), \dots, d(v_{k-1})\} \geq 4$ , then  $v$  sends at most  $2 \times \frac{3}{4} + (k - 1 - 2)$  to  $f_i, \dots, f_{k-1}$  by R2. If  $\max\{d(f_1), \dots, d(f_{k-1})\} \geq 5$ , then  $v$  sends at most  $\frac{1}{3} + (k - 1 - 1)$  to  $f_i, \dots, f_{k-1}$  by R3 and R4. Since  $2 \times \frac{3}{4} > 1 + \frac{1}{3}$ ,  $v$  sends at most  $\frac{3}{2} + (k - 3)$  to  $f_i, f_{i+1}, \dots, f_{k-1}$ . ■

*Case 1.*  $\Delta(G) = 7$ . Let  $v$  be a 7-vertex. Then  $ch(v) = 2 \times 7 - 6 = 8$ . We consider the following cases.

*Subcase 1.1.*  $n_2(v) = 6$ . By (c), any 2-vertex is not incident with a 4-face. Moreover,  $t = f_3(v) = 0$  and  $f_{6^+}(v) \geq 5$  by Lemma 4, configurations (1) and (4). So  $ch'(v) \geq ch(v) - \frac{6+2}{2} - 2 = 2 > 0$ .

*Subcase 1.2.*  $n_2(v) = 5$ . Then  $t \leq 1$  by Lemma 4, configurations (1) and (4). If  $t = 0$ , then  $f_{6^+}(v) \geq 3$  and  $f_{4^+}(v) \leq 4$  by Lemma 4, configuration (4), and it follows that  $ch'(v) \geq ch(v) - \frac{5+2}{2} - 4 \times 1 = \frac{1}{2} > 0$ . Otherwise,  $f_{6^+}(v) \geq 4$  and  $f_{4^+}(v) \leq 2$ . Thus  $ch'(v) \geq ch(v) - \frac{5+2}{2} - \frac{3}{2} - 2 \times 1 = 1 > 0$ .

*Subcase 1.3.*  $n_2(v) = 4$ . There are four possible configurations as shown in Figure 2.




Figure 2.  $n_2(v) = 4$ .

For Figure 2(a),  $t \leq 1$  and  $f_{6+}(v) \geq 3$  by Lemma 4, configurations (1) and (4). If  $t = 1$ , then  $f_{5+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{4+t}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 1 = \frac{11}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{4+t}{2} - \frac{3}{2} - 2 = \frac{3}{2} > 0$  by Lemma 5.

For Figure 2(b) and (c),  $t \leq 1$  and  $f_{6+}(v) \geq 2$  by Lemma 4, configurations (1) and (4). If  $t = 1$ , then  $f_{5+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{4+t}{2} - \frac{3}{2} - 3 \times \frac{1}{3} - \frac{3}{2} = \frac{4}{3} > 0$  by Lemma 5. Otherwise,  $ch'(v) \geq ch(v) - \frac{4+t}{2} - \frac{3}{2} - \frac{3}{2} - 1 = 1 > 0$  by Lemma 5.

For Figure 2(d),  $t = 0$  and  $f_{6+}(v) \geq 1$  by Lemma 4, configurations (1) and (4). Then  $ch'(v) \geq ch(v) - \frac{4+t}{2} - 3 \times \frac{3}{2} = \frac{1}{2} > 0$  by Lemma 5.

*Subcase 1.4.*  $n_2(v) = 3$ . There are four possible configurations as shown in Figure 3.




Figure 3.  $n_2(v) = 3$ .

For Figure 3(a),  $t \leq 2$  and  $f_{6+}(v) \geq 2$  by Lemma 4, configurations (1) and (4). If  $t = 2$ , then  $f_{5+}(v) \geq 3$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+t}{2} - 2 \times \frac{3}{2} - 3 \times \frac{1}{3} = \frac{3}{2} > 0$ . If  $t = 1$ , then  $f_{5+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+t}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 2 = \frac{4}{3} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{3+t}{2} - \frac{3}{2} - 3 = 1 > 0$  by Lemma 5.

For Figure 3(b),  $t \leq 1$  and  $f_{6+}(v) \geq 1$  by Lemma 4, configurations (1) and (4). If  $t = 1$ , then  $f_{5+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+t}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 1 - \frac{3}{2} = \frac{5}{6} > 0$  by Lemma 5. Otherwise,  $ch'(v) \geq ch(v) - \frac{3+t}{2} - \frac{3}{2} - \frac{3}{2} - 2 = \frac{1}{2} > 0$  by Lemma 5.

For Figure 3(c),  $t \leq 2$  and  $f_{6+}(v) \geq 1$  by Lemma 4, configurations (1) and (4). If  $t = 2$ , then  $f_{5+}(v) \geq 4$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+t}{2} - 2 \times$



$\frac{3}{2} - 4 \times \frac{1}{3} = \frac{7}{6} > 0$ . If  $t = 1$ , then  $f_{5^+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - \frac{3}{2} - 1 = \frac{5}{6} > 0$  by Lemma 5. Otherwise,  $ch'(v) \geq ch(v) - \frac{3+2}{2} - 2 \times (\frac{3}{2} + 1) = \frac{1}{2} > 0$  by Lemma 5.

For Figure 3(d),  $t \leq 1$  by Lemma 4, configuration (1). If  $t = 1$ , then  $f_{5^+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 2 \times \frac{3}{2} = \frac{1}{3} > 0$  by Lemma 5. Otherwise,  $t = 0$ . If  $v$  is adjacent to at least four  $4^+$ -vertices, then  $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - \frac{3}{2} - 3 \times \frac{3}{4} = \frac{1}{4} > 0$  by Lemma 5. Otherwise,  $v$  is adjacent to at least one  $5^+$ -vertex or incident with at least one  $5^+$ -face by Lemma 4, configuration (6), and Lemma 3. If  $v$  is adjacent to at least one  $5^+$ -vertex, then  $ch'(v) \geq ch(v) - \frac{3+2}{2} - \max\{2 \times \frac{2}{3} + 1 + 2 \times \frac{3}{2}, \frac{3}{2} + 1 + 2 \times \frac{2}{3} + \frac{3}{2}\} = \frac{1}{6} > 0$  by R3 and Lemma 5. If  $v$  is incident with at least one  $5^+$ -face, then  $ch'(v) \geq ch(v) - \frac{3+2}{2} - \max\{\frac{1}{3} + 2 + 2 \times \frac{3}{2}, \frac{3}{2} + 1 + \frac{1}{3} + \frac{3}{2} + 1\} = \frac{1}{6} > 0$  by R4 and Lemma 5.

*Subcase 1.5.*  $n_2(v) = 2$ . There are three possible configurations as shown in Figure 4.




Figure 4.  $n_2(v) = 2$ .

For Figure 4(a), we have that the face  $f_6$  is a  $5^+$ -face, and it follows that  $f_6$  receives at most  $\frac{1}{3}$  from  $v$  by R4. So  $ch(v) - \frac{2+2}{2} - \frac{1}{3} = 8 - 2 - \frac{1}{3} = \frac{17}{3}$ . Moreover,  $t \leq 2$  by Lemma 4, configuration (1), and (a). If  $1 \leq t \leq 2$ , then  $f_{5^+}(v) \geq (t + 1)$  (except  $f_6$ ), and it follows that  $ch'(v) \geq \frac{17}{3} - t \times \frac{3}{2} - (t + 1) \times \frac{1}{3} - (7 - 1 - t - t - 1) = \frac{2+t}{6} > 0$ . Otherwise,  $ch'(v) \geq \frac{17}{3} - (\frac{3}{2} + 4) = \frac{1}{6} > 0$ .

For Figure 4(b),  $t \leq 2$  by Lemma 4, configuration (1), and (a). If  $t = 2$ , then  $f_{5^+}(v) \geq 3$ , it follows that  $ch'(v) \geq ch(v) - \frac{2+2}{2} - 2 \times \frac{3}{2} - 3 \times \frac{1}{3} - \frac{3}{2} = \frac{1}{2} > 0$ . If  $t = 1$ , then  $f_{5^+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - \frac{1}{3} \times 2 - 2 \times 1 - \frac{3}{2} = \frac{1}{3} > 0$ . Suppose  $t = 0$ . If  $v$  is adjacent to at least three  $4^+$ -vertices, then  $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - 3 \times \frac{3}{4} - 2 = \frac{1}{4} > 0$  by Lemma 5. Otherwise,  $v$  is adjacent to at least one  $5^+$ -vertex or incident with at least one  $5^+$ -face by Lemma 4, configurations (7)–(8). By the same argument as above, we can obtain  $ch'(v) \geq ch(v) - \frac{2+2}{2} - \max\{\frac{3}{2} + 3 + 2 \times \frac{2}{3}, \frac{3}{2} + 4 + \frac{1}{3}\} = \frac{1}{6} > 0$ .

For Figure 4(c),  $t \leq 2$  by Lemma 4, configuration (1), and (a). If  $t = 2$ , then  $f_{5^+}(v) \geq 4$ , it follows that  $ch'(v) \geq ch(v) - \frac{2+2}{2} - 2 \times \frac{3}{2} - 4 \times \frac{1}{3} - 1 = \frac{2}{3} > 0$ . If  $t = 1$ , then  $f_{5^+}(v) \geq 2$ , and it follows that  $ch'(v) \geq ch(v) - \frac{2+2}{2} -$

$\max \left\{ \frac{3}{2} + \frac{1}{3} \times 2 + \frac{3}{2} + 2, \frac{3}{2} + \frac{1}{3} \times 2 + 1 + \frac{3}{2} + 1 \right\} = \frac{1}{3} > 0$ . Suppose  $t = 0$ . If  $v$  is adjacent to at least three  $4^+$ -vertices, then  $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - 3 \times \frac{3}{4} - 2 = \frac{1}{4} > 0$  by Lemma 5. Otherwise,  $v$  is adjacent to at least one  $5^+$ -vertex or incident with at least one  $5^+$ -face by Lemma 4, configurations (9)–(10). By the same argument as above, we can obtain  $ch'(v) \geq ch(v) - \frac{2+2}{2} - \max \left\{ \frac{3}{2} + 3 + 2 \times \frac{2}{3}, \frac{3}{2} + 4 + \frac{1}{3} \right\} = \frac{1}{6} > 0$ .

*Subcase 1.6.*  $n_2(v) = 1$ . Note that  $n_{4^+}(v) \geq 1$  by (e) and  $t \leq 3$  by (a). Suppose  $t = 0$ . If  $v$  is adjacent to at least two  $4^+$ -vertices, then  $ch'(v) \geq ch(v) - \frac{1+2}{2} - 3 \times \frac{3}{4} - 4 = \frac{1}{4} > 0$ . Otherwise,  $v$  is adjacent to at least one  $5^+$ -vertex or incident with at least one  $5^+$ -face by Lemma 4, configurations (11)–(13), then  $ch'(v) \geq ch(v) - \frac{1+2}{2} - 2 \times \frac{2}{3} - 5 = \frac{1}{6} > 0$  by R3 or  $ch'(v) \geq ch(v) - \frac{1+2}{2} - \frac{1}{3} - 6 = \frac{1}{6} > 0$  by R4. Suppose  $1 \leq t \leq 3$ . If  $v$  is incident with a  $(2, 7, 7)$ -face, then the other face incident with the 2-vertex is a  $6^+$ -face. Moreover, the other 3-faces incident with  $v$  are  $(4, 5^+, 5^+)$ -faces by Lemma 4, configuration (3), and  $v$  is incident with at least  $t$   $5^+$ -faces. Then we can obtain that  $ch'(v) \geq ch(v) - \frac{1+2}{2} - \frac{3}{2} - (t-1) \times \frac{5}{4} - \frac{1}{3} \times t - (7-1-t-t) = \frac{3+5t}{12} > 0$ , where  $v$  sends at most  $\frac{5}{4}$  to each of its incident  $(4, 5^+, 5^+)$ -faces by R2. Otherwise, 2-vertex is not incident with any 3-face. Then  $v$  is incident with at least  $(t+1)$   $5^+$ -faces, and it follows that  $ch'(v) \geq ch(v) - \frac{1+2}{2} - t \times \frac{3}{2} - \frac{1}{3} \times (t+1) - (7-t-t-1) = \frac{1+t}{6} > 0$ .

*Subcase 1.7.*  $n_2(v) = 0$ . Note that  $t \leq 3$  by (a). If  $t = 0$ , then  $ch'(v) \geq ch(v) - 7 \times 1 = 1 > 0$ . Otherwise, by the same argument as above, we can obtain that  $ch'(v) \geq ch(v) - t \times \frac{3}{2} - \frac{1}{3} \times (t+1) - (7-t-t-1) = \frac{10+t}{6} > 0$ .

*Case 2.*  $\Delta(G) \geq 8$ . In [23], Theorem 1 was established for  $\Delta \geq 9$ . So we assume that  $\Delta = 8$ . Then  $ch(v) = 2 \times 8 - 6 = 10$ . By the same argument as above, we consider the following cases.

*Subcase 2.1.*  $n_2(v) = 7$ . Then  $t = 0$  and  $f_{6^+}(v) \geq 6$  by Lemma 4, configurations (1) and (4). So  $ch'(v) \geq ch(v) - \frac{7+2}{2} - 2 = \frac{7}{2} > 0$ .

*Subcase 2.2.*  $n_2(v) = 6$ . Then  $t \leq 1$  and  $f_{6^+}(v) \geq 4$  by Lemma 4, configurations (1) and (4). If  $t = 0$ , then  $ch'(v) \geq ch(v) - \frac{6+2}{2} - 4 = 2 > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{6+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} = \frac{23}{6} > 0$ .

*Subcase 2.3.*  $n_2(v) = 5$ . Then  $t \leq 1$  and  $f_{6^+}(v) \geq 2$  by Lemma 4, configurations (1) and (4). If  $t = 0$ , then  $ch'(v) \geq ch(v) - \frac{5+2}{2} - 6 = \frac{1}{2} > 0$ . Otherwise,  $f_{5^+}(v) \geq 2$ . Thus  $ch'(v) \geq ch(v) - \frac{5+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 3 = \frac{4}{3} > 0$ .

*Subcase 2.4.*  $n_2(v) = 4$ . Then  $t \leq 2$  by Lemma 4, configuration (1). If  $1 \geq t \geq 2$ , then  $f_{6^+}(v) \geq 1$  and  $f_{5^+}(v) \geq t+1$ , and it follows that  $ch'(v) \geq ch(v) - \frac{4+2}{2} - t \times \frac{3}{2} - (t+1) \times \frac{1}{3} - (8-1-t-t-1) = \frac{4+t}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{4+2}{2} - \max \left\{ \frac{3}{2} + 3, \frac{3}{2} \times 2 + 2, \frac{3}{2} \times 3 + 1, \frac{3}{2} \times 4 \right\} = 1 > 0$ .

*Subcase 2.5.*  $n_2(v) = 3$ . Then  $t \leq 2$  by Lemma 4 configuration (1). If  $1 \geq t \geq 2$ , then  $f_{5^+}(v) \geq t+1$ , and it follows that  $ch'(v) \geq ch(v) - \frac{3+2}{2} - t \times \frac{3}{2} -$

$(t+1) \times \frac{1}{3} - (8-t-t-1) = \frac{1+t}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{3+t}{2} - \max\{\frac{3}{2} + 4, \frac{3}{2} \times 2 + 3, \frac{3}{2} \times 3 + 2\} = 1 > 0$ .

*Subcase 2.6.*  $n_2(v) = 2$ . Then  $t \leq 2$  by Lemma 4 configuration (1). If  $1 \geq t \geq 2$ , then  $f_{5^+}(v) \geq t+1$ , and it follows that  $ch'(v) \geq ch(v) - \frac{2+t}{2} - t \times \frac{3}{2} - (t+1) \times \frac{1}{3} - (8-t-t-1) = \frac{4+t}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{2+t}{2} - \max\{\frac{3}{2} + 5, \frac{3}{2} \times 2 + 4\} = 1 > 0$ .

*Subcase 2.7.*  $n_2(v) = 1$ . Then  $t \leq 4$  by (a). If  $1 \geq t \geq 4$ , then  $f_{5^+}(v) \geq t$ , and it follows that  $ch'(v) \geq ch(v) - \frac{1+t}{2} - t \times \frac{3}{2} - t \times \frac{1}{3} - (8-t-t) = \frac{3+t}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - \frac{1+t}{2} - 8 = \frac{1}{2} > 0$ .

*Subcase 2.8.*  $n_2(v) = 0$ . Then  $t \leq 4$  by (a). If  $1 \geq t \geq 4$ , then  $f_{5^+}(v) \geq t$ , and it follows that  $ch'(v) \geq ch(v) - t \times \frac{3}{2} - t \times \frac{1}{3} - (8-t-t) = \frac{12+t}{6} > 0$ . Otherwise,  $ch'(v) \geq ch(v) - 8 = 2 > 0$ .

Finally, according to the above argument, we have checked  $ch'(x) \geq 0$  for all  $x \in V \cup F$  and  $ch'(x) > 0$  for any  $5^+$ -vertex  $x \in V$ . By Lemma 1, we have  $\Delta(G) \geq 6$ . So  $\sum_{x \in V \cup F} ch'(x) > 0$ . Hence we complete the proof of Theorem 2.

## REFERENCES

- [1] M. Behzad, *Graphs and Their Chromatic Numbers* (Ph.D. Thesis, Michigan State University, 1965).
- [2] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications* (Macmillan Press Ltd., London, 1976).
- [3] O.V. Borodin, *On the total coloring of planar graphs*, J. Reine Angew. Math. **394** (1989) 180–185.
- [4] O.V. Borodin, *Coupled colourings of graphs on a plane*, Metody Diskret. Anal. **45** (1987) 21–27, in Russian.
- [5] O.V. Borodin, A.V. Kostochka and D.R. Woodall, *List edge and list total colourings of multigraphs*, J. Combin. Theory Ser. B **71** (1997) 184–204.  
doi:10.1006/jctb.1997.1780
- [6] O.V. Borodin A.V. Kostochka and D.R. Woodall, *Total colorings of planar graphs with large maximum degree*, J. Graph Theory **26** (1997) 53–59.  
doi:10.1002/(SICI)1097-0118(199709)26:1<53::AID-JGT6>3.0.CO;2-G
- [7] G.J. Chang, J. Hou and N. Roussel, *Local condition for planar graphs of maximum degree 7 to be 8-totally colorable*, Discrete Appl. Math. **159** (2011) 760–768.  
doi:10.1016/j.dam.2011.01.001
- [8] D. Du, L. Shen and Y. Wang, *Planar graphs with maximum degree 8 and without adjacent triangles are 9-totally-colorable*, Discrete Appl. Math. **157** (2009) 2778–2784.  
doi:10.1016/j.dam.2009.02.011
- [9] T.R. Jensen and B. Toft, *Graph Coloring Problems* (Wiley Interscience, 1995).

- [10] L. Kowalik, J.-S. Sereni and R. Škrekovski, *Total-colorings of plane graphs with maximum degree nine*, SIAM J. Discrete Math. **22** (2008) 1462–1479.  
doi:10.1137/070688389
- [11] L. Shen and Y.Q. Wang, *Total colorings of planar graphs with maximum degree at least 8*, Sci. China Ser A: Math. **52** (2009) 1733–1742.  
doi:10.1007/s11425-008-0155-3
- [12] L. Shen and Y. Wang, *On the 7 total colorability of planar graphs with maximum degree 6 and without 4-cycles*, Graphs Combin. **25** (2009) 401–407.  
doi:10.1007/s00373-009-0843-y
- [13] A.V. Kostochka, *The total coloring of a multigraph with maximal degree 4*, Discrete Math. **17** (1977) 161–163.  
doi:10.1016/0012-365X(77)90146-7
- [14] A.V. Kostochka, *An analogue of Shannon's estimate for complete colorings*, Metody Diskret. Anal. **30** (1977) 13–22, in Russian.
- [15] A.V. Kostochka, *The total chromatic number of any multigraph with maximum degree five is at most seven*, Discrete Math. **162** (1996) 199–214.  
doi:10.1016/0012-365X(95)00286-6
- [16] B. Liu, J.F. Hou, J.L. Wu and G.Z. Liu, *Total colorings and list total colorings of planar graphs without intersecting 4-cycles*, Discrete Math. **309** (2009) 6035–6043.  
doi:10.1016/j.disc.2009.05.006
- [17] D.P. Sanders and Y. Zhao, *On total 9-coloring planar graphs of maximum degree seven*, J. Graph Theory **31** (1999) 67–73.  
doi:10.1002/(SICI)1097-0118(199905)31:1(67::AID-JGT6)3.0.CO;2-C
- [18] V.G. Vizing, *Some unsolved problems in graph theory*, Uspekhi Mat. Nauk **23** (1968) 117–134, in Russian.
- [19] B. Wang and J.-L. Wu, *Total coloring of planar graphs with maximum degree seven*, Inform. Process. Lett. **111** (2011) 1019–1021.  
doi:10.1016/j.ipl.2011.07.012
- [20] P. Wang and J.-L. Wu, *A note on total colorings of planar graphs without 4-cycles*, Discuss. Math. Graph Theory **24** (2004) 125–135.  
doi:10.7151/dmgt.1219
- [21] H.J. Wang, L.D. Wu, W.L. Wu, P.M. Pardalos and J.L. Wu, *Minimum total coloring of planar graph*, J. Global Optim. **60** (2014) 777–791.  
doi:10.1007/s10898-013-0138-y
- [22] H.J. Wang, B. Liu, J.L. Wu and G.Z. Liu, *Total coloring of embedded graphs with maximum degree at least seven*, Theoret. Comput. Sci. **518** (2014) 1–9.  
doi:10.1016/j.tcs.2013.04.030
- [23] H.J. Wang, B. Liu, J.L. Wu and B. Wang, *Total coloring of graphs embedded in surfaces of nonnegative Euler characteristic*, Sci. China Math. **57** (2014) 211–220.  
doi:10.1007/s11425-013-4576-2

- [24] J.L. Wu and P. Wang, *List-edge and list-total colorings of graphs embedded on hyperbolic surfaces*, *Discrete Math.* **308** (2008) 6210–6215.  
doi:10.1016/j.disc.2007.11.044
- [25] W.F. Wang, *Total chromatic number of planar graphs with maximum degree ten*, *J. Graph Theory* **54** (2007) 91–102.  
doi:10.1002/jgt.20195

Received 30 September 2016

Revised 30 March 2017

Accepted 30 March 2017