

## **$\mathcal{P}$ -BIPARTITIONS OF MINOR HEREDITARY PROPERTIES**

PIOTR BOROWIECKI

*Institute of Mathematics*  
*Technical University*  
*Podgórna 50, 65-246 Zielona Góra, Poland*  
**e-mail:** p.borowiecki@im.pz.zgora.pl

AND

JAROSLAV IVANČO

*Department of Geometry and Algebra*  
*P.J. Šafárik University*  
*Jesenná 5, 041 54 Košice, Slovakia*  
**e-mail:** ivanco@duro.upjs.sk

### **Abstract**

We prove that for any two minor hereditary properties  $\mathcal{P}_1$  and  $\mathcal{P}_2$ , such that  $\mathcal{P}_2$  covers  $\mathcal{P}_1$ , and for any graph  $G \in \mathcal{P}_2$  there is a  $\mathcal{P}_1$ -bipartition of  $G$ . Some remarks on minimal reducible bounds are also included.

**Keywords:** minor hereditary property of graphs, generalized colouring, bipartitions of graphs.

**1991 Mathematics Subject Classification:** 05C70, 05C15.

### 1. INTRODUCTION AND NOTATION

According to [3] we denote by  $\mathcal{I}$  the class of all finite simple graphs. A *graph property* is a nonempty isomorphism-closed subclass of  $\mathcal{I}$ . We also say that a graph has the property  $\mathcal{P}$  if  $G \in \mathcal{P}$ . For properties  $\mathcal{P}_1, \mathcal{P}_2$  of graphs a vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -*partition* of a graph  $G$  is a partition  $(V_1, V_2)$  of  $V(G)$  such that the subgraph  $G[V_i]$  induced by the set  $V_i$  has the property  $\mathcal{P}_i$  for each  $i = 1, 2$ . The class of all vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partitionable graphs is denoted by  $\mathcal{P}_1 \circ \mathcal{P}_2$ . If  $\mathcal{P}_1 = \mathcal{P}_2 = \mathcal{P}$ , then a  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition (as in [4]) we call a  $\mathcal{P}$ -*bipartition*.

Let be given a graph  $G \in \mathcal{I}$ . A *contraction* of the graph  $G$  is a graph obtained from  $G$  by repeated contractions of edges, where *contraction of an edge*  $(v_1, v_2)$  of the graph  $G$  is obtained by deleting  $v_1$  and  $v_2$  and all incident edges from  $G$  and adding a new vertex  $u$  and all the edges required to satisfy the following condition  $N(u) = N(v_1) \cup N(v_2) \setminus \{v_1, v_2\}$ .

A graph  $H$  obtained from  $G$  by deletions of vertices or edges, or contractions of edges is called a *minor* of  $G$ . So, the graph  $H$  is a minor of the graph  $G$  if  $H$  is a subgraph of  $G$  or can be obtained from a subgraph of  $G$  by contractions of edges. We express this relation between the graphs  $H$  and  $G$  by  $H < G$ .

A property  $\mathcal{P}$  of graphs is called *minor hereditary* (*hereditary*) if it is closed under minors (subgraphs), i.e., if whenever  $G \in \mathcal{P}$  and  $H$  is a minor (subgraph) of  $G$ , then also  $H \in \mathcal{P}$ .

Any minor hereditary property  $\mathcal{P}$  can be uniquely determined by the set of *forbidden minors* which can be defined in the following way:

$$\mathbf{F}_M(\mathcal{P}) = \{G \in \mathcal{I} : G \notin \mathcal{P} \text{ but each minor } H \text{ of } G, H \neq G, \text{ belongs to } \mathcal{P}\}.$$

A property  $\mathcal{P}$  is called *additive* if it is closed under disjoint union of graphs, i.e., if for each graph  $G$  all of whose connected components have a property  $\mathcal{P}$  it follows that  $G$  has a property  $\mathcal{P}$ , too. It is easy to see that a minor hereditary property  $\mathcal{P}$  is additive if and only if all minors  $H \in \mathbf{F}_M(\mathcal{P})$  are connected.

Many well-known properties of graphs are both minor hereditary and additive. According to [2], [3] we list some of them to introduce the necessary notions which will be used in the paper. It is convenient to work with an arbitrary nonnegative integer  $k$ .

$$\begin{aligned} \mathcal{O} &= \{G \in \mathcal{I} : G \text{ is edgeless, i.e., } E(G) = \emptyset\}, \\ \mathcal{D}_1 &= \{G \in \mathcal{I} : G \text{ is 1-degenerate, i.e., the minimum degree } \delta(H) \leq 1 \\ &\quad \text{for each } H \subseteq G\}, \\ \mathcal{T}_k &= \{G \in \mathcal{I} : G \text{ contains no subgraph homeomorphic to } K_{k+2} \\ &\quad \text{or } K_{\lfloor \frac{k+3}{2} \rfloor, \lceil \frac{k+3}{2} \rceil}, k \leq 3\}, \\ \mathcal{SP} &= \{G \in \mathcal{I} : G \text{ contains no subgraph homeomorphic to } K_4\}. \end{aligned}$$

We have  $\mathcal{D}_1 = \mathcal{T}_1$  to be the class of all forests,  $\mathcal{T}_2$  and  $\mathcal{T}_3$  the class of all outerplanar and all planar graphs, respectively and  $\mathcal{SP}$  the class of all series-parallel graphs.

For the properties given above we have:

$$\begin{aligned} \mathbf{F}_M(\mathcal{O}) &= \{K_2\}, \\ \mathbf{F}_M(\mathcal{D}_1) &= \{K_3\}, \end{aligned}$$

$$\begin{aligned} \mathbf{F}_M(\mathcal{T}_2) &= \{K_4, K_{2,3}\}, \\ \mathbf{F}_M(\mathcal{T}_3) &= \{K_5, K_{3,3}\}, \\ \mathbf{F}_M(\mathcal{SP}) &= \{K_4\}. \end{aligned}$$

Let us define the next properties.

$$\begin{aligned} \mathbf{F}_M(\mathcal{LF}) &= \{K_3, K_{1,3}\}, \\ \mathbf{F}_M(\mathcal{S}) &= \{K_4, K_{1,3} + K_1\}. \end{aligned}$$

All additive minor hereditary (hereditary) properties of graphs, partially ordered by a set-inclusion, form a lattice  $\mathbf{T}^a$ , ( $\mathbf{L}^a$ ) with  $\cap$  as a meet operation and  $\mathcal{O}$  as the smallest element (see [2]).

All the above listed properties form in  $\mathbf{T}^a$  the following chain:

$$\mathcal{O} \subset \mathcal{LF} \subset \mathcal{D}_1 \subset \mathcal{T}_2 \subset \mathcal{S} \subset \mathcal{SP} \subset \mathcal{T}_3.$$

## 2. $\mathcal{P}$ -BIPARTITION THEOREM

**Definition.** Let  $\mathcal{P}_1$  and  $\mathcal{P}_2$  be two additive minor hereditary properties. We say that  $\mathcal{P}_2$  covers  $\mathcal{P}_1$  whenever for every graph  $G_1 \in \mathbf{F}_M(\mathcal{P}_1)$  there exists a graph  $G_2 \in \mathbf{F}_M(\mathcal{P}_2)$  such that  $G_2 - v$  is a minor of  $G_1$  for some vertex  $v \in V(G_2)$ .

**Theorem 1.** *If  $\mathcal{P}_2$  covers  $\mathcal{P}_1$ , then the vertex set of a graph  $G \in \mathcal{P}_2$  can be partitioned into two subsets such that each of them induces a subgraph of  $G$  belonging to  $\mathcal{P}_1$ .*

**Proof.** Let us consider a given graph  $G \in \mathcal{P}_2$  with an arbitrarily chosen vertex  $v$ . It is sufficient to consider a case when  $G$  is connected. We define the subsets  $U_k = \{u \in V(G) : d(v, u) = k\}$ , where  $d(u, v)$  is the length of the shortest path between  $v$  and  $u$ . Put  $e = \max\{k : U_k \neq \emptyset\}$ . Then  $U_0, U_1, \dots, U_e$  is a partition of  $V(G)$  into  $e + 1$  pairwise disjoint subsets. Moreover, a subgraph induced by  $U_0 = \{v\}$  belongs to  $\mathcal{P}_1$ . Now, let us assume to the contrary, that one of the subsets  $U_k$ ,  $k = 1, \dots, e$ , induces a subgraph of  $G$ , which is not in  $\mathcal{P}_1$ . Thus there is a minor  $H$  of  $G[U_k]$  belonging to  $\mathbf{F}_M(\mathcal{P}_1)$ . Since the subgraph of  $G$  induced by  $U' = \bigcup_{i=0}^{k-1} U_i$  is connected and every vertex of  $U_k$  is adjacent to a vertex of  $U_{k-1} \subseteq U'$ , then the graph  $H + K_1$  is a minor of  $G$ . Since  $\mathcal{P}_2$  covers  $\mathcal{P}_1$ , then  $\mathbf{F}_M(\mathcal{P}_2)$  contains a graph  $H'$  such that  $H' - u$  is a minor of  $H$ , for some  $u \in V(H')$ . Obviously,  $H'$  is a minor of  $H + K_1$ . Hence, since  $H + K_1$  is a minor of  $G$ , then  $H'$  is a minor of  $G$ , contrary to  $G \in \mathcal{P}_2$ . Therefore, each of the subsets  $U_i, i = 0, 1, \dots, e$  induces a subgraph of  $G$  belonging to  $\mathcal{P}_1$ . Since vertices

$u \in U_i$  and  $w \in U_j$ , for  $|i - j| > 1$  are non-adjacent in  $G$ , then both of the sets  $V_1 = \bigcup_{i=1}^{\lceil e/2 \rceil} U_{2i-1}$  and  $V_2 = \bigcup_{i=0}^{\lfloor e/2 \rfloor} U_{2i}$  induce subgraphs of  $G$  belonging to  $\mathcal{P}_1$ , i.e., the partition  $(V_1, V_2)$  is the required  $\mathcal{P}_1$ -bipartition of  $V(G)$ . ■

From the theorem given above, a series of well-known results follows:

- (a)  $\mathcal{D}_1 \subset \mathcal{O}^2$ ,
- (b)  $\mathcal{T}_2 \subset \mathcal{LF}^2$   
proved by Mihók [10], Broere and Mynhardt [5], Wang [13], and Goddard [8],
- (c)  $\mathcal{SP} \subset \mathcal{D}_1^2$   
which is the result of Dirac [7],
- (d)  $\mathcal{T}_3 \subset \mathcal{T}_2^2$   
proved by Broere and Mynhardt [5], Wang [13] and Poh [12].

The new conclusions can be drawn, too. For the class  $\mathcal{S}$  defined by  $\mathbf{F}_M(\mathcal{S}) = \{K_4, K_{1,3} + K_1\}$  we have:

- (e)  $\mathcal{S} \subset \mathcal{LF}^2$ .

### 3. MINIMAL REDUCIBLE BOUNDS

An additive hereditary property  $\mathcal{R}$  is called *reducible* in  $\mathbb{L}^a$ , if there exist additive hereditary properties  $\mathcal{P}_1, \mathcal{P}_2$  such that  $\mathcal{P} = \mathcal{P}_1 \circ \mathcal{P}_2$ , and it is called *irreducible*, otherwise.

For a given property  $\mathcal{P}$ , a reducible property  $\mathcal{R}$  is called *minimal reducible bound* for  $\mathcal{P}$  if  $\mathcal{P} \subset \mathcal{R}$  and there is no reducible property  $\mathcal{R}' \subset \mathcal{R}$  satisfying  $\mathcal{P} \subseteq \mathcal{R}'$ . The set of all minimal reducible bounds for  $\mathcal{P}$  will be denoted by  $\mathbf{B}(\mathcal{P})$ . The notion of minimal reducible bounds have been introduced in [11]. In this paper Mihók proved that the class  $\mathcal{T}_2$  of outerplanar graphs has exactly two minimal reducible bounds, i.e.,  $\mathbf{B}(\mathcal{T}_2) = \{\mathcal{LF}^2, \mathcal{O} \circ \mathcal{D}_1\}$ . A similar results for  $\mathcal{SP}$  and  $\mathcal{D}_2$  can be found in [1], namely,  $\mathbf{B}(\mathcal{SP}) = \mathbf{B}(\mathcal{D}_2) = \{\mathcal{O} \circ \mathcal{D}_1\}$ .

By the transitivity and Mihók's proof (see [11]) we have the following minimal reducible bounds for the property  $\mathcal{S} \supset \mathcal{T}_2$ .

**Theorem 2.**  $\mathbf{B}(\mathcal{S}) = \{\mathcal{LF}^2, \mathcal{O} \circ \mathcal{D}_1\}$ .

## References

- [1] M. Borowiecki, I. Broere and P. Mihók, *Minimal reducible bounds for planar graphs* (submitted).

- [2] M. Borowiecki, I. Broere, M. Frick, P. Mihók and G. Semanišin, *A survey of hereditary properties of graphs*, *Discussiones Mathematicae Graph Theory* **17** (1997) 5–50.
- [3] M. Borowiecki and P. Mihók, *Hereditary Properties of Graphs*, in: *Advances in Graph Theory* (Vishwa Intern. Publications, 1991) 41–68.
- [4] P. Borowiecki, *P-Bipartitions of Graphs*, *Vishwa Intern. J. Graph Theory* **2** (1993) 109–116.
- [5] I. Broere and C.M. Mynhardt, *Generalized colourings of outerplanar and planar graphs*, in: *Graph Theory with Applications to Algorithms and Computer Science* (Wiley, New York, 1985) 151–161.
- [6] G. Chartrand and L. Lesniak, *Graphs and Digraphs* (Second Edition, Wadsworth & Brooks/Cole, Monterey, 1986).
- [7] G. Dirac, *A property of 4-chromatic graphs and remarks on critical graphs*, *J. London Math. Soc.* **27** (1952) 85–92.
- [8] W. Goddard, *Acyclic colorings of planar graphs*, *Discrete Math.* **91** (1991) 91–94.
- [9] T.R. Jensen and B. Toft, *Graph Colouring Problems* (Wiley-Interscience Publications, New York, 1995).
- [10] P. Mihók, *On the vertex partition numbers of graphs*, in: M. Fiedler, ed., *Graphs and Other Combinatorial Topics*, Proc. Third Czech. Symp. Graph Theory, Prague, 1982 (Teubner-Verlag, Leipzig, 1983) 183–188.
- [11] P. Mihók, *On the minimal reducible bound for outerplanar and planar graphs*, *Discrete Math.* **150** (1996) 431–435.
- [12] K.S. Poh, *On the Linear Vertex-Arboricity of a Planar Graph*, *J. Graph Theory* **14** (1990) 73–75.
- [13] J. Wang, *On point-linear arboricity of planar graphs*, *Discrete Math.* **72** (1988) 381–384.

Received 25 February 1997