

UNIQUELY PARTITIONABLE PLANAR GRAPHS  
WITH RESPECT TO PROPERTIES HAVING  
A FORBIDDEN TREE

JOZEF BUCKO

*Department of Mathematics, Technical University*  
*Hlavná 6, 040 01 Košice, Slovak Republic*

**e-mail:** bucko@ccsun.tuke.sk

AND

JAROSLAV IVANČO

*Department of Geometry and Algebra*  
*P.J. Šafárik University, Jesenná 5*  
*041 54 Košice, Slovak Republic*

**e-mail:** ivanco@duro.upjs.sk

**Abstract**

Let  $\mathcal{P}_1, \mathcal{P}_2$  be graph properties. A vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition of a graph  $G$  is a partition  $\{V_1, V_2\}$  of  $V(G)$  such that for  $i = 1, 2$  the induced subgraph  $G[V_i]$  has the property  $\mathcal{P}_i$ . A property  $\mathcal{R} = \mathcal{P}_1 \circ \mathcal{P}_2$  is defined to be the set of all graphs having a vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition. A graph  $G \in \mathcal{P}_1 \circ \mathcal{P}_2$  is said to be uniquely  $(\mathcal{P}_1, \mathcal{P}_2)$ -partitionable if  $G$  has exactly one vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition. In this note, we show the existence of uniquely partitionable planar graphs with respect to hereditary additive properties having a forbidden tree.

**Keywords:** uniquely partitionable planar graphs, forbidden graphs.

**1991 Mathematics Subject Classification:** 05C15, 05C70.

1. INTRODUCTION

Let us denote by  $\mathcal{I}$  the class of all finite undirected graphs without loops and multiple edges. If  $\mathcal{P}$  is a proper isomorphism closed subclass of  $\mathcal{I}$ , then  $\mathcal{P}$  will also denote the property that a graph is a member of the set  $\mathcal{P}$ . We shall use the terms *set of graphs* and *property of graphs* interchangeably.

A property  $\mathcal{P}$  is said to be *hereditary* if, whenever  $G \in \mathcal{P}$  and  $H$  is a subgraph of  $G$ , then also  $H \in \mathcal{P}$ . A property  $\mathcal{P}$  is called *additive* if for each graph  $G$  all of whose components have the property  $\mathcal{P}$  it follows that  $G \in \mathcal{P}$ , too.

For every hereditary property  $\mathcal{P}$  there is a nonnegative integer  $c(\mathcal{P})$  such that  $K_{c(\mathcal{P})+1} \in \mathcal{P}$  but  $K_{c(\mathcal{P})+2} \notin \mathcal{P}$  called the *completeness* of  $\mathcal{P}$ . For example  $c(\mathcal{O}) = 0$ ,  $c(\mathcal{D}_1) = 1$ ,  $c(\mathcal{T}_2) = 2$ ,  $c(\mathcal{T}_3) = 3$ , where  $\mathcal{O}$  is the class of all totally disconnected graphs,  $\mathcal{D}_1$  is the class of acyclic graphs,  $\mathcal{T}_2$  is the class of outerplanar graphs and  $\mathcal{T}_3$  is the class of planar graphs.

Any hereditary property  $\mathcal{P}$  is uniquely determined by the set

$$\mathbf{F}(\mathcal{P}) = \{G \in \mathcal{I} \mid G \notin \mathcal{P}, \text{ but each proper subgraph } H \text{ of } G \text{ belongs to } \mathcal{P}\}$$

of its minimal forbidden subgraphs.

Let  $\mathcal{P}_1, \mathcal{P}_2$  be arbitrary hereditary properties of graphs. A *vertex*  $(\mathcal{P}_1, \mathcal{P}_2)$ -*partition* of a graph  $G$  is a partition  $\{V_1, V_2\}$  of  $V(G)$  such that for  $i = 1, 2$  the induced subgraph  $G[V_i]$  has the property  $\mathcal{P}_i$ .

A property  $\mathcal{R} = \mathcal{P}_1 \circ \mathcal{P}_2$  is defined to be the set of all graphs having a vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition. It is easy to see that if  $\mathcal{P}_1, \mathcal{P}_2$  are additive and hereditary, then  $\mathcal{R} = \mathcal{P}_1 \circ \mathcal{P}_2$  is additive and hereditary, too.

A graph  $G \in \mathcal{P}_1 \circ \mathcal{P}_2$  is said to be *uniquely*  $(\mathcal{P}_1, \mathcal{P}_2)$ -*partitionable* if  $G$  has exactly one (unordered) vertex  $(\mathcal{P}_1, \mathcal{P}_2)$ -partition. For the concept of uniquely partitionable graphs we refer the reader to [1]. Basic properties of uniquely partitionable graphs are discussed in [1] and [4].

**Proposition 1** [1]. *Let  $\mathcal{P}$  be an additive hereditary property. Then there exists a uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable graph  $G$  if and only if  $\mathcal{P} \neq \mathcal{O} \circ \mathcal{Q}$ .*

The proof used non-planar graphs. The constructions of uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable outerplanar and planar graphs were presented in [2]. The following results have been proved:

**Proposition 2** [2]. *Let  $\mathcal{P}$  be an additive hereditary property of completeness 1. Then there exists a uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable outerplanar graph  $G$  if and only if there is a tree  $T$  which is forbidden for  $\mathcal{P}$ .*

**Proposition 3** [2]. *Let  $\mathcal{P}$  be an additive hereditary property of completeness 1. Then there exists a uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable planar graph  $G$  if and only if either some odd cycle  $C_{2q+1}$  has property  $\mathcal{P}$  or there is a bipartite planar graph  $H$  which is forbidden for  $\mathcal{P}$ .*

Our first result shows that the restriction on the completeness is not necessary for the existence of uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable planar graphs.

**Theorem 1.** *Let  $\mathcal{P}$  be an additive hereditary property. If there is a tree  $T \in \mathbf{F}(\mathcal{P})$ , then there exists a uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable planar graph.*

Furthermore, let us consider  $(\mathcal{D}_1, \mathcal{D}_1)$ -partitions of planar graphs. The following result is presented in [3]:

**Proposition 4** [3]. *There are no uniquely  $(\mathcal{D}_1, \mathcal{D}_1)$ -partitionable planar graph.*

In this note, we shall show that the property  $\mathcal{D}_1 \circ \mathcal{D}_1$  is in some sense "a minimal property" having no uniquely partitionable planar graphs. More precisely, we will prove the following result:

**Theorem 2.** *Let  $\mathcal{P}, \mathcal{Q}$  be the additive hereditary properties of graphs with completeness 1. If there is a tree  $T \in \mathbf{F}(\mathcal{P})$ , then there exists a uniquely  $(\mathcal{P}, \mathcal{Q})$ -partitionable planar graph.*

2. PROOFS OF THE MAIN RESULTS

**Proof of Theorem 1.** Let  $T$  be a forbidden tree for a property  $\mathcal{P}$ . As every connected bipartite planar graph is uniquely  $(\mathcal{O}, \mathcal{O})$ -partitionable, we can assume that  $T$  has at least 3 vertices. Then  $T$  contains a path  $wuv_1$ , where  $v_1$  is an end vertex of  $T$ . Denote by  $T'$  the graph which we obtain from  $T$  by adding the edge  $wv_1$ .  $T'$  is outerplanar and so the join  $K_1 + T'$  is a planar graph. Let  $G(T, 1)$  be the graph which we obtain from  $K_1 + T'$  by deleting the edge  $av_1$ , where  $a$  denotes the vertex of  $K_1$ . Evidently,  $G(T, 1)$  may be embedded on the plane such that the vertices  $a$  and  $v_1$  lie in the exterior face (see Figure 1).





Figure 1

$G(T, k)$ , for  $k > 1$ , is a planar graph which we obtain from  $G(T, 1)$  by adding the vertices  $v_2, v_3, \dots, v_k$  and edges  $uv_2, uv_3, \dots, uv_k, wv_2, wv_3, \dots, wv_k$ . The vertex  $a$  is called the *root* of  $G(T, k)$  and vertices  $v_1, v_2, \dots, v_k$ , are called *leaves* of  $G(T, k)$ . Moreover, for every leaf  $v_i$  we define its successor  $s(v_i)$  by  $s(v_1) = a$  and  $s(v_i) = v_{i-1}$ , if  $i = 2, 3, \dots, k$ . Obviously,  $G(T, k)$  may be embedded on the plane such that both vertices  $v_i$  and  $s(v_i)$  lie in a common face (see Figure 1).

Now we construct a planar graph  $H(T, k, d)$  using the induction on  $d$ .  $H(T, k, 1)$  is a graph which we obtain from  $k$  copies of  $G(T, k)$  by identifying their roots. The vertex arisen by the identification is called the *root* of  $H(T, k, 1)$ . The *leaves* of copies of  $G(T, k)$  are leaves of  $H(T, k, 1)$ . Similarly, the successor of a leaf in  $H(T, k, 1)$  is equal to the successor of this leaf in the corresponding copy of  $G(T, k)$ . For  $d > 1$ ,  $H(T, k, d)$  is a planar graph which we obtain from  $H(T, k, 1)$  and  $k^2$  copies of  $H(T, k, d-1)$  by identifying each leaf of  $H(T, k, 1)$  with the root of a copy of  $H(T, k, d-1)$ . Evidently, a copy of  $H(T, k, d-1)$  can be inserted into a face of  $H(T, k, 1)$  which contains a corresponding leaf  $x$  of  $H(T, k, 1)$  and its successor  $s_1(x)$  in  $H(T, k, 1)$  (see Figure 2).






Figure 2

The root of  $H(T, k, d)$  is the root of  $H(T, k, 1)$  and the leaves of  $H(T, k, d)$  are leaves of copies of  $H(T, k, d-1)$ . Denote by  $s_d(y)$  and  $s_{d-1}(y)$  the successor of a leaf  $y$  in  $H(T, k, d)$  and in a corresponding copy of  $H(T, k, d-1)$ . Then

$$s_d(y) = \begin{cases} s_1(x), & \text{if } s_{d-1}(y) \text{ was identified with } x, \\ s_{d-1}(y), & \text{otherwise.} \end{cases}$$

Finally,  $H^*(T, k, d)$  is a planar graph which we obtain from  $H(T, k, d)$  such that we connect each leaf of  $H(T, k, d)$  with its successor by a copy of  $G(T, 1)$  identifying the leaf with the root of  $G(T, 1)$  and the successor with the leaf of  $G(T, 1)$  (see Figure 2).

Put  $V_1 = \{x \in V(H^*(T, k, d)) \mid d(r, x) \equiv 0 \pmod{2}\}$ , where  $r$  denotes the root of  $H(T, k, d)$  and  $d(y, z)$  is the length of the shortest path between  $y$  and  $z$  in  $H(T, k, d)$ . The vertices belonging to  $V_1$  are depicted by white in Figure 2. It is easy to see that  $V_1$  is an independent set of  $H^*(T, k, d)$ . Moreover, the set  $V_2 = V(H^*(T, k, d)) - V_1$  induces a subgraph of  $H^*(T, k, d)$  each of whose components is isomorphic to  $T - v_1$ . So,  $\{V_1, V_2\}$  is a vertex  $(\mathcal{O}, \mathcal{P})$ -partition of  $H^*(T, k, d)$ .

Suppose that  $\{U_1, U_2\}$  is a vertex  $(\mathcal{O}, \mathcal{P})$ -partition of  $H^*(T, k, d)$ . Consider two cases:

*Case 1.*  $U_1 \cap V_1 \neq \emptyset$ . Let  $x \in U_1 \cap V_1$  and let  $y$  be any vertex of  $V_1 - \{x\}$ . From the construction of  $H^*(T, k, d)$  it can easily be seen that there exists

a sequence  $x = x_1, x_2, \dots, x_t = y$  satisfying: For every  $i = 1, \dots, t - 1$ , there is a subgraph  $G_i$  of  $H^*(T, k, d)$  isomorphic to  $G(T, k)$  (or  $G(T, 1)$ ), where  $x_i$  is its root and  $x_{i+1}$  is its leaf. As  $x_1$  belongs to  $U_1$ , all vertices of  $G_1$  adjacent to  $x_1$  belong to  $U_2$ . However, these neighbours of  $x_1$  together with  $x_2$  induce a subgraph of  $G_1$  containing  $T$ . Therefore,  $x_2 \in U_1$ , and by induction,  $y \in U_1$ . Since  $y$  is any vertex of  $V_1 - \{x\}$ ,  $V_1 \subseteq U_1$ . The set  $V_1$  is a domination set of  $H^*(T, k, d)$ , and so,  $V_1 = U_1$ , i.e.,  $\{U_1, U_2\} = \{V_1, V_2\}$ .

*Case 2.*  $V_1 \subseteq U_2$ . It is easy to see that every block of  $H(T, k, d)$  is a copy of  $G(T, k)$ , where the root and leaves of the copy belong to  $V_1$ . As the vertices of a block corresponding to  $u$  and  $w$  are adjacent, at least one of them belongs to  $U_2$ . Thus, vertices of a block belonging to  $U_2$  induce a graph containing a star  $K_{1, k+1}$ . From the construction of  $H(T, k, d)$  one can see that vertices of  $H(T, k, d)$  belonging to  $U_2$  induce a graph containing a complete  $k$ -ary tree with  $2d+1$  levels. Therefore, for  $k \geq \Delta(T)$  and  $d \geq \frac{1}{2}rad(T)$ ,  $H^*(T, k, d)[U_2]$  contains a subgraph isomorphic to  $T$ , a contradiction. Thus, for  $k \geq \Delta(T)$  and  $d \geq \frac{1}{2}rad(T)$ , the graph  $H^*(T, k, d)$  is uniquely  $(\mathcal{O}, \mathcal{P})$ -partitionable. ■

**Proof of Theorem 2.** To construct the planar graph  $H_r(s)$ , for  $r \geq 1$ ,  $s \geq 2$  we will use the induction on  $r$ . The first step is the construction of planar graph  $H_1(s)$ :

$H_1(s) = K_2 + \cup_{i=1}^s K_2$ , where  $V(K_2) = \{x_1, x_2\}$  and  $V(\cup_{i=1}^s K_2) = \{y_{1i}, y_{2i} \mid i = 1, 2, \dots, s\}$ . The edge  $x_1x_2$  of  $H_1(s)$  we will call the "major" edge of  $H_1(s)$  and edges  $y_{1i}y_{2i}$ ,  $i = 1, 2, \dots, s$  we will call "minor" edges of  $H_1(s)$ . For the construction of  $H_1(3)$  see Figure 3.




Figure 3. The graph  $H_1(3)$

Let us construct the graph  $H_{k+1}(s)$  in the following way:

We insert  $s$  copies of graph  $H_k(s)$  to graph  $H_1(s)$  such that we identify the "major" edges of copies of graphs  $H_k(s)$  with "minor" edges of  $H_1(s)$ . For the construction of  $H_2(3)$  see Figure 4.

It is easy to see from the construction, that  $H_r(s)$  is a planar graph. Now we shall show, that if the maximum degree  $\Delta(T)$  of the tree  $T \in \mathbf{F}(\mathcal{P})$  is  $\Delta(T) \leq s$  and radius  $rad(T)$  of the tree  $T$  is  $rad(T) \leq r$ , then the planar graph  $H_r(s)$  is uniquely  $(\mathcal{P}, \mathcal{Q})$ -partitionable.

Let us distinguish two "possible" vertex partitions of the graph  $H_r(s)$ :  
 1. The end vertices  $x_1, x_2$  of "major" edge of  $H_1(s)$  belong to different classes of the vertex partition. From the fact that  $K_3$  is forbidden for both properties  $\mathcal{P}, \mathcal{Q}$ , it follows that vertices of "minor" edges of  $H_1(s)$  belong to different classes of the vertex partition, too. By induction on  $r$  in both classes of the partition, it grove the complete  $s$ -ary tree with  $1 + r$  levels, which is, for  $r \geq rad(T)$  and  $s \geq \Delta(T)$ , a supergraph of the forbidden tree  $T$ . It means, that it is not a  $(\mathcal{P}, \mathcal{Q})$ -partition of  $H_r(s)$ .




Figure 4. The graph  $H_2(3)$

2. Hence the end vertices  $x_1, x_2$  of "major" edge of the graph  $H_1(s)$  have to belong to the same class of a vertex partition. From the fact that  $K_3$  is forbidden for both properties  $\mathcal{P}, \mathcal{Q}$ , it follows, that vertices of "minor" edges of  $H_1(s)$  have both to belong to the second class of the vertex partition. From

the construction of  $H_r(s)$  and from the fact that  $K_3$  is forbidden it is easy to see that the partition of  $H_r(s)$  is a  $(\mathcal{P}, \mathcal{Q})$ -partition of  $H_r(s)$ . Thus  $H_r(s)$ , for  $r \geq \text{rad}(T)$  and  $s \geq \Delta(T)$  is a uniquely  $(\mathcal{P}, \mathcal{Q})$ -partitionable graph. ■

## REFERENCES

- [1] J. Bucko, M. Frick, P. Mihók and R. Vasky, *Uniquely partitionable graphs*, Discuss. Math. Graph Theory **17** (1997) 103–114.
- [2] J. Bucko, P. Mihók and M. Voigt, *Uniquely partitionable planar graphs*, Discrete Math. **191** (1998) 149–158.
- [3] M. Borowiecki, J. Bucko, P. Mihók, Z. Tuza and M. Voigt, *Remarks on the existence of uniquely partitionable planar graphs*, 13. Workshop on Discrete Optimization, Burg, abstract, 1998.
- [4] P. Mihók, *Additive hereditary properties and uniquely partitionable graphs*, in: M. Borowiecki and Z. Skupień, eds., Graphs, hypergraphs and matroids (Zielona Góra, 1985) 49–58.

Received 14 July 1998

Revised 24 November 1998