

## ISOMORPHIC COMPONENTS OF KRONECKER PRODUCT OF BIPARTITE GRAPHS

PRANAVA K. JHA

*Department of Computer Engineering*  
*Delhi Institute of Technology: Delhi Kashmere Gate*  
*Delhi 110 006, India*  
**e-mail:** pkj@dit.ernet.in

SANDI KLAVŽAR\*

AND

BLAŽ ZMAZEK

*Department of Mathematics, PEF, University of Maribor*  
*Koroška cesta 160, 2000 Maribor, Slovenia*  
**e-mail:** sandi.klavzar@uni-lj.si

### Abstract

Weichsel (Proc. Amer. Math. Soc. 13 (1962) 47–52) proved that the Kronecker product of two connected bipartite graphs consists of two connected components. A condition on the factor graphs is presented which ensures that such components are isomorphic. It is demonstrated that several familiar and easily constructible graphs are amenable to that condition. A partial converse is proved for the above condition and it is conjectured that the converse is true in general.

**Keywords:** Kronecker product, bipartite graphs, graph isomorphism.

**1991 Mathematics Subject Classification:** 05C60.

### 1. INTRODUCTION

The Kronecker product is one of the (four) most important graph products, and can be viewed as the product in the category of graphs, cf. Hell [4]. The product was used by Greenwell and Lovász [2] to demonstrate that for all  $n \geq 3$ , there is a uniquely  $n$ -colorable graph without odd cycles shorter than a given number  $s$ . Every graph is an induced subgraph of a Kronecker

---

\*This work was supported in part by the Ministry of Science and Technology of Slovenia under the grants P1-0206-101 and J1-7036.

product of certain complete graphs, see Nešetřil [11]. Graph retracts, see for instance Pesch [12], form another area where the Kronecker product construction turned out to be useful. The product has several other applications, for instance in modelling concurrency in multiprocessor systems [10] and in automata theory. (For example, closure of regular sets, or closure of context-free sets with a regular set, under intersection may be proved by taking a Kronecker product of the respective machines.)

Weichsel [14] proved that the *Kronecker product* of two nontrivial graphs is connected if and only if both factors are connected and at least one of them possesses an odd cycle. If both factors are connected and bipartite, then their Kronecker product consists of two connected components. When are these components isomorphic? This natural question plays the central role in our paper.

The *Kronecker product*  $G \times H$  of graphs  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$  and edge set  $\{(u, x), (v, y) \mid \{u, v\} \in E(G) \text{ and } \{x, y\} \in E(H)\}$ . The terminology is justified by the fact that the adjacency matrix of a Kronecker graph product is given by the Kronecker matrix product of the adjacency matrices of the factor graphs; see [14] for details. However, this product is also known under several different names including *categorical product*, *tensor product*, *direct product*, *weak direct product*, *cardinal product* and *graph conjunction*.

The Kronecker product is commutative and associative in an obvious way. It is also distributive with respect to edge-disjoint union of graphs. In particular, if  $G$  and  $H$  are bipartite graphs which respectively appear as subgraphs of (not necessarily bipartite) graphs  $G_1$  and  $H_1$ , then the two components of  $G \times H$  appear as vertex-disjoint subgraphs of  $G_1 \times H_1$ . This lends some structure to the graph  $G_1 \times H_1$ . In fact, this view has been effectively used in Hamiltonian decompositions of certain Kronecker-product graphs [8].

The rest of the paper is organized as follows. In the next section, we give some more definitions, recall Weichsel's result and give an alternative proof of it. In Section 3, we prove that a certain condition on factor graphs is sufficient for the existence of an isomorphism between components of the Kronecker product of two bipartite graphs. It is also shown that several familiar and easily constructible graphs are amenable to that condition. A partial converse to the sufficient condition is established in the last section and it is conjectured that it holds in general. We conclude the paper with some remarks on the algorithmic aspects of related problems.

## 2. PRELIMINARIES

By a graph is meant a finite, simple, undirected graph having at least two vertices. Unless indicated otherwise, graphs are also connected.

Another graph product which is of relevance here is the Cartesian product. The *Cartesian product*  $G \square H$  of graphs  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$  and  $\{(u, x), (v, y)\} \in E(G \square H)$  whenever  $\{u, v\} \in E(G)$  and  $x = y$ , or  $u = v$  and  $\{x, y\} \in E(H)$ . Note that the Cartesian product of two (or finitely many) graphs is connected if and only if every factor is. Observe also that  $G \square H$  is bipartite if and only if both  $G$  and  $H$  are bipartite [13] and that  $G \times H$  is bipartite if and only if at least one of  $G$  and  $H$  is bipartite [3].

We have already mentioned the following result of Weichsel:

**Theorem 2.1** [14]. *Let  $G$  and  $H$  be connected graphs. Then  $G \times H$  is connected if and only if at least one of  $G$  and  $H$  contains an odd cycle.* ■

The next result refines a part of Theorem 2.1.

**Lemma 2.2** [7]. *If  $G = (V_0 \cup V_1, E)$  and  $H = (W_0 \cup W_1, F)$  are bipartite graphs, then  $(V_0 \times W_0) \cup (V_1 \times W_1)$  and  $(V_0 \times W_1) \cup (V_1 \times W_0)$  are vertex sets of the two components of  $G \times H$ .* ■

While Lemma 2.2 is an alternative proof of the “only if” part of Theorem 2.1, the following argument (which is different from Weichsel’s) takes care of the converse. Recall that  $\times$ -product is distributive with respect to the edge-disjoint union of graphs.

Let  $G = (V, E)$  be a non-bipartite graph, and let  $H = (W, F)$  be any graph. Choose spanning subgraphs  $G'$  and  $H'$  of  $G$  and  $H$ , respectively, and an edge  $e = \{a, b\}$  of  $G \setminus G'$  such that

- (i)  $G'$  and  $H'$  are connected, bipartite graphs and
- (ii)  $G' + e$  is a non-bipartite graph.

$G'$  and  $H'$  can always be constructed by successively removing edges from odd cycles. Let  $V_0$  and  $V_1$  (resp.  $W_0$  and  $W_1$ ) be the partite sets of  $G'$  (resp.  $H'$ ). Note that  $a, b \in V_0$  or  $a, b \in V_1$ . It suffices to show that  $(G' + e) \times H'$  is a connected graph.

Without loss of generality, let  $a, b \in V_0$ . By Lemma 2.2,  $G' \times H'$  consists of two connected components having vertex sets  $(V_0 \times W_0) \cup (V_1 \times W_1)$  and  $(V_0 \times W_1) \cup (V_1 \times W_0)$ . Let  $\{c, d\} \in E(H')$ , where  $c \in W_0$  and  $d \in W_1$ .

It is clear that  $(a, c) \in V_0 \times W_0$ ,  $(b, d) \in V_0 \times W_1$ , and  $\{(a, c), (b, d)\}$  is an edge of  $(G' + e) \times H'$  which runs between  $(V_0 \times W_0) \cup (V_1 \times W_1)$  and  $(V_0 \times W_1) \cup (V_1 \times W_0)$ . It immediately follows that the graph  $(G' + e) \times H'$  is connected, and hence the result.

### 3. PROPERTY $\pi$

This is our main definition.

**Definition 3.1.** *A bipartite graph  $G = (V_0 \cup V_1, E)$  is said to have a property  $\pi$  if  $G$  admits an automorphism  $\varphi$  such that  $x \in V_0$  if and only if  $\varphi(x) \in V_1$ . ■*

Note that a bipartite graph which satisfies the foregoing definition possesses a certain symmetry, and has the same number of vertices in each partite set. Examples of graphs possessing the property  $\pi$  include even paths, even cycles, complete bipartite graphs with equal-sized partite sets, and the graph obtained from two copies of a star  $K_{1,n}$  by introducing an edge between the center vertices of the two stars.

**Theorem 3.2.** *If  $G$  and  $H$  are bipartite graphs one of which has the property  $\pi$ , then the two components of  $G \times H$  are isomorphic.*

**Proof.** Let  $G = (V_0 \cup V_1, E)$  and  $H = (W_0 \cup W_1, F)$  be bipartite graphs. Assume that the graph  $G$  possesses the property  $\pi$  with  $\varphi$  as an appropriate automorphism. The sets  $(V_0 \times W_0) \cup (V_1 \times W_1)$  and  $(V_0 \times W_1) \cup (V_1 \times W_0)$  correspond to vertex sets of the two components of the graph  $G \times H$ . Consider the mapping given by  $(u, x) \mapsto (\varphi(u), x)$  between vertex sets of these components. It is straightforward to see that this is a well-defined bijection which corresponds to a desired isomorphism. ■

We next show that graphs with the property  $\pi$  enjoy certain interesting closure properties with respect to the Kronecker product and the Cartesian product.

**Proposition 3.3.** *If  $G$  and  $H$  are bipartite graphs having the property  $\pi$ , then so is each of the following:*

- (i)  $\times$ -product of  $G$  and a non-bipartite graph,
- (ii) each component of the  $\times$ -product of  $G$  and  $H$ , and
- (iii)  $\square$ -product of  $G$  and a bipartite graph.

**Proof.** Let  $G = (V_0 \cup V_1, E)$  and  $H = (W_0 \cup W_1, F)$  be bipartite graphs which are respectively equipped with automorphisms  $\varphi$  and  $\psi$  in the sense of Definition 3.1.

(i) Let  $K = (X, D)$  be a non-bipartite graph. The two partite sets of the connected, bipartite graph  $G \times K$  are  $V_0 \times X$  and  $V_1 \times X$ . The mapping given by  $(v, x) \mapsto (\varphi(v), x)$  constitutes an automorphism of  $G \times K$  corresponding to the property  $\pi$ .

(ii) Consider the component of the (bipartite) graph  $G \times H$  on vertex set  $(V_0 \times W_0) \cup (V_1 \times W_1)$ . Note that  $V_0 \times W_0$  and  $V_1 \times W_1$  are the two partite sets of this component. The mapping given by  $(v, w) \mapsto (\varphi(v), \psi(w))$  constitutes a desired automorphism of this component.

(iii) Let  $K = (X_0 \cup X_1, D)$  be a bipartite graph. The two partite sets of the (connected) bipartite graph  $G \square K$  are  $(V_0 \times X_0) \cup (V_1 \times X_1)$  and  $(V_0 \times X_1) \cup (V_1 \times X_0)$ . The mapping given by  $(v, x) \mapsto (\varphi(v), x)$  constitutes a desired automorphism of this graph. ■

It follows from Proposition 3.3 that each of the following graphs has the property  $\pi$ : (i) the  $n$ -cube  $Q_n$  ( $Q_1 = K_2$ ,  $Q_n = K_2 \square Q_{n-1}$ ,  $n \geq 2$ ), (ii) the planar grid  $P_m \square P_n$ , where  $m$  or  $n$  is even, (iii) the graph  $C_{n_1} \square \dots \square C_{n_r}$ , where each  $n_i$  is even, and (iv) each component of the graph  $C_{n_1} \times \dots \times C_{n_r}$ , where at least one  $n_i$  is even.

The following is a relevant result. It is a generalization of a similar result from [6] for the graph  $C_m \times P_n$ .

**Proposition 3.4.** *If  $m$  is even,  $m/2$  is odd and  $G$  is a bipartite graph, then each component of the graph  $C_m \times G$  is isomorphic to  $C_{m/2} \times G$ .*

**Proof.** Let  $m$  and  $G = (V_0 \cup V_1, E)$  be as stated. Denote the consecutive vertices of  $C_m$  by  $0, 1, \dots, m-1$ . Then vertex sets of the two components of the graph  $C_m \times G$  are

$$(\{0, 2, \dots, m-2\} \times V_0) \cup (\{1, 3, \dots, m-1\} \times V_1)$$

and

$$(\{0, 2, \dots, m-2\} \times V_1) \cup (\{1, 3, \dots, m-1\} \times V_0).$$

By Theorem 3.2, these two components are isomorphic. Consider the mapping from the vertex set of one of these components to that of  $C_{m/2} \times G$  given by  $(i, x) \mapsto (i \bmod (m/2), x)$ . That this is a well-defined bijection corresponding to a desired isomorphism is left to the reader. ■

4. A CONJECTURE

We conjecture that the converse of Theorem 3.2 holds as well:

**Conjecture 4.1.** *Let  $G$  and  $H$  be bipartite graphs. Then the components of  $G \times H$  are isomorphic if and only if at least one of  $G$  and  $H$  has the property  $\pi$ .*

In the rest of the paper, let  $G = (V_0 \cup V_1, E)$  and  $H = (W_0 \cup W_1, F)$  be bipartite graphs and let  $X$  and  $Y$  be the two connected components of  $G \times H$  induced by  $(V_0 \times W_0) \cup (V_1 \times W_1)$  and  $(V_0 \times W_1) \cup (V_1 \times W_0)$ , respectively.

Note first that if  $X$  and  $Y$  are isomorphic, then at least one of  $|V_0| = |V_1|$  or  $|W_0| = |W_1|$  holds. Indeed, since  $|X| = |Y|$  we have  $|V_0||W_0| + |V_1||W_1| = |V_0||W_1| + |V_1||W_0|$ . Therefore  $|V_0|(|W_0| - |W_1|) = |V_1|(|W_0| - |W_1|)$  and thus the observation. However, much more is true, but first we need some more notation.

As usual, for  $x \in V(G)$ , let  $N_G(x) = \{y \mid \{x, y\} \in E(G)\}$  and for  $Q \subseteq V(G)$ , let  $N_G(Q) = \bigcup_{x \in Q} N_G(x)$ . We will also write  $N(Q)$  if the graph  $G$  will be clear from the context. For a graph  $G$  and  $Q \subseteq V(G)$  let  $\deg_G(Q)$  be the multiset  $\{\deg_G(v) \mid v \in Q\}$ . Also, for a set  $Q \subseteq V(G)$  let  $N^i(Q)$ ,  $i \geq 1$ , be defined by  $N^1(Q) = N(Q)$  and  $N^i(Q) = N(N^{i-1}(Q))$ ,  $i \geq 2$ .

**Theorem 4.2.** *Let  $\varphi : X \rightarrow Y$  be an isomorphism which maps  $V'_0 \times W'_0$  onto  $V'_1 \times W'_0$ , where  $V'_0 \subseteq V_0$ ,  $V'_1 \subseteq V_1$  and  $W'_0 \subseteq W_0$ . Then for all  $i \geq 1$ ,  $\deg_G(N^i(V'_0)) = \deg_G(N^i(V'_1))$  hold.*

**Proof.** Since  $\varphi$  is an isomorphism, we have

$$\varphi(N_{G \times H}(V'_0 \times W'_0)) = N_{G \times H}(\varphi(V'_0 \times W'_0)) = N_{G \times H}(V'_1 \times W'_0)$$

and from the definition of the Kronecker product it follows

$$N_{G \times H}(V'_1 \times W'_0) = N_G(V'_1) \times N_H(W'_0).$$

Therefore,  $\varphi$  maps  $N_{G \times H}(V'_0 \times W'_0) = N_G(V'_0) \times N_H(W'_0)$  onto  $N_G(V'_1) \times N_H(W'_0)$ .

Let  $v_0, v_1$  and  $w$  be vertices of maximum degrees in  $N_G(V'_0)$ ,  $N_G(V'_1)$  and  $N_H(W'_0)$ , respectively. Then  $(v_0, w) \in N_G(V'_0) \times N_H(W'_0)$  and  $(v_1, w) \in N_G(V'_1) \times N_H(W'_0)$  have the same (largest) degree and thus  $v_0$  and  $v_1$  must

have the same degree in  $G$ . It follows that  $\deg_{G \times H}(\{v_0\} \times N_H(W'_0)) = \deg_{G \times H}(\{v_1\} \times N_H(W'_0))$ , and therefore

$$\deg_{G \times H}((N(V'_0) \setminus \{v_0\}) \times N(W'_0)) = \deg_{G \times H}((N(V'_1) \setminus \{v_1\}) \times N(W'_0)).$$

Repeating the above argument, we conclude  $\deg_G(N(V'_0)) = \deg_G(N(V'_1))$ . This proves the theorem for  $i = 1$ .

We have seen above that  $\varphi$  maps  $N_G(V'_0) \times N_H(W'_0)$  isomorphically onto  $N_G(V'_1) \times N_H(W'_0)$ . But this is just the theorem assumption, hence as above we obtain  $\deg_G(N^2(V'_0)) = \deg_G(N^2(V'_1))$ . Repeating the argument, we prove the equality for any  $i \geq 1$ . ■

**Corollary 4.3.** *Let  $A_0, A_1, B_0$  and  $B_1$  be the sets of vertices with maximum (minimum) degree in  $V_0, V_1, W_0$  and  $W_1$ , respectively. Then if  $X$  and  $Y$  are isomorphic, at least one of  $\deg_G(N^i(A_0)) = \deg_G(N^i(A_1))$  or  $\deg_G(N^i(B_0)) = \deg_G(N^i(B_1))$  hold for all  $i \geq 1$ .*

**Proof.** Let  $\varphi : X \rightarrow Y$  be an isomorphism. As  $X$  and  $Y$  are connected bipartite graphs,  $\varphi$  maps  $V_0 \times W_0$  either onto  $V_1 \times W_0$  or onto  $V_0 \times W_1$ . In the former case  $\varphi(A_0 \times B_0) = A_1 \times B_0$  and by Theorem 4.2,  $\deg_G(N^i(A_0)) = \deg_G(N^i(A_1))$ . In the later case  $\varphi(A_0 \times B_0) = A_0 \times B_1$  and by Theorem 4.2 and commutativity of the Kronecker product,  $\deg_G(N^i(B_0)) = \deg_G(N^i(B_1))$ . ■

Consider for example the graph  $G$  in Figure 1.  $G$  fulfils the conclusion of Corollary 4.3. On the other hand, the connected components of  $G \times P_3$  are not isomorphic. Indeed, there is a vertex of degree 6 which is adjacent to vertices of degrees 2, 2, 3, 3, 4 and 4 in one of the components and there is no such vertex in the other component.




Figure 1. The graph  $G$

Theorem 4.2 also implies that if  $X$  and  $Y$  are isomorphic then in at least one of the factors the partite sets have the same degree sequences.

**Corollary 4.4.** *If  $X$  and  $Y$  are isomorphic, then at least one of  $\deg_G(V_0) = \deg_G(V_1)$  or  $\deg_H(W_0) = \deg_H(W_1)$  holds.*

**Proof.** Let  $\varphi : X \rightarrow Y$  be an isomorphism. As in Corollary 4.3  $\varphi$  maps  $V_0 \times W_0$  either onto  $V_1 \times W_0$  or onto  $V_0 \times W_1$ . Assume the former case and let  $A_0$  and  $A_1$  be as in Corollary 4.3. Clearly, for any  $i \geq 1$ ,  $N^i(A_0) \subseteq N^{i+2}(A_0)$  and  $N^i(A_1) \subseteq N^{i+2}(A_1)$ . Furthermore, since  $G$  is connected, there must be some  $i$  such that  $N^i(A_0) = V_0$  and  $N^i(A_1) = V_1$ . By Corollary 4.3,  $\deg_G(V_0) = \deg_G(V_1)$ .

In the case when  $\varphi$  maps  $V_0 \times W_0$  onto  $V_0 \times W_1$ , we analogously obtain  $\deg_H(W_0) = \deg_H(W_1)$ . ■

#### CONCLUDING ALGORITHMIC REMARKS

The Kronecker product seems to be interesting also from the algorithmic point of view. As we learned from Babai (at the Workshop on Cayley Graphs held in September 1996 in Montréal), Lalonde [9] proved that it is NP-complete to decide whether a given bipartite graph  $G$  has the property  $\pi$  provided in addition that the corresponding automorphism is of order 2. However, the problem of deciding whether  $G$  has the property  $\pi$  becomes isomorphism complete, cf. [1, page 1518].

Finally, we mention a recent very interesting algorithmic result of Imrich [5]. He namely proved that the unique prime factor decomposition of finite, nonbipartite, connected graphs with or without loops can be determined in polynomial time.

## References

- [1] L. Babai, *Automorphism Groups, Isomorphism, Reconstruction*, Chapter 27 in *Handbook of Combinatorics* (R.L. Graham, M. Grötschel, L. Lovász, eds.) Elsevier, Amsterdam, (1995) 1447–1540.
- [2] D. Greenwell and L. Lovász, *Applications of product colouring*, Acta Math. Acad. Sci. Hungar. **25** (1974) 335–340.
- [3] S. Hedetniemi, *Homomorphisms of graphs and automata*, University of Michigan, Technical Report 03105-44-T, (1966).
- [4] P. Hell, *An introduction to the category of graphs*, Ann. New York Acad. Sci. **328** (1979) 120–136.
- [5] W. Imrich, *Factorizing cardinal product graphs in polynomial time*, Discrete Math., to appear.

- [6] P.K. Jha, *Decompositions of the Kronecker product of a cycle and a path into long cycles and long paths*, Indian J. Pure Appl. Math. **23** (1992) 585–602.
- [7] P.K. Jha, *Hamiltonian decompositions of products of cycles*, Indian J. Pure Appl. Math. **23** (1992) 723–729.
- [8] P.K. Jha, N. Agnihotri and R. Kumar, *Edge exchanges in Hamiltonian decompositions of Kronecker-product graphs*, Comput. Math. Applic. **31** (1996) 11–19.
- [9] F. Lalonde, *Le probleme d'etoiles pour graphes est NP-complet*, Discrete Math. **33** (1981) 271–280.
- [10] R.H. Lamprey and B.H. Barnes, *Product graphs and their applications, Modelling and Simulation*, **5** (1974) 1119–1123 (Proc Fifth Annual Pittsburgh Conference, Instrument Society of America, Pittsburgh, PA, 1974).
- [11] J. Nešetřil, *Representations of graphs by means of products and their complexity*, Lecture Notes in Comput. Sci. **118** (1981) 94–102.
- [12] E. Pesch, *Minimal extensions of graphs to absolute retracts*, J. Graph Theory **11** (1987) 585–598.
- [13] V.G. Vizing, *The Cartesian product of graphs*, Vyc. Sis. **9** (1963) 30–43.
- [14] P.M. Weichsel, *The Kronecker product of graphs*, Proc. Amer. Math. Soc. **13** (1962) 47–52.

Received 18 February 1997

Revised 12 May 1997