

A NOTE ON STRONG AND CO-STRONG PERFECTNESS OF THE X -JOIN OF GRAPHS

ALINA SZELECKA

Institute of Mathematics, Technical University of Zielona Góra
Podgórna 50, 65–246 Zielona Góra, Poland

e-mail: A.Szelecka@im.pz.zgora.pl

AND

ANDRZEJ WŁOCH

Department of Mathematics, Technical University of Rzeszów
W. Pola 2, 35–359 Rzeszów, Poland

e-mail: awloch@ewa.prz.rzeszow.pl

Abstract

Strongly perfect graphs were introduced by C. Berge and P. Duchet in [1]. In [4], [3] the following was studied: the problem of strong perfectness for the Cartesian product, the tensor product, the symmetrical difference of n , $n \geq 2$, graphs and for the generalized Cartesian product of graphs. Co-strong perfectness was first studied by G. Ravindra and D. Basavayya [5]. In this paper we discuss strong perfectness and co-strong perfectness for the generalized composition (the lexicographic product) of graphs named as the X -join of graphs.

Keywords: strongly perfect graphs, co-strongly perfect graphs, the X -join of graphs.

1991 Mathematics Subject Classification: 05C75, 05C60.

1. INTRODUCTION

Let G be a finite undirected connected simple graph. By $V(G)$ and $E(G)$ we denote its vertex set and edge set, respectively. The notation $H = \langle V_0 \rangle_G$, $V_0 \subseteq V(G)$ means that H is the subgraph of G induced by V_0 . A subset $S \subset V(G)$ is said to be *stable* in G if no two distinct vertices of S are adjacent in G . A subset $Q \subseteq V(G)$ is a *clique* of G if $\langle Q \rangle_G$ is a complete subgraph of G . If the stable set S meets every maximal (with respect to the set inclusion) clique Q , then we will call it a *stable*

transversal of G . A graph G is called *strongly perfect* ([1]) if its every induced subgraph (including G itself) has a stable transversal. We call G *co-strongly perfect* ([5]) if G and the complementary graph \overline{G} to G are strongly perfect. Let G_1, \dots, G_n , $n \geq 2$, be graphs of the same order $m \geq 2$ with the vertex sets $V(G_i) = V = \{y_1, \dots, y_m\}$ for $i = 1, \dots, n$ and X be a graph such that $V(X) = \{x_1, \dots, x_n\}$. The X -join ([2]) of the sequence of graphs G_1, \dots, G_n and the graph X is the graph $X[G_1, \dots, G_n]$ with the vertex set $V(X) \times V$ and the edge set $\{[(x_j, y_p), (x_k, y_q)] : j = k \text{ and } [y_p, y_q] \in E(G_i) \text{ or } [x_j, x_k] \in E(X)\}$.

Observe that if $G_1 = G_2 = \dots = G_n = Y$, then we obtain the *composition* (the *lexicographic product*) of graphs Y and X denoted by $X[Y]$.

Let $V_0 \subseteq V(X) \times V$. By the *projection* $Pr_X V_0$ of the subset V_0 on the graph X we mean the set $Pr_X V_0 = \{x \in V(X) : \text{there exists } y \in V(G_i), 1 \leq i \leq n, \text{ that } (x, y) \in V_0\}$.

2. RESULTS

Put $G = X[G_1, \dots, G_n]$, for convenience. Let H be a connected induced subgraph of G such that it is not isomorphic to any induced subgraph H' of the graph X or G_i , for $i = 1, \dots, n$.

Let $Pr_X V(H) = \{x_{i1}, \dots, x_{ip}\}$, $2 \leq p \leq n$.

We partition the set $V(H)$ on p -disjoint sets $V_{ij}(H)$ such that $Pr_X V_{ij}(H) = \{x_{ij}\}$ for $j = 1, \dots, p$. For an arbitrary subset $R \subseteq V(H)$, in a natural way we can write $R = \bigcup_{j=1}^t R \cap V_{ij}(H)$, where $1 \leq t \leq p$.

For G and H given above it follows immediately.

Lemma 1. *If Q is a maximal clique of H , then $Pr_X Q$ is a maximal clique of $\langle Pr_X V(H) \rangle$.*

Lemma 2. *A subset $Q \subseteq V(H)$ is a maximal clique of H if and only if*

- (1) $Q \cap V_{ij}(H)$ is a maximal clique of $\langle V_{ij}(H) \rangle$ for $j = 1, \dots, p$ or $Q \cap V_{ij}(H) = \emptyset$ and
- (2) $Pr_X Q$ is a maximal clique of $\langle Pr_X V(H) \rangle$.

Proof. I. Assume that Q is a maximal clique of H . We can write $Q = \bigcup_{j=1}^t Q \cap V_{ij}(H)$ where $1 \leq t \leq p$ with $Q \cap V_{ij}(H) \neq \emptyset$ for each $j = 1, \dots, t$. Moreover, each of the sets $Q \cap V_{ij}(H)$ must be a clique of $\langle V_{ij}(H) \rangle$. Suppose there exists j , $1 \leq j \leq t$ such that $Q \cap V_{ij}(H)$, is not maximal. In consequence, there exists a vertex $(x_{ij}, y_r) \in V_{ij}(H) \setminus Q \cap V_{ij}(H)$, $1 \leq j \leq t$

(of course $(x_{ij}, y_r) \notin Q$) which is adjacent to each vertex from $Q \cap V_{ij}(H)$. Moreover, by the definition of G and from the fact that $Q \cap V_{ij}(H) \subset Q$ it follows that (x_{ij}, y_r) must be adjacent to each vertex from $Q \setminus Q \cap V_{ij}(H)$. In consequence, (x_{ij}, y_r) is adjacent to all vertices from Q and $(x_{ij}, y_r) \notin Q$, a contradiction with the assumption that Q is a maximal clique of H . This shows that the condition in (1) holds.

Condition (2) follows from Lemma 1.

II. Suppose that conditions (1) and (2) hold. We can write $Q = \bigcup_{j=1}^t Q \cap V_{ij}(H)$, $1 \leq t \leq p$. Note that $|Q| > 1$, by the assumption about H . Firstly, we shall show that Q is a clique of H . Let $(x_{ij}, y_r), (x_{ik}, y_s)$ be two distinct vertices from Q . If $j = k$, then they belong to $Q \cap V_{ij}(H)$ and are adjacent by (1). If $j \neq k$, then $x_{ij}, x_{ik} \in Pr_X Q$ and by (2) they are adjacent in X . Thus, by the definition of G the vertices $(x_{ij}, y_r), (x_{ik}, y_s)$ are adjacent in G . This proves that Q is a clique of H .

Assume that Q is not maximal. This means that there exists $(x_{il}, y_r) \notin Q$ but it is adjacent to each vertex from Q . Moreover, by the definition of G , the vertex x_{il} is adjacent to all vertices from $Pr_X Q$. This implies that $x_{il} \in Pr_X Q$ by (2). In consequence, it must be that $(x_{il}, y_r) \in V_{il}(H) \setminus Q \cap V_{il}(H)$ (evidently $(x_{il}, y_r) \notin Q \cap V_{il}(H)$). Since $Q \cap V_{il}(H) \subset Q$ and (x_{il}, y_r) is adjacent to each vertex from Q , then it is adjacent to each vertex from $Q \cap V_{il}(H)$. Hence by (1) it must be that $(x_{il}, y_r) \in Q \cap V_{il}(H)$, a contradiction. Hence, Q is a maximal clique of H and this completes the proof of the lemma. ■

Using the same method as in the proof of Lemma 2 we prove.

Lemma 3. *A subset $S \subset V(H)$ is a maximal stable set of H if and only if*

- (1) *$S \cap V_{ij}(H)$ is a maximal stable set of $\langle V_{ij}(H) \rangle$ for $j = 1, \dots, s$ or $S \cap V_{ij}(H) = \emptyset$ and*
- (2) *$Pr_X S$ is a maximal stable set of $\langle Pr_X V(H) \rangle$.*

Lemma 4 follows directly from the definition of the graph $X[G_1, \dots, G_n]$.

Lemma 4. $\overline{X[G_1, \dots, G_n]} = \overline{X[G_1, \dots, G_n]}$.

Theorem 1. *$X[G_1, \dots, G_n]$ is strongly perfect if and only if the graphs X, G_1, \dots, G_n are strongly perfect.*

Proof. I. Let $X[G_1, \dots, G_n]$ be strongly perfect. Then X, G_1, \dots, G_n are strongly perfect since they are isomorphic to some induced subgraphs of G .

II. Suppose that the graphs X, G_1, \dots, G_n are strongly perfect. We shall show that G is strongly perfect. Let H be a connected induced subgraph of G . We shall prove that H has a stable transversal.

If H is an induced subgraph of X or $G_i, 1 \leq i \leq n$, then H has a stable transversal, by the assumption that X, G_1, \dots, G_n are strongly perfect.

Let H be not induced subgraph of $X, G_i, i = 1, \dots, n$. Assume that H does not have a stable transversal, i.e., for every maximal stable set $S \subseteq V(H)$ there exists a maximal clique $Q \subseteq V(H)$ such that $S \cap Q = \emptyset$. Moreover, by the definition of G and Lemmas 2, 3 we have that for every maximal stable set $Pr_X S$ of $\langle Pr_X V(H) \rangle$ there exists a maximal clique $Pr_X Q$ of $\langle Pr_X V(H) \rangle$ such that $Pr_X S \cap Pr_X Q = \emptyset$. This is a contradiction, since $\langle Pr_X V(H) \rangle$ has a stable transversal.

This proves that $X[G_1, \dots, G_n]$ is strongly perfect and the proof is complete. \blacksquare

For $G_1 = G_2 = \dots = G_n = Y$ we obtain

Corollary 1. *The composition $X[Y]$ of graphs X and Y is strongly perfect if and only if both X and Y are strongly perfect.*

Using Lemma 4 and Theorem 1 we obtain

Corollary 2. *$\overline{X[G_1, \dots, G_n]}$ is strongly perfect if and only if the graphs $\overline{X}, \overline{G_1}, \dots, \overline{G_n}$ are strongly perfect.*

In consequence, it follows immediately

Theorem 2. *$X[G_1, \dots, G_n]$ is co-strongly perfect if and only if the graphs X, G_1, \dots, G_n are co-strongly perfect.*

REFERENCES

- [1] C. Berge and P. Duchet, *Strongly perfect graphs*, Ann. Disc. Math. **21** (1984) 57–61.
- [2] M. Borowiecki and A. Szelecka, *One factorizations of the generalized Cartesian product and of the X -join of regular graphs*, Discussiones Mathematicae **13** (1993) 15–19.
- [3] M. Kwaśnik and A. Szelecka, *Strong perfectness of the generalized Cartesian product of graphs*, accepted for publication in the special issue of Discrete Math., devoted to the Second Kraków Conference on Graph Theory, Zakopane 1994.

- [4] E. Mandrescu, *Strongly perfect product of graphs*, Czech. Math. Journal, **41** (116) (1991) 368–372.
- [5] G. Ravindra and D. Basavayya, *Co-strongly perfect bipartite graphs*, Jour. Math. Phy. Sci. **26** (1992) 321–327.

Received 26 April 1996
Revised 8 October 1996