

COLOURING GRAPHS WITH PRESCRIBED INDUCED CYCLE LENGTHS

BERT RANDEATH

Institut für Informatik
Universität zu Köln
D–50969 Köln, Germany

e-mail: randerath@informatik.uni-koeln.de

AND

INGO SCHIERMEYER

Fakultät für Mathematik und Informatik
TU Bergakademie Freiberg
D–09596 Freiberg, Germany

e-mail: schierme@mathe.tu-freiberg.de

Abstract

In this paper we study the chromatic number of graphs with two prescribed induced cycle lengths. It is due to Sumner that triangle-free and P_5 -free or triangle-free, P_6 -free and C_6 -free graphs are 3-colourable. A canonical extension of these graph classes is $\mathcal{G}^I(4, 5)$, the class of all graphs whose induced cycle lengths are 4 or 5. Our main result states that all graphs of $\mathcal{G}^I(4, 5)$ are 3-colourable. Moreover, we present polynomial time algorithms to 3-colour all triangle-free graphs G of this kind, i.e., we have polynomial time algorithms to 3-colour every $G \in \mathcal{G}^I(n_1, n_2)$ with $n_1, n_2 \geq 4$ (see Table 1). Furthermore, we consider the related problem of finding a χ -binding function for the class $\mathcal{G}^I(n_1, n_2)$. Here we obtain the surprising result that there exists no linear χ -binding function for $\mathcal{G}^I(3, 4)$.

Keywords: colouring, chromatic number, induced subgraphs, graph algorithms, χ -binding function.

2000 Mathematics Subject Classification: 05C15, 05C75, 05C85.

1. Introduction and Results

We consider finite undirected simple graphs. For terminology and notation not defined here we refer to [1]. As introduced by Gyárfás [6], a family \mathcal{G} of graphs is called χ -bound with χ -binding function f , if $\chi(G') \leq f(\omega(G'))$ holds whenever G' is an induced subgraph of $G \in \mathcal{G}$.

Erdős [4] showed that for each pair g, k with $g, k \geq 4$ there exist graphs with girth g and chromatic number k . Hence, triangle-free graphs can have arbitrary large chromatic number. Sumner [13] showed that triangle-free and P_5 -free or triangle-free, P_6 -free and C_6 -free graphs are 3-colourable.

For $t \geq 5$ define \mathcal{G}_t as the class of all triangle-free graphs which are P_t -free and C_i -free for $6 \leq i \leq t$. For $k \geq 1$ and $3 \leq n_1 < n_2 < \dots < n_k$ let $\mathcal{G}^I(n_1, n_2, \dots, n_k)$ be the class of all graphs whose induced cycle lengths are equal to one of n_1, n_2, \dots, n_k . Thus

$$\mathcal{G}_5 \subset \mathcal{G}_6 \subset \mathcal{G}_7 \subset \dots \subset \mathcal{G}^I(4, 5)$$

and all graphs G of \mathcal{G}_5 and \mathcal{G}_6 are 3-colourable by Sumners result. Note that all graphs of \mathcal{G}_t have diameter at most $t - 2$ whereas graphs of $\mathcal{G}^I(4, 5)$ can have arbitrary diameter.

Our research was motivated by the question whether 3-colourability still holds for a superclass \mathcal{G}_t (of \mathcal{G}_5 and \mathcal{G}_6) for some $t \geq 7$. Theorem 1 states that all graphs of $\mathcal{G}^I(4, 5)$ are 3-colourable. Hence, the answer is affirmative for each $t \geq 7$. Moreover, we can guarantee a 3-colouring with some additional properties. For a fixed integer $p \geq 2$ we call a graph $G \in \mathcal{G}^I(4, 2p + 1)$ 3^* -colourable with root v , if there is a 3-colouring of G such that $G[N_G^p(v)]$ is coloured with two colours, where $N_G^p(v)$ is the set of vertices having distance p from v . Observe that this definition implies the following useful property: If G is 3^* -colourable with root v , then we can choose a 3-colouring such that $G[N_G^i(v)]$ is coloured monochromatic for every $1 \leq i < p$ and $G[N_G^p(v)]$ is coloured with two colours. Furthermore, if this property holds for every vertex of $G \in \mathcal{G}^I(4, 2p + 1)$, then we call G 3^* -colourable. This definition is motivated by the following observation.

If $G_1, G_2 \in \mathcal{G}^I(4, 2p + 1)$ and $v_i \in G_i$ for $i = 1, 2$, then the new graph G^* with vertex set $V(G^*) = V(G_1 - v_1) \cup V(G_2 - v_2)$ and edge set $E(G^*) = E(G_1 - v_1) \cup E(G_2 - v_2) \cup \{u_1 u_2 \mid u_i \in N_{G_i}(v_i) \text{ for } i = 1, 2\}$ is likewise a member of $\mathcal{G}^I(4, 2p + 1)$. The invariance of $\mathcal{G}^I(4, 2p + 1)$ concerning this graph operation reasons the equivalence of 3^* - and 3-colourability for the class $\mathcal{G}^I(4, 2p + 1)$.

Theorem 1. *Every graph of $\mathcal{G}^I(4, 2p + 1)$ with $p \geq 2$ is 3^* -colourable.*

The proof of Theorem 1 is based on decomposition and provides a polynomial time algorithm to 3^* -colour a given graph $G \in \mathcal{G}^I(4, 2p + 1)$. Note that the class $\mathcal{G}^I(4, 5)$ is a canonical extension of $\mathcal{G}^I(4)$, which are the well-known chordal bipartite graphs (e.g. see [2]). Very recently Brandt [3] examined the maximal (with respect to edge addition) triangle-free members of the class $\mathcal{G}^I(4, 5)$ with emphasis on graph homomorphisms. Brandt also observed that the class $\mathcal{G}^I(4, 5)$ is a natural extension of $\mathcal{G}^I(4)$ — the chordal bipartite graphs — and he introduced for members of $\mathcal{G}^I(4, 5)$ the terminology of *chordal triangle-free graphs*.

Motivated by the first theorem we consider next the classes $\mathcal{G}^I(2q, 2p + 1)$ and $\mathcal{G}^I(2p' + 1, 2q')$ for $q, q' \geq 3$ and $p, p' \geq 2$. But first we will examine the larger class $\mathcal{G}^I(n_1, n_2, \dots, n_k)$ with $n_1 \geq 5$. A graph G is *r-degenerate*, if there exists an ordering (v_1, \dots, v_n) of $V(G)$ such that $d_{G[\{v_i, \dots, v_n\}]}(v_i) \leq r$ for all $1 \leq i \leq n$.

Theorem 2. *Every graph of $\mathcal{G}^I(n_1, n_2, \dots, n_k)$ with $k \geq 1$ and $n_1 \geq 5$ is $(k + 1)$ -degenerate. Especially, every vertex v of G being an endvertex of a longest induced path of G satisfies $d_G(v) \leq k + 1$.*

Corollary 3. *Every graph of $\mathcal{G}^I(n_1, n_2, \dots, n_k)$ with $k \geq 1$ and $n_1 \geq 5$ is $(k + 2)$ -colourable.*

The last result reveals an interesting relation to the colouring properties of graphs of the class $\mathcal{G}(n_1, n_2, \dots, n_k)$, the class of all graphs whose (not necessarily induced) cycle lengths are equal to one of n_1, n_2, \dots, n_k . Now let G be a graph with r different odd and s different even cycle lengths (which need not to be induced). In [10] Mihók and Schiermeyer presented a polynomial time colouring algorithm, called MAXBIP, which recursively constructs maximal bipartite subgraphs. Based on MAXBIP they proved the following theorem, answering thereby a question of B. Bollobás and P. Erdős [5].

Theorem 4 (Mihók and Schiermeyer [10], 1997). $\chi(G) \leq \min\{2r + 2, 2s + 3\}$.

With $k = r + s$ this also implies $\chi(G) \leq k + 2$. The question of B. Bollobás and P. Erdős [5] only concerned $(2r + 2)$ -colourability of graphs with r different odd cycle lengths (which need not to be induced). This question was

first answered affirmative by A. Gyárfás [7]. Additional informations and a related conjecture can be found in the excellent book [8] of T. Jensen and B. Toft on graph colouring problems.

Obviously, Corollary 3 is best possible for $k = 1$. But for $k = 2$ we are able to improve Corollary 3. For the next theorem we need to recall the definition of the famous Petersen graph P^* . This 3-regular, non-bipartite graph P^* of order 10 is a member of the class $\mathcal{G}^I(5, 6)$. The Petersen graph P^* consists of two disjoint induced 5-cycles $C^1 = a_0a_1a_2a_3a_4a_0$ and $C^2 = b_0b_1b_2b_3b_4b_0$ and the additional edges $a_0b_0, a_1b_3, a_2b_1, a_3b_4$ and a_4b_2 . Obviously P^* is 3-colourable.

Theorem 5. *Every graph G of $\mathcal{G}^I(2q, 2p + 1)$ or $\mathcal{G}^I(2p' + 1, 2q')$ with $q, q' \geq 3$ and $p, p' \geq 2$ fulfills at least one of the following properties:*

1. G is bipartite;
2. G satisfies $\delta(G) \leq 2$;
3. $G \in \mathcal{G}^I(5, 6)$ and one of the following properties holds:
 - (a) $G \cong P^*$;
 - (b) G contains a clique cutset, i.e., K_1 or K_2 clique cutset.

Every graph G of $\mathcal{G}^I(2q, 2p + 1)$ or $\mathcal{G}^I(2p' + 1, 2q')$ with $q, q' \geq 3$ and $p, p' \geq 2$ not isomorphic to P^* fulfills at least one of the three properties because of Theorem 5. Testing whether G is bipartite, has minimal degree two or contains a complete cutset of size at most two can be done very efficiently. Moreover, if $G \in \mathcal{G}^I(5, 6)$ is non-bipartite, $\delta(G) \geq 3$ and contains no complete cutset, then $G \cong P^*$, which obviously is 3-colourable. Hence, Theorem 5 provides a polynomial time algorithm to 3-colour a given graph $G \in \mathcal{G}^I(2q, 2p + 1)$ or $G \in \mathcal{G}^I(2p' + 1, 2q')$ with $q, q' \geq 3$ and $p, p' \geq 2$. This algorithm (recursively) makes use of the fact that the graph (in question) is bipartite, has a vertex of degree at most two, is isomorphic to the Petersen graph or the graph can be decomposed into two smaller graphs according to a complete cutset of size at most two.

Corollary 6. *Every graph G of $\mathcal{G}^I(2q, 2p + 1)$ or $\mathcal{G}^I(2p' + 1, 2q')$ with $q, q' \geq 3$ and $p, p' \geq 2$ is 3-colourable.*

Now we consider the related problem of finding a (best possible) χ -binding function f^* for $\mathcal{G}^I(n_1, n_2)$ and for completeness also for its subclasses $\mathcal{G}^I(n_1)$.

Recall that a graph is *perfect* if for each induced subgraph H of G the chromatic number $\chi(H)$ equals the corresponding clique number $\omega(H)$. Furthermore the *lexicographic product* $G_1[G_2]$ of two graphs G_1 and G_2 contains the vertex set $V(G_1[G_2]) = V(G_1) \times V(G_2)$ and two vertices (a, b) and (c, d) are adjacent in $G_1[G_2]$ if a is adjacent to c in G_1 or $a = c$ and b is adjacent to d in G_2 .

For convenience we drop the condition that n_1 is always smaller than n_2 in the definition of $\mathcal{G}^I(n_1, n_2)$.

(I) n_1, n_2 are even :

For even n_1 and n_2 all graphs of $\mathcal{G}^I(n_1)$ and $\mathcal{G}^I(n_1, n_2)$ are bipartite and thus perfect with $f^*(\omega) = \omega \leq 2$.

(II) n_1 is even, n_2 is odd : (A) $n_2 \geq 5$:

By our results (Theorem 1, Corollary 6) every graph of $\mathcal{G}^I(n_2)$ and $\mathcal{G}^I(n_1, n_2)$ is 3-colourable, i.e., with $\omega \leq 2$ we have $f^*(\omega) = \omega + 1 \leq 3$.

(II) n_1 is even, n_2 is odd : (B1) $n_2 = 3$ and $n_1 \geq 6$:

Recently, Rusu [11] proved that all members of a superclass of $\mathcal{G}^I(3, 2q)$ are perfect for any $q \geq 3$. Hence, we also have $f(\omega) = \omega$. A well-known subclass of $\mathcal{G}^I(3, 2q)$ is $\mathcal{G}^I(3)$ containing the chordal graphs.

(II) n_1 is even, n_2 is odd : (B2) $n_2 = 3$ and $n_1 = 4$:

In 1987 Gyárfás [6] conjectured (motivated by the Strong Perfect Graph Conjecture) that there exists a χ -binding function for $\mathcal{G}^I(3, 4)$. But this Conjecture is still open. In [6] Gyárfás also suggested to examine whether there exists a linear χ -binding function for hereditary classes of graphs. For $\mathcal{G}^I(3, 4)$ we have constructed the following sequence of graphs (H_i) . Starting with $H_1 = \bar{C}_7$, the complement of the 7-cycle, we define $H_{i+1} = \bar{C}_7[H_i]$, the lexicographic product of the graphs \bar{C}_7 and H_i . Note that $\omega(H_{i+1}) = 3\omega(H_i)$. Furthermore, in any colouring of H_{i+1} we need for each copy of H_i at least $\chi(H_i)$ different colours. With $\alpha(\bar{C}_7) = 2$ we then observe that every colour of a colouring of H_{i+1} appears in at most two different copies of H_i . Hence, H_i has the order $n(H_i) = 7^i$, the independence number $\alpha(H_i) = 2^i$ and the clique number $\omega(H_i) = 3^i$. Therefore, its chromatic number $\chi(H_i)$ is at least $(7/2)^i$. Thus, the χ -binding function f^* for $\mathcal{G}^I(3, 4)$ satisfies $f^*(\omega) \geq (7/6)^i \omega$ for every integer i . Hence, we obtain the following surprising result:

Theorem 7. *There exists no linear χ -binding function for $\mathcal{G}^I(3, 4)$.*

It is noteworthy to mention that $\mathcal{G}^I(3, 4)$ contains all weakly triangulated graphs. Recently, Scott [12] achieved some related results.

(III) n_1, n_2 are odd : (A1) $n_1, n_2 \geq 5$:

Markossian, Gasparian and Reed [9] showed that all triangle-free and even-hole-free graphs are 2-degenerate and thus are 3-colourable. Hence, $f^*(\omega) = \omega + 1 \leq 3$ is a χ -binding function for $\mathcal{G}^I(n_1)$ and $\mathcal{G}^I(n_1, n_2)$.

(III) n_1, n_2 are odd : (A2) $n_1 = 3$:

It is an open problem, whether there exists a linear χ -binding function for $\mathcal{G}^I(3, n_2)$. The graph-sequence $G_r = C_{n_2}[K_r]$, the lexicographic product of the odd cycle C_{n_2} and the complete graph K_r , reveals that we have $f^*(\omega) \geq ((n_2 + 1)/(n_2 - 1))\omega$ for every χ -binding function. We expect that

$$f^*(\omega) = ((n_2 + 1)/(n_2 - 1))\omega.$$

Table 1. χ -binding function f^* for $\mathcal{G}^I(n_1, n_2)$.

$\begin{matrix} \rightarrow \\ n_1, n_2 \\ \downarrow \end{matrix}$	3	4	odd ≥ 5	even ≥ 6
3	$f^*(\omega) = \omega$ chordal	\nexists linear f^* Theorem 7 Conj.[6]: $\exists f^*$	$f^*(\omega) \geq ((n_2 + 1)/(n_2 - 1))\omega$ Conj.: " = "	$f^*(\omega) = \omega$ Rusu [11]
4		$f^*(\omega) = \omega \leq 2$ chordal bipartite	$f^*(\omega) = \omega + 1 \leq 3$ Theorem 1	$f^*(\omega) = \omega \leq 2$ \subset bipartite
odd ≥ 5			$f^*(\omega) = \omega + 1 \leq 3$ Markossian,... [9]	$f^*(\omega) = \omega + 1 \leq 3$ Corollary 6
even ≥ 6				$f^*(\omega) = \omega \leq 2$ \subset bipartite

2. Proofs

The following well-known easy observation provides a very useful property. If a graph G contains a pair u, v of nonadjacent vertices with $N_G(u) \subseteq N_G(v)$, then any proper k -colouring of $G - u$ can easily be extended to a proper k -colouring of G .

Therefore we only have to consider those graphs G having the property (*):

(*) If $uv \notin E(G)$, then there exist vertices (*private neighbours*) $p_u \in (N_G(u) - N_G(v))$ and $p_v \in (N_G(v) - N_G(u))$.

The next lemma provides a useful property of triangle-free and C_6 -free graphs, which will be used in the proof of Theorem 1. This class of graphs forms a superclass of all classes $\mathcal{G}^I(4, 2p+1)$ with $p \geq 2$.

Lemma 8. *Let G be a triangle-free and C_6 -free graph satisfying property (*). Then for every vertex x of degree $d_G(x) = k \geq 3$ with neighbours x_1, x_2, \dots, x_k there exists a pair x_i, x_j of neighbours such that $N_G(x_i) \cap N_G(x_j) \cap N_G^2(x) = \emptyset$.*

Proof. Let G be a triangle-free and C_6 -free graph satisfying property (*). Further suppose to the contrary that there exists a vertex x of degree $d_G(x) = k \geq 3$ with neighbours x_1, x_2, \dots, x_k such that $N_G(x_i) \cap N_G(x_j) \cap N_G^2(x) \neq \emptyset$ for all pairs i, j with $1 \leq i < j \leq k$. Choose a vertex $v \in N_G^2(x)$ such that $|N_G(x) \cap N_G(v)| = p$ is maximum. By (*) we know that $p \leq k - 1$. We may assume that $N_G(x) \cap N_G(v) = \{x_1, \dots, x_p\}$. By the assumption there exists a vertex $w \in N_G^2(x)$ with w likewise adjacent to a vertex $x_i \in \{x_1, \dots, x_p\}$ and to a vertex $x_j \in \{x_{p+1}, \dots, x_k\}$. Hence there is a vertex $w \in N_G^2(x) - \{v\}$ with $N_G(v) \cap N_G(w) \cap N_G(x) \neq \emptyset$ and $N_G(w) \cap \{x_{p+1}, \dots, x_k\} \neq \emptyset$. Choose a vertex w with $N_G(w) \cap \{x_{p+1}, \dots, x_k\} \neq \emptyset$ such that $|N_G(v) \cap N_G(w) \cap N_G(x)| = q$ is maximum. Hence $1 \leq q \leq p - 1$ by the choice of v and w . We may assume that $N_G(v) \cap N_G(w) \cap N_G(x) = \{x_1, \dots, x_q\}$ and that $x_{p+1} \in N_G(w)$. By the hypothesis there exists a vertex $u \in N_G^2(x) - \{v, w\}$ such that $x_p, x_{p+1} \in N_G(u)$. By the choice of p and q there is a vertex $x_s \in \{x_1, \dots, x_q\}$ such that $x_s \notin N_G(u)$. Now $G' := G[\{x_s, x_p, x_{p+1}, v, w, u\}] \cong C_6$ or G' contains a triangle, a contradiction. ■

Proof of Theorem 1. We will prove the theorem by induction. We may assume that G is a connected graph and fulfills property (*). By the induction hypothesis there is a 3^* -colouring with root y for every vertex y of any induced subgraph H of G with $|V(H)| < |V(G)|$. Now let x be an arbitrary vertex of G . If $\text{dist}_G(x, y) < p$ for all vertices $y \in V(G - x)$, then we can easily 3^* -colour G with root x . Hence we may assume that $\text{dist}_G(x, y) \geq p$ for at least one vertex $y \in V(G - x)$.

Case 1. Assume that $d_G(x) = 1$ and let $xz \in E(G)$.

Then by induction hypothesis there is a 3*-colouring with root z of $G - x$, which can be easily extended to a 3*-colouring with root x of G .

Case 2. Assume that $d_G(x) \geq 2$.

Also, if x is a cutvertex of G , then we can easily 3*-colour G with root x . Hence we may assume that x is not a cutvertex of G . Note that x and every neighbour of x has a degree of at least two.

We now consider $N_G^i(x)$ for $1 \leq i < p$. If there is a vertex $y \in N_G^i(x)$ such that

$$N_G(y) \cap N_G^{i+1}(x) \subseteq \left(\bigcup_{z \in N_G^i(x) - \{y\}} N_G(z) \right) \cap N_G^{i+1}(x),$$

then the levels $N_G^h(x)$ with $1 \leq h < p$ and $h \neq i$ are the same in G and $G - y$. Thus, we can reduce G to $G - y$ and a 3*-colouring with root x of $G - y$ can be extended to a 3*-colouring with root x of G since $G[N_G^{i-1}(x)]$ is independent and monochromatic, $G[N_G^i(x) - \{y\}]$ is independent and monochromatic and $G[N_G^{i+1}(x)]$ is 2-coloured. Consequently, we can assume for the remaining part of the proof that

Claim 1. Every vertex $y \in N_G^i(x)$ has a 'private neighbour' in $N_G^{i+1}(x)$ for $1 \leq i < p$.

Subcase 2.1. Assume that $d_G(x) \geq 3$.

Let u^1, v^1 and w^1 be three vertices of $N_G^1(x)$. Then by Claim 1 there are three 'paths of private neighbours' $u^1 u^2 \dots u^p, v^1 v^2 \dots v^p$ and $w^1 w^2 \dots w^p$. Set $u = u^{p-1}, v = v^{p-1}$ and $w = w^{p-1}$. Let U^p, V^p, W^p be the set of private neighbours of u, v and w , respectively.

If $p \geq 3$, then $N_G^p(x) \cap N_G(y) \cap N_G(z) = \emptyset$ for all pairs $y, z \in \{u, v, w\}$. Since otherwise there would be an induced C_{2p} - a contradiction. If $p = 2$, then we can choose u and v by Lemma 8 such that $N_G^2(x) \cap N_G(u) \cap N_G(v) = \emptyset$.

Now observe that $G[U^p, V^p], G[U^p, W^p]$ and $G[W^p, V^p]$ are $2K_2$ -free (and bipartite), since otherwise there would be an induced C_6 .

A useful property of every bipartite and $2K_2$ -free graph H is the existence of a labelling of the vertices of each partition set $X = \{x_1, \dots, x_k\}$, such that $N_H(x_i) \subset N_H(x_j)$ if $i \leq j$.

Then we easily deduce that there are sets $U_{(c)}^p, U_{(e)}^p, V_{(c)}^p$ and $V_{(e)}^p$ such that $U^p = U_{(c)}^p \cup U_{(e)}^p, V^p = V_{(c)}^p \cup V_{(e)}^p, G[U_{(e)}^p \cup V_{(e)}^p]$ is edgeless and

$G[U_{(c)}^p \cup V_{(c)}^p]$ is complete bipartite. Note that the partition of U^p and V^p into $U_{(c)}^p, U_{(e)}^p, V_{(c)}^p$ and $V_{(e)}^p$ is not necessarily unique. But, if $G[U^p, V^p]$ is not a complete bipartite graph, it is always possible to choose a partition of U^p and V^p into $U_{(c)}^p, U_{(e)}^p, V_{(c)}^p$ and $V_{(e)}^p$ such that $U_{(e)}^p \neq \emptyset$ and $V_{(e)}^p \neq \emptyset$!

Subcase 2.1.1. $G[U^p, V^p]$ is not a complete bipartite subgraph.

Then there are two non-adjacent vertices $y \in U^p$ and $z \in V^p$. Now we can choose a partition of U^p and V^p into $U_{(c)}^p, U_{(e)}^p, V_{(c)}^p$ and $V_{(e)}^p$ such that $y \in U_{(e)}^p$ and $z \in V_{(e)}^p$. Set $S = \{v \in V(G) \mid \text{dist}_G(x, v) \leq p-1\} \cup U_{(c)}^p \cup V_{(c)}^p$.

Subcase 2.1.1.1. For every vertex $y \in U_{(e)}^p$ and $z \in V_{(e)}^p$ there exists a path connecting y and z in $G-S$.

Now we choose $y \in U_{(e)}^p$ and $z \in V_{(e)}^p$ such that $\text{dist}_{G-S}(y, z) = \min\{\text{dist}_{G-S}(y', z') \mid y' \in U_{(e)}^p \text{ and } z' \in V_{(e)}^p\}$. Then obviously there would be an induced cycle of length at least $(2p+1)+1 > 2p+1 > 4$, a contradiction.

Subcase 2.1.1.2. There exist vertices $y \in U_{(e)}^p$ and $z \in V_{(e)}^p$ such that there is no path connecting y and z in $G-S$.

Now let H_1 be a component of $G-S$ and H_2 be the remaining part of $G-S$. Set $G_i = G[V(H_i) \cup S]$ for $i = 1, 2$. Then there is a 3^* -colouring with root x for each G_i with $i = 1, 2$. We can choose these 3^* -colourings in such a way that the vertices of S always receive the same colours. Hence we obtain a 3^* -colouring with root x of G . In the following, if we will apply this subprocedure, we shortly refer that we *apply decomposition*.

Subcase 2.1.2. $G[U^p, V^p]$ is a complete bipartite subgraph.

If $p \geq 3$, then $G[U^p, W^p]$ and $G[W^p, V^p]$ are complete as well contradicting that G is triangle-free.

Let $p = 2$. If we also have $N_G^2(x) \cap N_G(u) \cap N_G(w) = \emptyset$ or $N_G^2(x) \cap N_G(v) \cap N_G(w) = \emptyset$, then we either apply decomposition or we deduce that $G[U^p, W^p]$ or $G[W^p, V^p]$ is complete bipartite. But then there exists a path ayz or azy with $a \in W^p, y \in U^p$ and $z \in V^p$ implying the existence of an induced cycle C_6 or ayz induces a triangle – a contradiction. Hence $N_G^2(x) \cap N_G(u) \cap N_G(w) \neq \emptyset$ and $N_G^2(x) \cap N_G(v) \cap N_G(w) \neq \emptyset$. Let $a \in N_G^2(x) \cap N_G(u) \cap N_G(w)$ and $b \in N_G^2(x) \cap N_G(v) \cap N_G(w)$. We now consider the subgraph $G[U'^2, V'^2]$ with $U'^2 = N_G^2(x) \cap (N_G(u) - N_G(v))$ and $V'^2 = N_G^2(x) \cap (N_G(v) - N_G(u))$. Recall that $N_G^2(x) \cap (N_G(u) \cup N_G(v)) = U'^2 \cup V'^2$ and $G[U^2, V^2]$ is a subgraph of $G[U'^2, V'^2]$. Note that we obtain analogously that $G[U'^2, V'^2]$ is $2K_2$ -free (and bipartite). With $a \in U'^2, b \in V'^2, \{a, b\} \subset N_G(w)$ and G being triangle-free we deduce that $G[U'^2, V'^2]$ is not complete.

But now we analogously can apply decomposition. This settles the case that $d_G(x) \geq 3$.

Subcase 2.2. Suppose that $d_G(x) = 2$.

By the induction hypothesis there is a 3^* -colouring with root y for every vertex y with $d_G(y) \neq 2$ and for every y of any induced subgraph H of G with $|V(H)| < |V(G)|$. Note that $N_G(u^1) \cap N_G(v^1) \cap N_G^2(x) = \emptyset$ by (*).

Let $p = 2$. If $d_G(u^1) = d_G(v^1) = 2$, then because the vertices u^2, u^1, x, v^1 and v^2 are lying on a cycle (! x is no cutvertex of G !) we obtain adjacency of v^2 and u^2 . Now it is not very difficult to extend an arbitrary 3-colouring of $G - \{u^1, x, v^1\}$ to a 3^* -colouring with root x of G .

Let $d_G(u^1) \geq 3$. By the induction hypothesis there is a 3^* -colouring with root u^1 of G . Since $\{v^1\} \cup (N_G(v^1) \cap N_G^2(x))$ induces a star we may choose the colours in such a way that (e.g.) u^1 and v^1 receive colour 1, x and all vertices of $N_G(u^1) \cap N_G^2(x)$ get colour 2 and finally all vertices of $N_G(v^1) \cap N_G^2(x)$ receive colour 3. This gives a 3^* -colouring with root x of G .

Let $p \geq 3$. Since $N_G^i(x)$ is independent for $1 \leq i < p$ and G being C_{2q} -free for $q \geq 3$, we obtain that $N_G^i(u^1) \cap N_G^i(v^1) \cap N_G^{i+1}(x) = \emptyset$ for $2 \leq i < p$. Hence the set $N_G^p(x)$ is the disjoint union of the sets $N_G^{p-1}(u^1)$ and $N_G^{p-1}(v^1)$. Now the C_{2q+1} -freeness of G ($1 \leq q < p$) forces that $N_G^{p-1}(u^1)$ as well as $N_G^{p-1}(v^1)$ form independent sets. Moreover, it is not difficult to see that $N_G^p(x)$ induces a $2K_2$ -free bipartite graph. Likewise to the previous case $d_G(x) \geq 3$ we can apply decomposition or $N_G^p(x)$ induces a complete bipartite graph. In the latter case we can proceed analogously to the $p = 2$ subcase.

If all vertices of $N' := \{v \in V(G) \mid \text{dist}_G(x, v) \leq p - 1\}$ have degree two, then again $N_G^p(x)$ induces an edge. Now it is not very difficult to extend an arbitrary 3-colouring of $G - N'$ to a 3^* -colouring with root x of G .

Now let u^i (with $1 \leq i < p$) be a vertex of degree at least three having a minimum distance concerning x . By the induction hypothesis there is a 3^* -colouring with root u^i of G . Note that $N_G^p(x)$ is bicoloured, since $G[U^p, V^p]$ is complete bipartite. Now again it is not very difficult to recolour this 3-colouring in such a way that we obtain a 3^* -colouring with root x of G . ■

Proof of Theorem 2. We will show that every vertex v of G being an endvertex of a longest induced path of G satisfies $d_G(v) \leq k + 1$. Observe that this implies $\delta(G) \leq k + 1$. Let $P := v_1v_2 \dots v_t$ be a longest induced path of G . Suppose $d_G(v_t) = s \geq k + 2$. Let $N_G(v_t) = \{v_{t-1}, u_1, \dots, u_{s-1}\}$. By the choice of P every vertex u_i has a neighbour on P . For each i (with

$1 \leq i \leq s-1$) let $j(i)$ be the largest integer less than or equal to $t-1$ such that $u_i v_{j(i)} \in E(G)$. Since G is triangle-free and contains no induced C_4 , no two values of $j(1), \dots, j(s-1)$ are equal. Hence there are $s-1 \geq k+1$ induced cycles of different lengths $t - j(i) + 2$ for $1 \leq i \leq s-1$, a contradiction. ■

Proof of Theorem 5.

Case 1. Suppose there is a graph $G \in \mathcal{G}^I(2q, 2p+1)$ with $q \geq 3$ and $p \geq 2$ (and $2p+1 > 2q$), which satisfies $\delta(G) \geq 3$ and is non-bipartite.

Let $P := v_1 v_2 \dots v_t$ be a longest induced path of G . We deduce that $d_G(v_t) = 3$ because of Theorem 2. Let $N_G(v_t) = \{v_{t-1}, u_1, u_2\}$. By the choice of P the vertices u_1 and u_2 each have at least one neighbour on P . For $i = 1, 2$ let $j(i)$ be the largest integer less than or equal to $t-1$ such that $u_i v_{j(i)} \in E(G)$. Hence, $\{j(1), j(2)\} = \{t - (2p-1), t - (2q-2)\}$ and say $u_1 v_{t-(2p-1)}, u_2 v_{t-(2q-2)} \in E(G)$. Furthermore, there exists a maximum $r \geq 1$ such that $r(2q-2) < 2p-1$ and $u_2 v_{t-i(2q-2)} \in E(G)$ for $1 \leq i \leq r$. Now the cycle $v_t u_1 v_{t-(2p-1)} v_{t-2p+2} \dots v_{t-r(2q-2)} u_2 v_t$ is induced and has odd length. Hence, $(t - r(2q-2)) - (t - (2p-1)) + 4 = 2p+1$ and by rearranging $2 = r(2q-2)$ contradicting $q \geq 3$.

Case 2. Suppose there is a graph $G \in \mathcal{G}^I(2p+1, 2q)$ with $q \geq 3$ and $p \geq 2$ (and $2p+1 < 2q$!), which satisfies $\delta(G) \geq 3$ and is non-bipartite.

Let again $P := v_1 v_2 \dots v_t$ be a longest induced path of G . In the remaining proof we will deduce several structural statements concerning this longest induced path P of G . It is important to note that these statements also hold for every longest induced path P' of G . For convenience we call a vertex $v \in V(G)$ an *i-type vertex*, if there exists a longest induced path $P' = v'_1 v'_2 \dots v'_t$ of G with $v = v'_i$. Again because of Theorem 2 we obtain $d_G(v_i) = 3$ or more generally

Claim 1. Every t -type vertex has the degree 3.

Now let $N_G(v_t) = \{v_{t-1}, u_1, u_2\}$. Again for $i = 1, 2$ let $j(i)$ be the largest integer less than or equal to $t-1$ such that $u_i v_{j(i)} \in E(G)$. Analogously, we have say $j(1) = t - (2q-2)$, $j(2) = t - (2p-1)$ and $u_1 v_{t-(2q-2)}, u_2 v_{t-(2p-1)} \in E(G)$. Furthermore, there exists a maximum $r \geq 1$ such that $r(2p-1) < 2q-2$ and $u_2 v_{t-i(2p-1)} \in E(G)$ for $1 \leq i \leq r$. Now the cycle $v_t u_1 v_{t-(2q-2)} v_{t-2q+3} \dots v_{t-r(2p-1)} u_2 v_t$ is induced and has odd length. Hence, $(t - r(2p-1)) - (t - (2q-2)) + 4 = 2p+1$ and by rearranging $2q = (r+1)(2p-1)$.

We now examine a special case. Suppose that G contains no induced path P_{2q} containing $2q$ vertices.

Markossian, Gasparian and Reed [9] showed that all triangle-free and even-hole-free graphs are 2-degenerate. Hence G has to contain an induced even cycle $C = c_0c_1 \dots c_{2q-2}c_{2q-1}c_0$. Furthermore, since $\delta(G) \geq 3$ each vertex of C is adjacent to at least one vertex not lying on the cycle C , e.g. $\{c_0d_0, c_1d_1\} \subset E(G)$. The P_{2q} -freeness, $G \in \mathcal{G}^I(2p+1, 2q)$ and $2q = (r+1)(2p-1)$ forces that $N_G(d_i) \cap V(C) = \{c_i, c_{i+(2p-1)}, c_{i+2(2p-1)} \dots, c_{i+r(2p-1)}\}$ for $i = 0, 1$. But then the vertex set $\{c_0, d_0, c_{2p-1}, c_{2p}, d_1, c_1\}$ induces a 6-cycle enforcing that $q = 3$ and $p = 2$. Since G is non-bipartite G has to contain an induced 5-cycle $C^1 = a_0a_1a_2a_3a_4a_0$. Furthermore, since $\delta(G) \geq 3$ each vertex of C^1 is adjacent to at least one vertex not lying on the cycle C^1 , e.g. $\{a_0b_0, a_1b_3, a_2b_1, a_3b_4, a_4b_2\} \subset E(G)$. The P_6 -freeness forces that $C^2 = b_0b_1b_2b_3b_4b_0$ induces a 5-cycle. Now the C_4 -freeness of G forces that all vertices of C^1 have degree 3. Analogously, every vertex of C^2 has degree 3. Thus, we have $N_G(V(C^i)) - V(C^i) = V(C^{3-i})$ for $i = 1, 2$, i.e., $G \cong P^*$. Hence, in the following a longest induced path P contains at least $2q$ vertices.

We now examine the $(t-1)$ -type vertex v_{t-1} . Since $\delta(G) \geq 3$, P is chordless and G is triangle-free, v_{t-1} has a neighbour $w \in V(G) - (V(P) \cup N_G(v_t))$.

Subcase 2.1. w has no neighbour on P .

Then $P' := v_1v_2 \dots v_{t-1}w$ is also a longest induced path of G and the t -type vertex w has (exactly) two neighbours $w_1, w_2 \in V(G) - (V(P) \cup N_G(v_t) \cup \{w\})$. Because w_1 corresponds to w_2 and w_2 to w_1 , we deduce that say $w_1v_{t-(2q-2)} \in E(G)$ and $w_2v_{t-(2p-1)} \in E(G)$. Now $C^1 := w_1v_{t-(2q-2)}u_1v_tv_{t-1}ww_1$ forms a cycle of length six. Therefore, the only possible case is again $q = 3$ and $p = 2$, i.e., $G \in \mathcal{G}^I(5, 6)$. Now the C_3 - and C_4 -freeness of G forces that the set $\{w, w_1, v_{t-1}, v_t, u_2, v_{t-4}, v_{t-3}\}$ induces a C_7 , a contradiction.

Note that the C_3 - and C_4 -freeness of G always forces that every 7-cycle is an induced one.

Subcase 2.2. w has a neighbour on P .

Then $wv_{t-(2q-1)} \in E(G)$ or $wv_{t-2p} \in E(G)$. Now $C^1 := wv_{t-(2q-1)}v_{t-2q-2}u_1v_tv_{t-1}w$ or $C^2 := wv_{t-2p}v_{t-(2p-1)}u_2v_tv_{t-1}w$ induces a cycle of length six, i.e. $G \in \mathcal{G}^I(5, 6)$. Therefore, we have either $wv_{t-5} \in E(G)$ or $wv_{t-4} \in E(G)$.

Subcase 2.2.1. $wv_{t-5} \in E(G)$.

Since G is C_4 -free, u_2 is not adjacent to w and we deduce a contradiction by considering the set $\{w, v_{t-5}, v_{t-4}, v_{t-3}, u_2, v_t, v_{t-1}\}$, which induces a C_7 .

Subcase 2.2.2. $wv_{t-4} \in E(G)$.

Claim 2. Every $(t-1)$ -type vertex has the degree 3, e.g. $d_G(v_{t-1}) = 3$.

Assume $d_G(v_{t-1}) > 3$, then there exists a vertex $\bar{w} \in (N_G(v_{t-1}) - (\{w\} \cup V(P)))$. Likewise, we deduce that \bar{w} is also adjacent to v_{t-4} . But then $\{v_{t-1}, w, \bar{w}, v_{t-4}\}$ induces a C_4 – a contradiction. Thus, we have $d_G(v_{t-1}) = 3$.

Recall that u_1 is adjacent to v_{t-4} and u_2 is adjacent to v_{t-3} .

Claim 3. Every $(t-2)$ -type vertex has the degree 3, e.g. $d_G(v_{t-2}) = 3$.

Let x be a neighbour of v_{t-2} not lying on P . If x is adjacent to v_{t-5} or v_{t-6} , then $xv_{t-5}v_{t-4}u_1v_tv_{t-1}v_{t-2}x$ or $xv_{t-6}v_{t-5}v_{t-4}wv_{t-1}v_{t-2}x$ is a cycle of length 7, a contradiction. Hence, x has no other neighbours on P . The C_4 -freeness of G forces that $xw \notin E(G)$. Furthermore, we have $xu_2 \notin E(G)$, since otherwise $xu_2v_tu_1v_{t-4}v_{t-3}v_{t-2}x$ is a cycle of length 7, a contradiction. Since $d_G(x) \geq \delta(G) = 3$, x has a neighbour $y \notin V(P) \cup \{w, u_1, u_2\}$. If y is adjacent to v_{t-4} or v_{t-5} , then $yv_{t-4}u_1v_tv_{t-1}v_{t-2}xy$ or $yv_{t-5}v_{t-4}wv_{t-1}v_{t-2}xy$ is a cycle of length 7, a contradiction. Hence, $P''' = v_1 \dots v_{t-2}xy$ is a longest induced path implying with Claim 1 and Claim 2 that $d_G(x) = d_G(y) = 3$. Observe that x is a $(t-1)$ -type and y a t -type vertex.

Claim 4. x is adjacent to u_1 .

Suppose to the contrary that $xu_1 \notin E(G)$. Analogously to P , we deduce for P''' that the $(t-1)$ -type vertex x is adjacent to a vertex $y^* \notin V(P) \cup \{y, w, u_1, u_2\}$. Because y^* corresponds to w we also deduce that y^* is adjacent to v_{t-4} . But then $y^*v_{t-4}u_1v_tv_{t-1}v_{t-2}xy^*$ is a cycle of length 7, a contradiction. Thus we have $xu_1 \in E(G)$.

Now suppose that $d_G(v_{t-2}) > 3$, i.e. there exists a vertex $x^* \in N_G(v_{t-2}) - \{v_{t-3}, v_{t-1}, x\}$. Analogously to Claim 4 we obtain $x^*u_1 \in E(G)$, a contradiction since G is C_4 -free. Therefore, we have $d_G(v_{t-2}) = 3$.

For convenience we introduce the notation z -type vertex with $z \in \{w, u_1, u_2, x, y\}$. A vertex v of G is a z -type vertex, if there exists a longest induced path P''' , such that the role of v in P''' corresponds to the role of z for the longest induced path $P = v_1v_2 \dots v_t$.

Claim 5. w is adjacent to y .

Because the $(t-1)$ -type vertex x of $P''' = v_1v_2\dots v_{t-2}xy$ is adjacent to u_1 , we obtain that u_1 is a w -type vertex. Similarly, we deduce that v_{t-1} is a x -type vertex. Recall (Claim 4) that u_1 is adjacent x and the $(t-4)$ -type vertex v_{t-4} . Since w is adjacent to the x -type vertex v_{t-1} of P''' and the $(t-4)$ -type vertex v_{t-4} of P''' the C_4 -freeness of G forces that w is the u_1 -type vertex of P''' . Because y is the t -type vertex of P''' this implies Claim 5.

Claim 6. y is adjacent to u_2 and therefore $N_G(y) = \{x, w, u_2\}$.

Assume to the contrary that y is not adjacent to u_2 . Hence there exists x' with $\{x'\} = N_G(y) - \{x, w\}$. Note that $x' \notin V(P) \cup \{u_1, u_2, x, w\}$ and since $P''' = v_1\dots v_{t-2}xy$ is a longest induced path and w is adjacent to v_{t-4} , we obtain that x' is adjacent to v_{t-3} . But then $x'v_{t-3}u_2v_tv_{t-1}wyx'$ is a cycle of length 7, a contradiction. Thus we have $N_G(y) = \{x, w, u_2\}$. Since u_2 is a t -type vertex of the longest induced path $P'' = v_1\dots v_{t-4}u_1xyu_2$ we obtain the following Claim 7 by Claim 1.

Claim 7. $d_G(u_2) = 3$.

Summarizing all claims we deduce $G' := G[\{u_2, v_t, v_{t-1}, v_{t-2}, v_{t-3}, v_{t-4}, u_1, w, x, y\}] \cong P^*$ and all six non-neighbours of v_{t-4} have degree three. If $G \cong P^*$ we are done.

Assume $G \not\cong P^*$. Note that $CS := \{v_{t-4}, v_{t-3}, u_1, w\}$ is a cutset of G . If v_{t-4} and one of its three neighbours v_{t-3}, u_1 or w is already a (complete) cutset, then we are done. Hence w.l.o.g. suppose G_2 is a component of $G - CS$ different to the six cycle $G_1 := G[\{u_2, v_t, v_{t-1}, v_{t-2}, x, y\}]$ and say $G'' := \{u_1, w\} \cup V(G_2)$ is connected. The C_4 -freeness forces that a shortest path connecting u_1 and w in G'' contains at least 2 internal vertices and in $G''' := \{u_1, w\} \cup V(G_1)$ there exists a u_1 and w connecting induced path with 3 internal vertices. Then one can construct easily an induced cycle of length at least 7, a contradiction. This settles the proof of this theorem. ■

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory and Applications (Macmillan, London and Elsevier, New York, 1976).
- [2] A. Brandstädt, Van Bang Le and J.P. Spinrad, Graph classes: a survey, SIAM Monographs on Discrete Mathematics and Applications (SIAM, Philadelphia, PA, 1999).

- [3] S. Brandt, *Triangle-free graphs without forbidden subgraphs*, Electronic Notes in Discrete Math. Vol. 3.
- [4] P. Erdős, *Graph theory and probability*, Canad. J. Math. **11** (1959) 34–38.
- [5] P. Erdős, *Some of my favourite unsolved problems*, in: A. Baker, B. Bollobás and A. Hajnal, eds. A tribute to Paul Erdős (Cambridge Univ. Press, Cambridge, 1990) 467.
- [6] A. Gyárfás, *Problems from the world surrounding perfect graphs*, Zastos. Mat. **XIX** (1987) 413–441.
- [7] A. Gyárfás, *Graphs with k odd cycle lengths*, Discrete Math. **103** (1992) 41–48.
- [8] T.R. Jensen, B.Toft, *Graph Colouring problems* (Wiley-Interscience Publications, New York, 1995).
- [9] S.E. Markossian, G.S. Gasparian and B.A. Reed, *β -Perfect Graphs*, J. Combin. Theory (B) **67** (1996) 1–11.
- [10] P. Mihók and I. Schiermeyer, *Chromatic number of classes of graphs with prescribed cycle lengths*, submitted.
- [11] I. Rusu, *Berge graphs with chordless cycles of bounded length*, J. Graph Theory **32** (1999) 73–79.
- [12] A.D. Scott, *Induced Cycles and Chromatic Number*, J. Combin. Theory (B) **76** (1999) 70–75.
- [13] D.P. Sumner, *Subtrees of a Graph and the Chromatic Number*, in: G. Chartrand, Y. Alavi, D.L. Goldsmith, L. Lesniak-Foster, and D.R. Lick, eds, The Theory and Applications of Graphs, Proc. 4th International Graph Theory Conference (Kalamazoo, Michigan, 1980) 557–576, (Wiley, New York, 1981).

Received 28 December 2000

Revised 13 May 2001