

## A PROOF OF MENGER'S THEOREM BY CONTRACTION

FRANK GÖRING

*Department of Mathematics*  
*Technical University of Ilmenau*  
*D-98684 Ilmenau Germany*

### Abstract

A short proof of the classical theorem of Menger concerning the number of disjoint  $AB$ -paths of a finite graph for two subsets  $A$  and  $B$  of its vertex set is given. The main idea of the proof is to contract an edge of the graph.

**Keywords:** connectivity, disjoint paths, digraph, Menger.

**2000 Mathematics Subject Classifications:** 05C40.

Proofs of Menger's Theorem are given in [7, 6, 4, 8, 2]. A short proof is given by T. Böhme, F. Göring and J. Harant in [1]; another short proof based on edge deletion is given by the author in [5]. The new idea here is to get a short proof by contracting an arbitrary edge of the original graph.

For terminology and notation not defined here we refer to [3]. A graph with no edges is denoted by its vertex set. Let  $G$  be a finite graph (loops and multiple edges being allowed). For an edge  $e$  of  $G$  let  $G - e$  and  $G/e$  denote the graphs obtained from  $G$  by removing  $e$  and contracting  $e$  to one vertex  $v_e$ , respectively. For (possibly empty) sets of vertices  $A$  and  $B$  of  $G$  let an  $AB$ -separator be a set of vertices of  $G$  such that the graph obtained from  $G$  by deleting these vertices contains no path from  $A$  to  $B$ . Note that a single vertex of  $A \cap B$  is considered as a path from  $A$  to  $B$ , too. An  $AB$ -connector is a subgraph of  $G$  such that each of its components is a path from  $A$  to  $B$  having only one vertex in common with  $A$  and  $B$ , respectively. In particular the empty graph is also an  $AB$ -connector. If we contract an edge incident with a vertex of  $A$  or  $B$  then the resulting vertex is considered to be a vertex of  $A$  or  $B$ , respectively.

**Theorem** (Menger, 1927). *Let  $G$  be a finite graph,  $A$  and  $B$  sets of vertices of  $G$ , and  $s$  the minimum number of vertices forming an  $AB$ -separator. Then there is an  $AB$ -connector  $C$  with  $|C \cap A| = s$ .*

**Proof.** If  $G$  is edgeless then set  $C = A \cap B$ . Suppose,  $G$  is a counterexample with  $|E(G)|$  minimal. Then  $G$  contains an edge  $e$  from  $x$  to  $y$  and  $G/e$  has an  $AB$ -separator  $S$  with  $|S| < s$ , otherwise we are done. Obviously,  $v_e \in S$ . Then  $P = (S \setminus \{v_e\}) \cup \{x, y\}$  is an  $AB$ -separator of  $G$  with  $|P| = |S| + 1 = s$ . An  $AP$ -separator (as well as an  $PB$ -separator) of  $G - e$  is an  $AB$ -separator of  $G$ . Consequently,  $G - e$  has an  $AP$ -connector  $X$  and a  $PB$ -connector  $Y$  containing  $P$ . Since  $X \cap Y = P$ , the set  $C = (X \cup Y)$  is an  $AB$ -connector of  $G$  with  $|C \cap A| = s$ , a contradiction. ■

## References

- [1] T. Böhme, F. Göring and J. Harant, *Menger's Theorem*, J. Graph Theory **37** (2001) 35–36.
- [2] W. McCuaig, *A simple proof of Menger's theorem*, J. Graph Theory **8** (1984) 427–429.
- [3] R. Diestel, *Graph Theory* (2nd edition), (Springer-Verlag, New York, 2000).
- [4] G.A. Dirac, *Short proof of Menger's graph theorem*, Mathematika **13** (1966) 42–44.
- [5] F. Goering, *Short Proof of Menger's Theorem*, to appear in Discrete Math.
- [6] T. Grünwald (later Gallai), *Ein neuer Beweis eines Mengerschen Satzes*, J. London Math. Soc. **13** (1938) 188–192.
- [7] K. Menger, *Zur allgemeinen Kurventheorie*, Fund. Math. **10** (1927) 96–115.
- [8] J.S. Pym, *A proof of Menger's theorem*, Monatshefte Math. **73** (1969) 81–88.

Received 8 June 2000

Revised 21 May 2001