

TWO GRAPHS WITH A COMMON EDGE

LIDIA BADURA

*Mathematics and Computer Science Department
University of Opole
Oleska 48
45-052 Opole, Poland*

e-mail: badura@math.uni.opole.pl

Abstract

Let $G = G_1 \cup G_2$ be the sum of two simple graphs G_1, G_2 having a common edge or $G = G_1 \cup e_1 \cup e_2 \cup G_2$ be the sum of two simple disjoint graphs G_1, G_2 connected by two edges e_1 and e_2 which form a cycle C_4 inside G . We give a method of computing the determinant $\det A(G)$ of the adjacency matrix of G by reducing the calculation of the determinant to certain subgraphs of G_1 and G_2 . To show the scope and effectiveness of our method we give some examples.

Keywords: graph, adjacency matrix, determinant of graph, path, cycle.

2010 Mathematics Subject Classification: 05C50.

REFERENCES

- [1] A. Abdollahi, *Determinants of adjacency matrices of graphs*, Trans. Combin. **1**(4) (2012) 9–16.
- [2] F. Harary, *The Determinant of the adjacency matrix of a graph*, SIAM Rev. **4** (1961) 202–210.
doi:10.1137/1004057
- [3] L. Huang and W. Yan, *On the determinant of the adjacency matrix of a type of plane bipartite graphs*, MATCH Commun. Math. Comput. Chem. **68** (2012) 931–938.
- [4] H.M. Rara, *Reduction procedures for calculating the determinant of the adjacency matrix of some graphs and the singularity of square planar grids*, Discrete Math. **151** (1996) 213–219.
doi:10.1016/0012-365X(94)00098-4
- [5] P. Wojtylak and S. Arworn, *Paths of cycles and cycles of cycles*, (2010) preprint.

Received 30 October 2012

Revised 8 May 2013

Accepted 8 May 2013