

## SHARP UPPER BOUNDS ON THE SIGNLESS LAPLACIAN SPECTRAL RADIUS OF STRONGLY CONNECTED DIGRAPHS<sup>1</sup>

WEIGE XI AND LIGONG WANG<sup>2</sup>

*Department of Applied Mathematics  
School of Science, Northwestern Polytechnical University  
Xi'an, Shaanxi 710072, P.R. China*

**e-mail:** xiyanxwg@163.com  
lgwangmath@163.com

### Abstract

Let  $G = (V(G), E(G))$  be a simple strongly connected digraph and  $q(G)$  be the signless Laplacian spectral radius of  $G$ . For any vertex  $v_i \in V(G)$ , let  $d_i^+$  denote the outdegree of  $v_i$ ,  $m_i^+$  denote the average 2-outdegree of  $v_i$ , and  $N_i^+$  denote the set of out-neighbors of  $v_i$ . In this paper, we prove that:

(1)  $q(G) = d_1^+ + d_2^+$ , ( $d_1^+ \neq d_2^+$ ) if and only if  $G$  is a star digraph  $\overleftrightarrow{K}_{1,n-1}$ , where  $d_1^+, d_2^+$  are the maximum and the second maximum outdegree, respectively ( $\overleftrightarrow{K}_{1,n-1}$  is the digraph on  $n$  vertices obtained from a star graph  $K_{1,n-1}$  by replacing each edge with a pair of oppositely directed arcs).

(2)  $q(G) \leq \max \left\{ \frac{1}{2} \left( d_i^+ + \sqrt{d_i^{+2} + 8d_i^+ m_i^+} \right) : v_i \in V(G) \right\}$  with equality if and only if  $G$  is a regular digraph.

(3)  $q(G) \leq \max \left\{ \frac{1}{2} \left( d_i^+ + \sqrt{d_i^{+2} + \frac{4}{d_i^+} \sum_{v_j \in N_i^+} d_j^+ (d_j^+ + m_j^+)} \right) : v_i \in V(G) \right\}$ .

Moreover, the equality holds if and only if  $G$  is a regular digraph or a bipartite semiregular digraph.

(4)  $q(G) \leq \max \left\{ \frac{1}{2} \left( d_i^+ + 2d_j^+ - 1 + \sqrt{(d_i^+ - 2d_j^+ + 1)^2 + 4d_i^+} \right) : (v_j, v_i) \in E(G) \right\}$ . If the equality holds, then  $G$  is a regular digraph or  $G \in \Omega$ , where  $\Omega$  is a class of digraphs defined in this paper.

**Keywords:** digraph, signless Laplacian spectral radius.

**2010 Mathematics Subject Classification:** 05C50, 15A18.

<sup>1</sup>The research was supported by the National Natural Science Foundation of China (Grant No. 11171273).

<sup>2</sup>Corresponding author.

## REFERENCES

- [1] Ş.B. Bozkurt and D. Bozkurt, *On the signless Laplacian spectral radius of digraphs*, *Ars Combin.* **108** (2013) 193–200.
- [2] B.K. Bulter and P.H. Siegel, *Sharp bounds on the spectral radius of nonnegative matrices and digraphs*, *Linear Algebra Appl.* **439** (2013) 1468–1478.  
doi:10.1016/j.laa.2013.04.029
- [3] Y.Q. Chen and L.G. Wang, *Sharp bounds for the largest eigenvalue of the signless Laplacian of a graph*, *Linear Algebra Appl.* **433** (2010) 908–913.  
doi:10.1016/j.laa.2010.04.026
- [4] S.Y. Cui, C.X. Tian and J.J. Guo, *A sharp upper bound on the signless Laplacian spectral radius of graphs*, *Linear Algebra Appl.* **439** (2013) 2442–2447.  
doi:10.1016/j.laa.2013.06.015
- [5] K.Ch. Das, *A characterization on graphs which achieve the upper bound for the largest Laplacian eigenvalue of graphs*, *Linear Algebra Appl.* **376** (2004) 173–186.  
doi:10.1016/j.laa.2003.06.009
- [6] K.Ch. Das, *The Laplacian spectrum of a graph*, *Comput. Math. Appl.* **48** (2004) 715–724.  
doi:10.1016/j.camwa.2004.05.005
- [7] K.Ch. Das, *Maximizing the sum of the squares of the degrees of a graph*, *Discrete Math.* **285** (2004) 57–66.  
doi:10.1016/j.disc.2004.04.007
- [8] E. Gudiño and J. Rada, *A lower bound for the spectral radius of a digraph*, *Linear Algebra Appl.* **433** (2010) 233–240.  
doi:10.1016/j.laa.2010.02.012
- [9] A.D. Güngör, K.C. Das and A.S. Çevik, *Sharp upper bounds on the spectral radius of the signless Laplacian matrix of a graph*, *Appl. Math. Comput.* **219** (2013) 5025–5032.  
doi:10.1016/j.amc.2012.11.039
- [10] W.X. Hong and L.H. You, *Spectral radius and signless Laplacian spectral radius of strongly connected digraphs*, *Linear Algebra Appl.* **457** (2014) 93–113.  
doi:10.1016/j.laa.2014.05.007
- [11] R.A. Horn and C.R. Johnson, *Matrix Analysis* (Cambridge University Press, New York, 1985).  
doi:10.1017/CBO9780511810817
- [12] W.W. Lang and L.G. Wang, *Sharp bounds for the signless Laplacian spectral radius of digraphs*, *Appl. Math. Comput.* **238** (2014) 43–49.  
doi:10.1016/j.amc.2014.04.001
- [13] M.H. Liu, B.L. Liu and B. Cheng, *Ordering (signless) Laplacian spectral radii with maximum degrees of graphs*, *Discrete Math.* **338** (2015) 159–163.  
doi:10.1016/j.disc.2014.09.007

- [14] C.S. Oliveira, L.S. de Lima, N.M.M. de Abreu and P. Hansen, *Bounds on the index of the signless Laplacian of a graph*, Discrete Appl. Math. **158** (2010) 355–360.  
doi:10.1016/j.dam.2009.06.023
- [15] Y.L. Pan, *Sharp upper bounds for the Laplacian graph eigenvalues*, Linear Algebra Appl. **355** (2002) 287–295.  
doi:10.1016/S0024-3795(02)00353-1
- [16] G.H. Xu, K.F. Fang and J. Shen, *Bounds on the spectral radii of digraphs in terms of walks*, Appl. Math. Comput. **219** (2012) 3721–3728.  
doi:10.1016/j.amc.2012.09.073
- [17] G.L. Yu, Y.R. Wu and J.L. Shu, *Sharp bounds on the signless Laplacian spectral radii of graphs*, Linear Algebra Appl. **434** (2011) 683–687.  
doi:10.1016/j.laa.2010.09.029

Received 26 March 2015  
Revised 13 January 2016  
Accepted 13 January 2016