

## ONE-THREE JOIN: A GRAPH OPERATION AND ITS CONSEQUENCES

M.A. SHALU AND S. DEVI YAMINI

IITD & M, Kancheepuram, Chennai-600127, India

**e-mail:** shalu@iiitdm.ac.in  
mat10d001@iiitdm.ac.in

### Abstract

In this paper, we introduce a graph operation, namely *one-three join*. We show that the graph  $G$  admits a one-three join if and only if either  $G$  is one of the basic graphs (bipartite, complement of bipartite, split graph) or  $G$  admits a constrained homogeneous set or a bipartite-join or a join. Next, we define  $\mathcal{M}_H$  as the class of all graphs generated from the induced subgraphs of an odd hole-free graph  $H$  that contains an odd anti-hole as an induced subgraph by using one-three join and co-join recursively and show that the maximum independent set problem, the maximum clique problem, the minimum coloring problem, and the minimum clique cover problem can be solved efficiently for  $\mathcal{M}_H$ .

**Keywords:** one-three join, bipartite-join, homogeneous set, odd hole-free graphs.

**2010 Mathematics Subject Classification:** 05C75, 05C76.

### REFERENCES

- [1] R.K. Ahuja, T.L. Magnanti and J.B. Orlin, Network Flows (Prentice-Hall, Englewood Cliffs, NJ, 1993) 409–411.
- [2] M. Basavaraju, L.S. Chandran and T. Karthick, *Maximum weight independent sets in hole- and dart-free graphs*, Discrete Appl. Math. **160** (2012) 2364–2369. doi:10.1016/j.dam.2012.06.015
- [3] D. Bienstock, *On complexity of testing for odd holes and induced odd paths*, Discrete Math. **90** (1991) 85–92. doi:10.1016/0012-365X(91)90098-M
- [4] H.L. Bodlaender, A. Brandstädt, D. Kratsch, M. Rao and J. Spinrad, *On algorithms for  $\{P_5, gem\}$ -free graphs*, Theoret. Comput. Sci. **349** (2005) 2–21. doi:10.1016/j.tcs.2005.09.026

- [5] A. Brandstädt and V. Giakoumakis, *Addendum to: Maximum weighted independent sets in hole- and co-chair-free graphs*, Inform. Process. Lett. **115** (2015) 345–350.  
doi:10.1016/j.ipl.2014.09.019
- [6] A. Brandstädt, V. Giakoumakis and F. Maffray, *Clique separator decomposition of hole- and diamond-free graphs and algorithmic consequences*, Discrete Appl. Math. **160** (2012) 471–478.  
doi:10.1016/j.dam.2011.10.031
- [7] A. Brandstädt and T. Karthick, *Weighted efficient domination in two subclasses of  $P_6$ -free graphs*, Discrete Appl. Math. **201** (2016) 38–46.  
doi:10.1016/j.dam.2015.07.032
- [8] A. Brandstädt and R. Mosca, *Maximum weight independent sets in odd hole-free graphs without dart or without bull*, Graphs Combin. **31** (2015) 1249–1262.  
doi:10.1007/s00373-014-1461-x
- [9] M. Chudnovsky and P. Seymour, *The structure of claw-free graphs*, Surveys in Combinatorics, London Math. Soc. Lecture Note Ser. **327** (2005) 153–171.
- [10] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, *The strong perfect graph theorem*, Ann. of Math. **164** (2006) 51–229.  
doi:10.4007/annals.2006.164.51
- [11] M. Chudnovsky, I. Penev, A. Scott and N. Trotignon, *Substitution and  $\chi$ -boundedness*, J. Combin. Theory Ser. B **103** (2013) 567–586.  
doi:10.1016/j.jctb.2013.02.004
- [12] M. Conforti, G. Cornuéjols and K. Vušković, *Even-hole-free graphs, Part II: Recognition algorithm*, J. Graph Theory **40** (2002) 238–266.  
doi:10.1002/jgt.10045
- [13] M. Conforti, G. Cornuéjols and K. Vušković, *Decomposition of odd-hole-free graphs by double star cutsets and 2-joins*, Discrete Appl. Math. **141** (2004) 41–91.  
doi:10.1016/S0166-218X(03)00364-0
- [14] M. Conforti, G. Cornuéjols, X. Liu and K. Vušković and G. Zambelli, *Odd-hole recognition in graphs of bounded clique size*, SIAM J. Discrete Math. **20** (2006) 42–48.  
doi:10.1137/S089548010444540X
- [15] D.G. Corneil, H. Lerchs and L.S. Burlingham, *Complement reducible graphs*, Discrete Appl. Math. **3** (1981) 163–174.  
doi:10.1016/0166-218X(81)90013-5
- [16] T. Feder, P. Hell, S. Klein and R. Motwani, *Complexity of graph partition problems*, in: Proceedings of the Thirty-First Annual ACM Symposium on Theory of Computing (1999) 464–472.  
doi:10.1145/301250.301373
- [17] T. Feder, P. Hell, S. Klein and R. Motwani, *List partitions*, SIAM J. Discrete Math. **16** (2003) 449–478.  
doi:10.1137/S0895480100384055

- [18] S. Földes and P.L. Hammer, *Split graphs*, Congr. Numer. **19** (1977) 311–315.
- [19] M. Grötschel, L. Lovász and A. Schrijver, *Polynomial algorithms for perfect graphs*, Ann. Discrete Math. **21** (1984) 325–356.  
doi:10.1016/s0304-0208(08)72943-8
- [20] N.C. Lê, C. Brause and I. Schiermeyer, *Extending the MAX Algorithm for maximum independent set*, Discuss. Math. Graph Theory **35** (2015) 365–386.  
doi:10.7151/dmgt.1811
- [21] R. Mosca, *Stable sets for  $(P_6, K_{2,3})$ -free graphs*, Discuss. Math. Graph Theory **32** (2012) 387–401.  
doi:10.7151/dmgt.1598
- [22] D.B. West, *Introduction to Graph Theory* (Prentice Hall, USA, 1996).

Received 29 September 2015

Revised 2 June 2016

Accepted 2 June 2016