Discussiones Mathematicae
Graph Theory 39 (2019) 31–40
doi:10.7151/dmgt.2063

BOUNDING THE LOCATING-TOTAL DOMINATION NUMBER OF A TREE IN TERMS OF ITS ANNIHILATION NUMBER

Wenjie Ning

College of Science
China University of Petroleum (East China)
Qingdao 266580, China

e-mail: ningwenjie-0501@163.com

Mei Lu

Department of Mathematical Sciences
Tsinghua University, Beijing 100084, China

e-mail: mlu@math.tsinghua.edu.cn

and

Kun Wang

School of Mathematical Sciences
Anhui University, Hefei 230601, China

e-mail: wangkun26@163.com

Abstract

Suppose $G = (V, E)$ is a graph with no isolated vertex. A subset S of V is called a locating-total dominating set of G if every vertex in V is adjacent to a vertex in S, and for every pair of distinct vertices u and v in $V - S$, we have $N(u) \cap S \neq N(v) \cap S$. The locating-total domination number of G, denoted by $\gamma_t^L(G)$, is the minimum cardinality of a locating-total dominating set of G. The annihilation number of G, denoted by $a(G)$, is the largest integer k such that the sum of the first k terms of the nondecreasing degree sequence of G is at most the number of edges in G. In this paper, we show that for any tree of order $n \geq 2$, $\gamma_t^L(T) \leq a(T) + 1$ and we characterize the trees achieving this bound.

Keywords: total domination, locating-total domination, annihilation number, tree.

2010 Mathematics Subject Classification: 05C69.

References

[1] M. Blidia and W. Dali, *A characterization of locating-total domination edge critical graphs*, Discuss. Math. Graph Theory **31** (2011) 197–202.
doi:10.7151/dmgt.1538

[2] M. Chellali, *On locating and differentiating-total domination in trees*, Discuss. Math. Graph Theory **28** (2008) 383–392.
doi:10.7151/dmgt.1414

[3] M. Chellali and N. Jafari Rad, *Locating-total domination critical graphs*, Australas. J. Combin. **45** (2009) 227–234.

[4] X. Chen and M.Y. Sohn, *Bounds on the locating-total domination number of a tree*, Discrete Appl. Math. **159** (2011) 769–773.
doi:10.1016/j.dam.2010.12.025

[5] W.J. Desormeaux, T.W. Haynes and M.A. Henning, *Relating the annihilation number and the total domination number of a tree*, Discrete Appl. Math. **161** (2013) 349–354.
doi:10.1016/j.dam.2012.09.006

[6] W.J. Desormeaux, M.A. Henning, D.F. Rall and A. Yeos, *Relating the annihilation number and the* 2*-domination number of a tree*, Discrete Math. **319** (2014) 15–23.
doi:10.1016/j.disc.2013.11.020

[7] O. Favaron, M.A. Henning, J. Puecha and D. Rautenbach, *On domination and annihilation in graphs with claw-free blocks*, Discrete Math. **231** (2001) 143–151.
doi:10.1016/S0012-365X(00)00313-7

[8] J.R. Griggs and D.J. Kleitman, *Independence and the Havel-Hakimi residue*, Discrete Math. **127** (1994) 209–212.
doi:10.1016/0012-365X(92)00479-B

[9] T.W. Haynes, M.A. Henning and J. Howard, *Locating and total dominating sets in trees*, Discrete Appl. Math. **154** (2006) 1293–1300.
doi:10.1016/j.dam.2006.01.002

[10] M.A. Henning and N. Jafari Rad, *Locating-total domination in graphs*, Discrete Appl. Math. **160** (2012) 1986–1993.
doi:10.1016/j.dam.2012.04.004

[11] L. Jennings, New Sufficient Condition for Hamiltonian Paths (Ph.D. Dissertation, Rice University, 2008).

[12] C.E. Larson and R. Pepper, *Graphs with equal independence and annihilation numbers*, Electron. J. Combin. **18** (2011) #P180.

[13] R. Pepper, Binding Independence (Ph.D. Dissertation, University of Houston, 2004).

[14] R. Pepper, *On the annihilation number of a graph*, in: Recent Advances in Applied Mathematics and Computational and Information Sciences, Vol. I, K. Jegdic, P. Simeonov, V. Zafiris (Ed(s)), (WSEAS Press, 2009) 217–220.

Received 28 November 2016
Accepted 6 May 2017