

ALL TIGHT DESCRIPTIONS OF 3-STARS IN 3-POLYTOPES WITH GIRTH 5

OLEG V. BORODIN¹

*Institute of Mathematics Siberian Branch
Russian Academy of Sciences
Novosibirsk, 630090, Russia*

e-mail: brdnoleg@math.nsc.ru

AND

ANNA O. IVANOVA²

*Ammosov North-Eastern Federal University
Yakutsk, 677000, Russia*

e-mail: shmgnanna@mail.ru

Abstract

Lebesgue (1940) proved that every 3-polytope P_5 of girth 5 has a path of three vertices of degree 3. Madaras (2004) refined this by showing that every P_5 has a 3-vertex with two 3-neighbors and the third neighbor of degree at most 4. This description of 3-stars in P_5 s is tight in the sense that no its parameter can be strengthened due to the dodecahedron combined with the existence of a P_5 in which every 3-vertex has a 4-neighbor.

We give another tight description of 3-stars in P_5 s: there is a vertex of degree at most 4 having three 3-neighbors. Furthermore, we show that there are only these two tight descriptions of 3-stars in P_5 s.

Also, we give a tight description of stars with at least three rays in P_5 s and pose a problem of describing all such descriptions. Finally, we prove a structural theorem about P_5 s that might be useful in further research.

Keywords: 3-polytope, planar graph, structure properties, k -star.

2010 Mathematics Subject Classification: 05C15.

¹This work was supported by grants 16-01-00499 and 15-01-05867 of the Russian Foundation for Basic Research and President Grant for Government Support of the Leading Scientific Schools of the Russian Federation NSh-1939.2014.1.

²The author's work was performed as a part of government work "Organizing research" and supported by grant 15-01-05867 of the Russian Foundation for Basic Research.

REFERENCES

- [1] O.V. Borodin, *Solution of Kotzig's and Grünbaum's problems on the separability of a cycle in a planar graph*, Mat. Zametki **46** (1989) 9–12, in Russian.
- [2] O.V. Borodin, *Structural properties of planar maps with the minimal degree 5*, Math. Nachr. **158** (1992) 109–117.
doi:10.1002/mana.19921580108
- [3] O.V. Borodin and D. Sanders, *On light edges and triangles in planar graphs of minimal degree five*, Math. Nachr. **170** (1994) 19–24.
doi:10.1002/mana.19941700103
- [4] O.V. Borodin and D.R. Woodall, *Short cycles of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **18** (1998) 159–164.
doi:0.7151/dmgt.1071
- [5] O.V. Borodin and A.O. Ivanova, *Describing 4-stars at 5-vertices in normal plane maps with minimum degree 5*, Discrete Math. **313** (2013) 1710–1714.
doi:10.1016/j.disc.2013.04.025
- [6] O.V. Borodin, A.O. Ivanova and A.V. Kostochka, *Every 3-polytope with minimum degree 5 has a 6-cycle with maximum degree at most 11*, Discrete Math. **315–316** (2014) 128–134.
doi:10.1016/j.disc.2013.10.021
- [7] O.V. Borodin, A.O. Ivanova and D.R. Woodall, *Light C_4 and C_5 in 3-polytopes with minimum degree 5*, Discrete Math. **334** (2014) 63–69.
doi:10.1016/j.disc.2014.06.024
- [8] O.V. Borodin, A.O. Ivanova and T.R. Jensen, *5-stars of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **34** (2014) 539–546.
doi:10.7151/dmgt.1748
- [9] O.V. Borodin and A.O. Ivanova, *Every 3-polytope with minimum degree 5 has a 7-cycle with maximum degree at most 15*, Sibirsk. Mat. Zh. **56** (2015) 775–789, in Russian.
- [10] B. Ferencová and T. Madaras, *On the structure of polyhedral graphs with prescribed edge and dual edge weight*, Acta Univ. M. Belii Ser. Math. **12** (2005) 13–18.
- [11] B. Ferencová and T. Madaras, *Light graph in families of polyhedral graphs with prescribed minimum degree, face size, edge and dual edge weight*, Discrete Math. **310** (2010) 1661–1675.
doi:10.1016/j.disc.2009.11.027
- [12] Ph. Franklin, *The four-color problem*, Amer. J. Math. **44** (1922) 225–236.
doi:10.2307/2370527
- [13] P. Hudák and T. Madaras, *On doubly light triangles in plane graphs*, Discrete Math. **313** (2013) 1978–1988.
doi:10.1016/j.disc.2012.11.018

- [14] S. Jendrol' and T. Madaras, *On light subgraphs in plane graphs with minimum degree five*, Discuss. Math. Graph Theory **16** (1996) 207–217.
doi:10.7151/dmgt.1035
- [15] S. Jendrol', T. Madaras, R. Soták and Zs. Tuza, *On light cycles in plane triangulations*, Discrete Math. **197–198** (1999) 453–467.
doi:10.1016/S0012-365X(98)00254-4
- [16] S. Jendrol' and H.-J. Voss, *Light subgraphs of graphs embedded in the plane—a survey*, Discrete Math. **313** (2013) 406–421.
doi:10.1016/j.disc.2012.11.007
- [17] H. Lebesgue, *Quelques conséquences simples de la formule d'Euler*, J. Math. Pures Appl. **19** (1940) 27–43.
- [18] T. Madaras and R. Soták, *The 10-cycle C_{10} is light in the family of all plane triangulations with minimum degree five*, Tatra Mt. Math. Publ. **18** (1999) 35–56.
- [19] T. Madaras, *On the structure of plane graphs of minimum face size 5*, Discuss. Math. Graph Theory **24** (2004) 403–411.
doi:10.7151/dmgt.1239
- [20] T. Madaras, *Two variations of Franklin's theorem*, Tatra Mt. Math. Publ. **36** (2007) 61–70.
- [21] T. Madaras, R. Škrekovski and H.-J. Voss, *The 7-cycle C_7 is light in the family of planar graphs with minimum degree 5*, Discrete Math. **307** (2007) 1430–1435.
doi:10.1016/j.disc.2005.11.080
- [22] E. Steinitz, *Polyeder und Raumeinteilungen*, Enzykl. Math. Wiss. **3** (1922) 1–139.
- [23] P. Wernicke, *Über den kartographischen Vierfarbensatz*, Math. Ann. **58** (1904) 413–426.
doi:10.1007/BF01444968

Received 22 June 2015
 Revised 16 January 2016
 Accepted 16 January 2016