

NEIGHBOR SUM DISTINGUISHING TOTAL CHOOSABILITY OF IC-PLANAR GRAPHS

WEN-YAO SONG, LIAN-YING MIAO

School of Mathematics
China University of Mining and Technology
Xuzhou 221116, P.R. China

e-mail: songwenyao@cumt.edu.cn
miaolianying@cumt.edu.cn

AND

YUAN-YUAN DUAN

School of Mathematics and Statistics
Zaozhuang University
Zaozhuang 277160, P.R. China

e-mail: duanyy0827@sina.com

Abstract

Two distinct crossings are independent if the end-vertices of the crossed pair of edges are mutually different. If a graph G has a drawing in the plane such that every two crossings are independent, then we call G a plane graph with independent crossings or IC-planar graph for short. A proper total- k -coloring of a graph G is a mapping $c : V(G) \cup E(G) \rightarrow \{1, 2, \dots, k\}$ such that any two adjacent elements in $V(G) \cup E(G)$ receive different colors. Let $\sum_c(v)$ denote the sum of the color of a vertex v and the colors of all incident edges of v . A total- k -neighbor sum distinguishing-coloring of G is a total- k -coloring of G such that for each edge $uv \in E(G)$, $\sum_c(u) \neq \sum_c(v)$. The least number k needed for such a coloring of G is the neighbor sum distinguishing total chromatic number, denoted by $\chi''_{\Sigma}(G)$. In this paper, it is proved that if G is an IC-planar graph with maximum degree $\Delta(G)$, then $ch''_{\Sigma}(G) \leq \max\{\Delta(G) + 3, 17\}$, where $ch''_{\Sigma}(G)$ is the neighbor sum distinguishing total choosability of G .

Keywords: neighbor sum distinguishing total choosability, maximum degree, IC-planar graph, Combinatorial Nullstellensatz.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] M.O. Albertson, *Chromatic number, independence ratio, and crossing number*, Ars Math. Contemp. **1** (2008) 1–6.
- [2] N. Alon, *Combinatorial Nullstellensatz*, Combin. Probab. Comput. **8** (1999) 7–29. doi:10.1017/S0963548398003411
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications* (North-Holland, New York-Amsterdam-Oxford, 1982).
- [4] L. Ding, G. Wang and G. Yan, *Neighbor sum distinguishing total colorings via the Combinatorial Nullstellensatz*, Sci. China Math. **57** (2014) 1875–1882. doi:10.1007/s11425-014-4796-0
- [5] L. Ding, G. Wang, J. Wu and J. Yu, *Neighbor sum (set) distinguishing total choosability via the Combinatorial Nullstellensatz*, Graphs Combin. **33** (2017) 885–900. doi:10.1007/s00373-017-1806-3
- [6] A. Dong and G. Wang, *Neighbor sum distinguishing total colorings of graphs with bounded maximum average degree*, Acta Math. Sin. (Engl. Ser.) **30** (2014) 703–709. doi:10.1007/s10114-014-2454-7
- [7] D. Král and L. Stacho, *Coloring plane graphs with independent crossings*, J. Graph Theory **64** (2010) 184–205. doi:10.1002/jgt.20448
- [8] H. Li, B. Liu and G. Wang, *Neighbor sum distinguishing total colorings of K_4 -minor free graphs*, Front. Math. China **8** (2013) 1351–1366. doi:10.1007/s11464-013-0322-x
- [9] H. Li, L. Ding, B. Liu and G. Wang, *Neighbor sum distinguishing total colorings of planar graphs*, J. Comb. Optim. **30** (2015) 675–688. doi:10.1007/s10878-013-9660-6
- [10] S. Loeb, J. Przybyło and Y. Tang, *Asymptotically optimal neighbor sum distinguishing total colorings of graphs*, Discrete Math. **340** (2017) 58–62. doi:10.1016/j.disc.2016.08.012
- [11] M. Pilśniak and M. Woźniak, *On the total-neighbor-distinguishing index by sums*, Graphs Combin. **31** (2015) 771–782. doi:10.1007/s00373-013-1399-4
- [12] C. Qu, G. Wang, J. Wu and X. Yu, *On the neighbor sum distinguishing total coloring of planar graphs*, Theoret. Comput. Sci. **609** (2016) 162–170. doi:10.1016/j.tcs.2015.09.017
- [13] C. Qu, G. Wang, G. Yan and X. Yu, *Neighbor sum distinguishing total choosability of planar graphs*, J. Comb. Optim. **32** (2016) 906–916. doi:10.1007/s10878-015-9911-9
- [14] D.E. Scheim, *The number of edge 3-colorings of a planar cubic graph as a permanent*, Discrete Math. **8** (1974) 377–382. doi:10.1016/0012-365X(74)90157-5

- [15] J. Wang, J. Cai and Q. Ma, *Neighbor sum distinguishing total choosability of planar graphs without 4-cycles*, Discrete Appl. Math. **206** (2016) 215–219.
doi:10.1016/j.dam.2016.02.003
- [16] J. Yao and H. Kong, *Neighbor sum distinguishing total choosability of graphs with larger maximum average degree*, Ars Combin. **125** (2016) 347–360.
- [17] X. Zhang and J. Wu, *On edge colorings of 1-planar graphs*, Inform. Process. Lett. **111** (2011) 124–128.
doi:10.1016/j.ipl.2010.11.001
- [18] J. Yao, X. Yu, G. Wang and C. Xu, *Neighbor sum (set) distinguishing total choosability of d -degenerate graphs*, Graphs Combin. **32** (2016) 1611–1620.
doi:10.1007/s00373-015-1646-y
- [19] X. Cheng, D. Huang, G. Wang and J. Wu, *Neighbor sum distinguishing total colorings of planar graphs with maximum degree Δ* , Discrete Appl. Math. **190-191** (2015) 34–41.
doi:10.1016/j.dam.2015.03.013
- [20] Y. Lu, M.M. Han and R. Luo, *Neighbor sum distinguishing total coloring and list neighbor sum distinguishing total coloring*, Discrete Appl. Math. **237** (2018) 109–115.
doi:10.1016/j.dam.2017.12.001

Received 30 June 2017
Revised 21 March 2018
Accepted 21 March 2018