

ON LIST EQUITABLE TOTAL COLORINGS OF THE GENERALIZED THETA GRAPH

JEFFREY A. MUDROCK, MAX MARSH

AND

TIM WAGSTROM

Department of Mathematics
College of Lake County
Grayslake, IL 60030

e-mail: jumudrock@clcollinois.edu

Abstract

In 2003, Kostochka, Pelsmajer, and West introduced a list analogue of equitable coloring called equitable choosability. A k -assignment, L , for a graph G assigns a list, $L(v)$, of k available colors to each $v \in V(G)$, and an equitable L -coloring of G is a proper coloring, f , of G such that $f(v) \in L(v)$ for each $v \in V(G)$ and each color class of f has size at most $\lceil |V(G)|/k \rceil$. Graph G is equitably k -choosable if G is equitably L -colorable whenever L is a k -assignment for G . In 2018, Kaul, Mudrock, and Pelsmajer subsequently introduced the List Equitable Total Coloring Conjecture which states that if T is a total graph of some simple graph, then T is equitably k -choosable for each $k \geq \max\{\chi_\ell(T), \Delta(T)/2 + 2\}$ where $\Delta(T)$ is the maximum degree of a vertex in T and $\chi_\ell(T)$ is the list chromatic number of T . In this paper, we verify the List Equitable Total Coloring Conjecture for subdivisions of stars and the generalized theta graph.

Keywords: graph coloring, total coloring, equitable coloring, list coloring, equitable choosability.

2010 Mathematics Subject Classification: 05C15.

1. INTRODUCTION

In this paper, all graphs are nonempty, finite, simple graphs unless otherwise noted. Generally speaking we follow West [37] for terminology and notation. The set of natural numbers is $\mathbb{N} = \{1, 2, 3, \dots\}$. For $m \in \mathbb{N}$, we write $[m]$ for the

set $\{1, \dots, m\}$. For sets A and B , we write $A - B$ for the set of all elements of A that are not elements of B . If G is a graph and $S \subseteq V(G)$, we use $G[S]$ for the subgraph of G induced by S . For $v \in V(G)$, we write $d_G(v)$ for the degree of vertex v in the graph G and $\Delta(G)$ for the maximum degree of a vertex in G , and we write $N_G(v)$ for the neighborhood of vertex v in the graph G . If $e, w \in E(G)$ and $v \in V(G)$, we say e and v are incident if v is an endpoint of e , and we say e and w are adjacent if e and w share an endpoint. Also, G^k denotes the k^{th} power of graph G (i.e. G^k has the same vertex set as G and edges between any two vertices within distance k in G).

1.1. Total coloring, equitable coloring, and list coloring

In this paper, we study a conjecture that combines different types of colorings, namely total coloring, equitable coloring, and list coloring. So, we begin by briefly reviewing these three notions.

Given a graph G , in the classic vertex coloring problem we wish to color the elements of $V(G)$ with colors from the set $[m]$ so that adjacent vertices receive different colors, a so-called *proper m -coloring*. The *chromatic number* of G , denoted $\chi(G)$, is the smallest k such that a proper k -coloring of G exists.

1.1.1. Total coloring

A *total m -coloring* of G is a labeling $f : V(G) \cup E(G) \rightarrow [m]$ where $f(u) \neq f(v)$ whenever u and v are adjacent or incident in G . The *total chromatic number* of a graph G , denoted $\chi''(G)$, is the smallest k such that G has a total k -coloring. Clearly, for any graph G , $\chi''(G) \geq \Delta(G) + 1$. A famous open problem is the Total Coloring Conjecture (see [1, 34]) which states that for any graph G , $\chi''(G) \leq \Delta(G) + 2$. See [22] for some other applications of total coloring.

It is possible to rephrase total coloring in terms of classic vertex coloring. Specifically, the *total graph* of graph G , $T(G)$, is the graph with vertex set $V(G) \cup E(G)$ and vertices are adjacent in $T(G)$ if and only if the corresponding elements are adjacent or incident in G . Then G has a total k -coloring if and only if $T(G)$ has a proper k -coloring. It follows that $\chi''(G) = \chi(T(G))$.

Given a graph G , one can construct $T(G)$ in two steps: first subdivide every edge of G to get a new graph H , then take its square (i.e. $T(G) = H^2$). For example, $T(P_m) = P_{2m-1}^2$.

1.1.2. Equitable coloring

The study of equitable coloring began in 1964 with a conjecture of Erdős [9], but it was formally introduced by Meyer in the 1970's [27]. An *equitable k -coloring* of a graph G is a proper k -coloring of G , f , such that the sizes of the color classes differ by at most one (where a proper k -coloring has exactly k color classes). In

an equitable k -coloring, the color classes associated with the coloring are each of size $\lceil |V(G)|/k \rceil$ or $\lfloor |V(G)|/k \rfloor$. We say that a graph G is *equitably k -colorable* if there exists an equitable k -coloring of G . Many applications of equitable coloring exist, see for example [16, 17, 31, 33].

Unlike classic vertex coloring, increasing the number of colors can make equitable coloring more difficult in certain cases. Indeed for any $m \in \mathbb{N}$, $K_{2m+1, 2m+1}$ is equitably $2m$ -colorable, but it is not equitably $(2m + 1)$ -colorable. In 1970, Hajnal and Szemeredi [15] proved the 1964 conjecture of Erdős: every graph G has an equitable k -coloring when $k \geq \Delta(G) + 1$.

In 1994, Chen, Lih, and Wu [5] conjectured that the result of Hajnal and Szemeredi can be improved by 1 for most connected graphs. Their conjecture is known as the Δ -Equitable Coloring Conjecture, and it is still open. Formally, the Δ -Equitable Coloring Conjecture states: a connected graph G is equitably $\Delta(G)$ -colorable if it is different from K_m , C_{2m+1} , and $K_{2m+1, 2m+1}$. The Δ -Equitable Coloring Conjecture has been proven true for interval graphs, bipartite graphs, outerplanar graphs, subcubic graphs, certain planar graphs, and several other classes of graphs (see [5, 7, 8, 25, 26, 38]).

1.1.3. List coloring

List coloring is another variation on classic vertex coloring that was introduced independently by Vizing [35] and Erdős, Rubin, and Taylor [10] in the 1970's. In list coloring, we associate with graph G a *list assignment*, L , that assigns to each vertex $v \in V(G)$ a list, $L(v)$, of available colors. Graph G is said to be *L -colorable* if there exists a proper coloring f of G such that $f(v) \in L(v)$ for each $v \in V(G)$ (we refer to f as a *proper L -coloring* of G). A list assignment L is called a *k -assignment* for G if $|L(v)| = k$ for each $v \in V(G)$. We say G is *k -choosable* if G is L -colorable whenever L is a k -assignment for G . The *list chromatic number* of G , denoted $\chi_\ell(G)$, is the smallest k for which G is k -choosable. Since a k -assignment can assign the same k colors to every vertex of a graph, $\chi(G) \leq \chi_\ell(G)$.

1.2. Equitable total coloring and list equitable coloring

We now briefly review work that has been done on equitably coloring total graphs, and we review a list analogue of equitable coloring. We will then spend the remainder of the paper focused upon list equitable total coloring.

1.2.1. Equitable total coloring

The study of equitable total coloring was initiated by Fu in 1994 [11]. Specifically, Fu introduced the Equitable Total Coloring Conjecture which we now state.

Conjecture 1 (Equitable Total Coloring Conjecture [11]). *For every graph G , $T(G)$ has an equitable k -coloring for each $k \geq \max\{\chi(T(G)), \Delta(G) + 2\}$.*

The “ $\Delta(G) + 2$ ” is required because Fu [11] found an infinite family of graphs G with $\chi''(G) = \Delta(G) + 1$ but $T(G)$ is not equitably $(\Delta(G) + 1)$ -colorable (cf. Proposition 2.10 in [11]). Note that if the Total Coloring Conjecture is true, we would have $\max\{\chi''(G), \Delta(G) + 2\} = \Delta(G) + 2$.

Fu [11] showed that Conjecture 1 holds for complete bipartite graphs, complete t -partite graphs of odd order, trees, and certain split graphs. Equitable total coloring has also been studied for graphs with maximum degree 3 [36], joins of certain graphs [13, 14, 39], the Cartesian product of cycles [6], and the corona product of cubic graphs [12].

1.2.2. List equitable coloring

In 2003, Kostochka, Pelsmajer, and West introduced a list analogue of equitable coloring called equitable choosability [20]. Suppose that L is a k -assignment for graph G . An *equitable L -coloring* of G is a proper L -coloring, f , of G such that f uses no color more than $\lceil |V(G)|/k \rceil$ times¹. When an equitable L -coloring of G exists we say that G is *equitably L -colorable*. Graph G is *equitably k -choosable* if G is equitably L -colorable whenever L is a k -assignment for G .

We now mention a convention used in this paper. Suppose that H is a subgraph of G , and suppose that L is a k -assignment for G . When there is an equitable L' -coloring of H where L' is the k -assignment for H defined by $L'(v) = L(v)$ for each $v \in V(H)$, we say H has an equitable L -coloring. Notice an equitable L -coloring of H requires color classes of size at most $\lceil |V(H)|/k \rceil$ which may be more restrictive than the bound required for an equitable L -coloring of G .

It is important to note that, similar to equitable coloring, making the lists larger may make equitable list coloring more difficult in certain cases. Indeed $K_{1,9}$ is equitably 4-choosable, but it is not equitably 5-choosable. Also, equitable k -choosability does not imply equitable k -colorability unless $k = 2$. Indeed $K_{1,6}$ is equitably 3-choosable, but it is not equitably 3-colorable (see [29]). In [20], there are (perhaps surprising) conjectures that are list analogues of Hajnal and Szemeredi’s result and the Δ -Equitable Coloring Conjecture.

Conjecture 2 [20]. *Every graph G is equitably k -choosable when $k \geq \Delta(G) + 1$.*

Conjecture 3 [20]. *A connected graph G is equitably k -choosable for each $k \geq \Delta(G)$ if it is different from K_m , C_{2m+1} , and $K_{2m+1,2m+1}$.*

In [20], it is shown that Conjectures 2 and 3 hold for forests, connected interval graphs, and 2-degenerate graphs with maximum degree at least 5. It

¹When it comes to equitable choosability, the word equitable indicates that no color is used excessively often.

was also shown in [20] that if G is a graph and $k \geq \max\{\Delta(G), |V(G)|/2\}$, then G is equitably k -choosable unless G contains K_{k+1} or is $K_{k,k}$ with k odd in the latter case. Thus, Conjecture 3 is true for small graphs (at most $2k$ vertices). Conjectures 2 and 3 have also been verified for outerplanar graphs [42], powers of paths and cycles [18], series-parallel graphs [41], and certain planar graphs (see [23, 40, 43]). In 2013, Kierstead and Kostochka made substantial progress on Conjecture 2, as follows.

Theorem 4 [19]. *If G is any graph, then G is equitably k -choosable whenever*

$$k \geq \begin{cases} \Delta(G) + 1 & \text{if } \Delta(G) \leq 7, \\ \Delta(G) + \frac{\Delta(G)+6}{7} & \text{if } 8 \leq \Delta(G) \leq 30, \\ \Delta(G) + \frac{\Delta(G)}{6} & \text{if } \Delta(G) \geq 31. \end{cases}$$

1.3. List equitable total coloring

In 2018, Kaul, Pelsmajer, and the first author, began studying the equitable choosability of total graphs [18] which was originally suggested by Nakprasit [30]. Motivated by the Equitable Total Coloring Conjecture (Conjecture 1), they introduced the List Equitable Total Coloring Conjecture (LETCC for short).

Conjecture 5 (List Equitable Total Coloring Conjecture [18]). *For every graph G , $T(G)$ is equitably k -choosable for each $k \geq \max\{\chi_\ell(T(G)), \Delta(G) + 2\}$.*

Note that since $\Delta(T(G)) = 2\Delta(G)$, the LETCC is saying something stronger about total graphs than Conjectures 2 and 3 when $\Delta(G) > 2$. Also, Fu's infinite family of graphs G with $\chi''(G) = \Delta(G) + 1$ and $T(G)$ is not equitably $\Delta(G) + 1$ -colorable also has the property that $\chi_\ell(T(G)) = \Delta(G) + 1$ and $T(G)$ is not equitably $(\Delta(G) + 1)$ -choosable. So the LETCC would be sharp if true. The LETCC has been verified for all graphs G with $\Delta(G) \leq 2$, stars, double stars, and trees of maximum degree 3 (see [18, 28]).

1.4. Outline of results and an open question

In this paper, we study list equitable total coloring of generalized theta graphs. Suppose that $m \in \mathbb{N}$ and $l_1, \dots, l_m \in \mathbb{N}$ satisfy $l_1 \leq \dots \leq l_m$. Then, the *generalized theta graph* $\Theta(l_1, \dots, l_m)$ is the equivalence class of graphs consisting of two vertices joined by internally disjoint paths of lengths l_1, \dots, l_m . We will assume that $l_2 \geq 2$ when $m \geq 2$ since we will only be considering simple graphs in this paper.² Studying list equitable total coloring of generalized theta graphs is quite natural as theta graphs and generalized theta graphs have many interesting

²For the remainder of this paper, whenever we see $\Theta(l_1, \dots, l_m)$, we will always assume $m, l_1, \dots, l_m \in \mathbb{N}$ with $l_1 \leq \dots \leq l_m$ and $l_2 \geq 2$ when $m \geq 2$.

properties that have been studied by many researchers (see [2, 3, 4, 10, 21, 24, 32]). For this paper, we prove a positive answer to the question: Does the LETCC hold for generalized theta graphs?

In order to build up to the generalized theta graph, in Section 2 we study list equitable total coloring of subdivisions of stars. We say that H is a *subdivision of G* if H is a graph obtained from G by replacing the edges of G with internally disjoint paths. In Section 2, we prove the following theorem.

Theorem 6. *Suppose G is a subdivision of $K_{1,m}$. If $m = 1$, then $T(G)$ is equitably k -choosable whenever $k \geq 3$. Otherwise $T(G)$ is equitably k -choosable whenever $k \geq m + 1$.*

Suppose G is a subdivision of $K_{1,m}$. It is worth noting that Theorem 6 is the best result possible since $\chi_\ell(T(G)) \geq \max\{3, m + 1\}$. Also, the result of Theorem 6 is saying something stronger than: the LETCC holds for subdivisions of stars. This is because the LETCC only says that $T(G)$ should be equitably k -choosable for each $k \geq m + 2$.

Finally, in Section 3 we prove the following.

Theorem 7. *Suppose $G = \Theta(l_1, \dots, l_m)$, then $T(G)$ is equitably k -choosable whenever $k \geq m + 2$.*

So, the LETCC holds for generalized theta graphs. Notice that in Theorem 7, $T(G)$ is a path square and cycle square when m is 1 and 2 respectively. Since path squares with at least 3 vertices are not equitably 2-choosable, and all cycle squares with order not divisible by 3 are not equitably 3-choosable (see [18] for further details), one can see that in the case of $m = 1, 2$, $T(G)$ may not be equitably $(m + 1)$ -choosable. The question of whether $T(G)$ is equitably $(m + 1)$ -choosable when $m \geq 3$ is open.

Question 8. *Suppose $G = \Theta(l_1, \dots, l_m)$. If $m \geq 3$, does it follow that $T(G)$ is equitably $(m + 1)$ -choosable?*

2. SUBDIVISIONS OF STARS

In this section, we prove Theorem 6. Suppose that $m \in \mathbb{N}$ and $l_1, \dots, l_m \in \mathbb{N}$ satisfy $l_1 \leq \dots \leq l_m$. Then, we use $B(l_1, \dots, l_m)$ to denote the equivalence class of subdivisions of $K_{1,m}$ where the edges of $K_{1,m}$ have been replaced with internally disjoint paths of lengths: l_1, \dots, l_m .³

For the remainder of this paper when $G = B(l_1, \dots, l_m)$, we assume that $V(G) = \{u\} \cup \{v_{i,j} : i \in [m], j \in [l_i]\}$, and the edges of G are drawn so that for

³For the remainder of this paper, whenever we see $B(l_1, \dots, l_m)$, we will always assume $m, l_1, \dots, l_m \in \mathbb{N}$ with $l_1 \leq \dots \leq l_m$.

each $i \in [m]$ vertices are adjacent if and only if they appear consecutively in the ordering: $u, v_{i,1}, \dots, v_{i,l_i}$.

Note that when $G = B(l_1, \dots, l_m)$, $T(G)$ is isomorphic to a copy of $[B(2l_1, \dots, 2l_m)]^2$. So, in order to prove Theorem 6, we begin by proving the following result which will imply Theorem 6 for each $m \geq 3$.

Theorem 9. *For $m \geq 3$, $(B(l_1, \dots, l_m))^2$ is equitably k -choosable for each $k \geq m + 1$.*

Notice that for $m \geq 3$ Theorem 9 is saying something stronger than Theorem 6 since we are allowing any of the natural numbers l_1, \dots, l_m to be odd. We now prove a Lemma that is closely related to a Lemma appearing in [20]; we use this Lemma frequently to prove our results.

Lemma 10. *Let G be a graph and let L be a k -assignment for G . Suppose that $|V(G)| = kq + r$ where $1 \leq r \leq k$. Suppose t satisfies $r \leq t \leq k$. Let $S = \{x_1, \dots, x_t\}$ be a set of t distinct vertices in G . If $G - S$ has an equitable L -coloring and*

$$|N_G(x_i) - S| \leq k - i$$

for each $i \in [t]$, then G has an equitable L -coloring.

Proof. Suppose that f is an equitable L -coloring of $G - S$ (notice $G - S$ could be the empty graph). Note that no color is used more than q times by f . In an equitable L -coloring of G , we must use no color more than $\lceil (kq + r)/k \rceil = q + 1$ times. Let $L'(x_i) = L(x_i) - \{f(v) : v \in N_G(x_i) - S\}$ for each $i \in [t]$. Since $|N_G(x_i) - S| \leq k - i$, we know that $|L'(x_i)| \geq i$. So, there is a proper L' -coloring of $G[S]$ that uses t distinct colors. Such a coloring along with f completes an equitable L -coloring of G . ■

We will now prove five lemmas that will imply Theorem 9. The first two of the five lemmas will take care of the case where $k \geq m + 2$, and the last three of the five lemmas will deal with $k = m + 1$.

Lemma 11. *Suppose $m \geq 3$. If $H = B(l_1, l_2, \dots, l_m)$ and $G = H^2$, then G is equitably k -choosable whenever $k \geq m + 3$.*

Proof. The result is obvious when $k \geq |V(G)| = 1 + \sum_{i=1}^m l_i$. So, we may assume that L is an arbitrary k -assignment for G such that $m + 3 \leq k < 1 + \sum_{i=1}^m l_i$. We will show that G is equitably L -colorable.

Since $1 + \sum_{i=1}^m l_i > m + 3$, $l_m > 1$. Let $S_0 = \{v_{i,1} : i \in [m]\} \cup A$ where $A = \{u, v_{m,2}\}$ if $v_{m,3} \notin V(G)$ (i.e. $l_m = 2$) and $A = \{u, v_{m,2}, v_{m,3}\}$ otherwise. Note $m + 2 \leq |S_0| \leq m + 3$. Let $d = k - |S_0|$. Let S_1 be an arbitrary subset of $V(G) - S_0$ of size d , and let $S = S_0 \cup S_1$. Note that $G - S$ is a graph with

maximum degree at most 4. By Theorem 4, there is an equitable L -coloring of $G - S$.

Now, let $x_1 = u$, $x_{k-3} = v_{m-2,1}$, $x_{k-2} = v_{m-1,1}$, $x_{k-1} = v_{m,2}$, and $x_k = v_{m,1}$. We then arbitrarily name the remaining vertices in S : x_2, \dots, x_{k-4} in an injective fashion. By the way S is constructed, $|N_G(x_{k-i}) - S| \leq 2$ for $i = 2, 3$, $|N_G(x_{k-1}) - S| \leq 1$, and $|N_G(x_k) - S| = 0$. Moreover, $|N_G(x_1) - S| \leq m - 1 \leq (m + 3) - 1 \leq k - 1$. Finally, for $2 \leq i \leq k - 4$, $|N_G(x_i) - S| \leq 4 \leq k - i$. So, Lemma 10 implies that G is equitably L -colorable. ■

Lemma 12. *Suppose $m \geq 3$. If $H = B(l_1, l_2, \dots, l_m)$ and $G = H^2$, then G is equitably $(m + 2)$ -choosable.*

Proof. Note that the result is obvious when $l_m = 1$ since G is a copy of K_{m+1} when $l_m = 1$. So, we assume that $l_m > 1$. Let L be an arbitrary $(m + 2)$ -assignment for G . We will show that G is equitably L -colorable in the following cases: $l_m = 2$, $l_m = 3$, and $l_m \geq 4$.

In the case $l_m = 2$, let $S = \{v_{i,1} : i \in [m]\} \cup \{u, v_{m,2}\}$. We name the vertices of S as x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m+2} = v_{m,2}$, and for each $j \in [m]$, $x_{j+1} = v_{j,1}$. By Theorem 4, $G - S$ is equitably L -colorable. It is easy to see that $|N_G(x_1) - S| \leq m - 1 \leq (m + 2) - 1$, $|N_G(x_{m+2}) - S| = 0$, $|N_G(x_{m+1}) - S| = 0$, and for each $2 \leq j \leq m$, $|N_G(x_j) - S| \leq 1 \leq m + 2 - j$. Thus, G is equitably L -colorable by Lemma 10.

In the case $l_m = 3$, let $S = \{v_{i,1} : 2 \leq i \leq m\} \cup \{u, v_{m,2}, v_{m,3}\}$. We name the vertices of S as x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m+2} = v_{m,3}$, $x_{m+1} = v_{m,2}$, and for each $2 \leq i \leq m$, $x_i = v_{i,1}$. By Theorem 4, $G - S$ is equitably L -colorable. It is easy to see that $|N_G(x_1) - S| \leq m \leq (m + 2) - 1$, $|N_G(x_{m+2}) - S| = 0$, $|N_G(x_{m+1}) - S| = 0$, $|N_G(x_m) - S| = 1$, and for each $2 \leq j \leq m - 1$, $|N_G(x_j) - S| \leq 3 \leq m + 2 - j$. Thus, G is equitably L -colorable by Lemma 10.

In the case $l_m \geq 4$, let $S = \{v_{i,1} : 3 \leq i \leq m\} \cup \{u, v_{m,2}, v_{m,3}, v_{m,4}\}$. We name the vertices of S as x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m+2} = v_{m,2}$, $x_{m+1} = v_{m,3}$, $x_m = v_{m,4}$, and $x_{m-1} = v_{m,1}$. Finally, if $m \geq 4$, then for each $2 \leq j \leq m - 2$ we let $x_j = v_{j+1,1}$. Notice $G - S$ has maximum degree at most 4. So, by Theorem 4, $G - S$ is equitably L -colorable. It is easy to see that $|N_G(x_1) - S| \leq m + 1 \leq (m + 2) - 1$, $|N_G(x_{m+2}) - S| = 0$, $|N_G(x_{m+1}) - S| \leq 1$, $|N_G(x_m) - S| \leq 2$, and $|N_G(x_{m-1}) - S| = 2$. Finally, if $m \geq 4$ then for each $2 \leq j \leq m - 2$, $|N_G(x_j) - S| \leq 4 \leq m + 2 - j$. Thus, G is equitably L -colorable by Lemma 10. ■

We now turn our attention to the case of $k = m + 1$. Notice that in the case when $m = 3$, when we try to use Lemma 10, we will no longer be able to use Theorem 4 to show $G - S$ is equitably L -colorable. So, we need a result from [18].

Proposition 13 [18]. *For $p, n \in \mathbb{N}$, P_n^p is equitably k -choosable whenever $k \geq p + 1$.*

Notice that Proposition 13 immediately implies that if G is a spanning subgraph of a path square, then G is equitably k -choosable whenever $k \geq 3$.

Lemma 14. *Suppose $m \geq 3$. If $H = B(l_1, l_2, \dots, l_m)$, $G = H^2$, and $l_1 \leq l_2 \leq 2$, then G is equitably $(m + 1)$ -choosable.*

Proof. We may assume that $|V(G)| > m + 1$. Suppose L is an arbitrary $(m + 1)$ -assignment for G . In the case that $l_2 = 1$, let $S = \{u, v_{1,1}, v_{2,1}, \dots, v_{m,1}\}$. Clearly $G - S$ has an equitable L -coloring by Proposition 13. Let $x_1 = u$, and let $x_{i+1} = v_{m+1-i,1}$ for each $i \in [m]$. Note that $|N_G(x_1) - S| \leq m - 2 \leq (m + 1) - 1$, $|N_G(x_i) - S| \leq 2 \leq m + 1 - i$ for all $2 \leq i \leq m - 1$, and $|N_G(x_i) - S| = 0 \leq m + 1 - i$ when $i = m, m + 1$. By Lemma 10, we know that G has an equitable L -coloring.

In the case that $l_2 = 2$, let $S = \{u, v_{1,1}, v_{2,1}, \dots, v_{m-1,1}, v_{2,2}\}$. Clearly $G - S$ has an equitable L -coloring by Proposition 13. Let $x_1 = u$, $x_{m-1} = v_{1,1}$, $x_m = v_{2,1}$, and $x_{m+1} = v_{2,2}$. If $m \geq 4$, let $x_{i+1} = v_{m-i,1}$ for each $i \in [m - 3]$. Note: $|N_G(x_1) - S| \leq m + 1 - 1$, $|N_G(x_{m-1}) - S| \leq 2$, $|N_G(x_m) - S| = 1$, and $|N_G(x_{m+1}) - S| = 0$. Furthermore, if $m \geq 4$, for each $2 \leq i \leq m - 2$, $|N_G(x_i) - S| \leq 3 \leq m + 1 - i$. By Lemma 10, we know that G has an equitable L -coloring. ■

We now would like to prove for $m \geq 3$ that if $H = B(l_1, l_2, \dots, l_m)$ and $G = H^2$, then G is equitably $(m + 1)$ -choosable by induction on $\sum_{i=1}^m l_i$. Lemma 14 takes care of the base case. The next lemma takes care of a small issue in the inductive step when $m = 3$.

Lemma 15. *If $H = B(1, 3, 3)$ and $G = H^2$, then G is equitably 4-choosable.*

Proof. Suppose that L is an arbitrary 4-assignment for G . Let $S = \{v_{2,3}, v_{3,3}, v_{3,2}, v_{3,1}\}$. Note that $G - S$ is the square of a path. So, by Proposition 13 we know that $G - S$ has an equitable L -coloring. We then let $x_1 = v_{3,1}$, $x_2 = v_{2,3}$, $x_3 = v_{3,2}$, and $x_4 = v_{3,3}$. Note that $|N_G(x_i) - S| = 4 - i$ for all $i \in [4]$. So, by Lemma 10, we know that G has an equitable L -coloring. ■

We are finally ready to complete our proof of Theorem 9.

Lemma 16. *Suppose that $m \geq 3$. If $H = B(l_1, l_2, \dots, l_m)$ and $G = H^2$, then G is equitably $(m + 1)$ -choosable.*

Proof. We will prove the desired by induction on $\sum_{i=1}^m l_i = |V(G)| - 1$. Let $C = \sum_{i=1}^m l_i$. Let L be an arbitrary $(m + 1)$ -assignment for G . We will show an equitable L -coloring of G exists for each $C \geq m$. For the base case suppose that

$m \leq C \leq 3m - 3$. Since $C \leq 3m - 3$, we know that $l_1 \leq l_2 \leq 2$. So, the desired result holds by Lemma 14.

For the inductive step suppose that $C \geq 3m - 2$ and assume that the desired result holds for all natural numbers less than C and at least m . Note that when $l_2 \leq 2$ the result holds by Lemma 14; so, we may assume that $l_2 \geq 3$.

If $l_m \geq 4$ we let $S = \{v_{j,l_j} : 2 \leq j \leq m - 1\} \cup \{v_{m,l_m-2}, v_{m,l_m-1}, v_{m,l_m}\}$. By the inductive hypothesis, $G - S$ is equitably L -colorable. Now let $x_i = v_{i+1,l_{i+1}}$ for all $i \in [m - 2]$, $x_{m-1} = v_{m,l_m-2}$, $x_m = v_{m,l_m-1}$, and $x_{m+1} = v_{m,l_m}$. Note that $|N_G(x_i) - S| = 2 \leq (m + 1) - i$ for all $i \in [m - 1]$, $|N_G(x_m) - S| = 1$, and $|N_G(x_{m+1}) - S| = 0$. Thus, Lemma 10 implies that G is equitably L -colorable.

Now, assume that $l_m \leq 3$ (note that this means $l_m = 3$ since $l_2 \geq 3$). This implies that $3m - 2 \leq C \leq 3m$. Let $d = C - (2m + 1)$. In the case that $C = 3m - 2$, we may assume that $m \geq 4$ by Lemma 15. Note that $m - 3 \leq d \leq m - 1$ and that $|V(G)| - d = C + 1 - d = 2m + 2$. We let $S = \{v_{2,l_2}, \dots, v_{1+d,l_{1+d}}\}$. By the inductive hypothesis, we know that $G - S$ has an equitable L -coloring. Let $x_i = v_{i+1,l_{i+1}}$ for all $i \in [d]$. Note that $|N_G(x_i) - S| = 2 \leq (m + 1) - i$ for all $i \in [d]$. Lemma 10 then implies that G has an equitable L -coloring. The induction step is now complete. \blacksquare

Finally, notice the result of Theorem 6 is implied by Theorem 9 when $m \geq 3$, and the result of Theorem 6 is implied by Proposition 13 when $m = 1, 2$.

3. GENERALIZED THETA GRAPHS

In this Section, we will prove Theorem 7. Throughout this section if $G = \Theta(l_1, \dots, l_m)$, we will assume that the vertices that are the common endpoints in $V(G)$ are u and w . We also let the vertices of the i th path be⁴:

$$u, v_{i,1}, \dots, v_{i,l_i-1}, w.$$

When it comes to proving Theorem 7, it is crucial to note that if $G = \Theta(l_1, \dots, l_m)$, then $T(G)$ is a copy of $[\Theta(2l_1, \dots, 2l_m)]^2$ where $2l_2 \geq 4$ whenever $m \geq 2$. So, the results in this Section will focus upon the equitable choosability of the squares of generalized theta graphs with sufficiently long paths. Notice that if $G = \Theta(l_1, \dots, l_m)$, $T(G)$ is a path square and cycle square on at least 6 vertices when m is 1 and 2 respectively. So, when $m = 1, 2$ the result of Theorem 7 is implied by Proposition 13 and the following result.

Proposition 17 [18]. *Suppose that $p, n \in \mathbb{N}$ with $p \geq 2$ and $n \geq 2p + 2$. Then, C_n^p is equitably k -choosable for each $k \geq 2p$.*

⁴Notice that if $l_1 = 1$, the first path has no internal vertices.

So, to complete the proof of Theorem 7, we may focus our attention on the case where $m \geq 3$. We begin by proving the following result.

Theorem 18. *For $m \geq 3$, $l_1 \geq 2$, and $l_2 \geq 4$, $[\Theta(l_1, l_2, \dots, l_m)]^2$ is equitably k -choosable whenever $k \geq m + 3$.*

We will establish two lemmas that will immediately imply Theorem 18.

Lemma 19. *Suppose that $m \geq 3$. Suppose $H = \Theta(l_1, l_2, \dots, l_m)$ where $l_1 \geq 2$ and $l_2 \geq 4$. If $G = H^2$, then G is equitably k -choosable whenever $k \geq 2m + 2$.*

Proof. The result is obvious when $k \geq |V(G)|$; thus, we will assume that $2m + 2 \leq k < |V(G)| = 2 + \sum_{i=1}^m (l_i - 1)$. Suppose L is an arbitrary k -assignment for G . Let

$$S_0 = \{u, w, v_{m,2}, v_{m,3}\} \cup \{v_{i,1} : 1 \leq i \leq m\} \cup \{v_{i,l_i-1} : 3 \leq i \leq m\}.$$

Note that $G - S_0$ is a spanning subgraph of a disjoint union of path squares. Let $d = k - |S_0|$. Let S_1 be an arbitrary subset of $V(G) - S_0$ of size d . Then let $S = S_0 \cup S_1$. Note that $G - S$ has maximum degree at most 4. By Theorem 4, there is an equitable L -coloring of $G - S$. We will name the vertices of S : x_1, x_2, \dots, x_k such that $x_1 = w$, $x_2 = u$, $x_{k-3} = v_{2,1}$, $x_{k-2} = v_{m,3}$, $x_{k-1} = v_{m,2}$, $x_k = v_{m,1}$. We then arbitrarily name the remaining vertices in S : x_3, x_4, \dots, x_{k-4} in an injective fashion. For $i \in \{k-3, k-2, k-1, k\}$ we have $|N_G(x_i) - S| \leq k - i$. For $3 \leq i \leq k-4$ we have that $|N_G(x_i) - S| \leq 4 \leq k - i$. Finally, $|N_G(x_2) - S| \leq m - 1 \leq k - 2$ and $|N_G(x_1) - S| \leq m + 2 \leq k - 1$. By Lemma 10, there is an equitable L -coloring for G . ■

Lemma 20. *Suppose that $m \geq 3$. Suppose $H = \Theta(l_1, l_2, \dots, l_m)$ where $l_1 \geq 2$ and $l_2 \geq 4$. If $G = H^2$, then G is equitably k -choosable whenever $m + 3 \leq k \leq 2m + 1$.*

Proof. Suppose that L is an arbitrary k -assignment for G such that $m + 3 \leq k \leq 2m + 1$. We let $S = \{u, v_{m,3}\} \cup \{v_{i,1} : i \in [m]\} \cup \{v_{i,2} : 2m + 3 - k \leq i \leq m\}$. Note that $|S| = k$. Note that there is an $r \in \{m - 2, m - 1, m\}$ and natural numbers a_1, \dots, a_r with $a_1 \leq \dots \leq a_r$ such that $G - S$ is isomorphic to $[B(a_1, a_2, \dots, a_r)]^2$. When $r \leq 2$, we know that $G - S$ has an equitable L -coloring by Theorem 4 since $\Delta(G - S) \leq 4$. When $r \geq 3$, we know that $G - S$ has an equitable L -coloring by Lemma 11. Let $x_1 = v_{m,3}$, $x_2 = v_{1,1}$, $x_3 = u$, $x_{k-1} = v_{m,2}$, and $x_k = v_{m,1}$. Finally, we arbitrarily name the remaining vertices in S : x_3, x_4, \dots, x_{k-2} in an injective fashion. Note that $|N_G(x_1) - S| \leq m \leq k - 1$, $|N_G(x_2) - S| \leq m \leq k - 2$, $|N_G(x_3) - S| \leq m - 1 \leq k - 3$, $|N_G(x_{k-1}) - S| = 1$, and $|N_G(x_k) - S| = 0$. Finally note that $|N_G(x_i) - S| \leq 2 \leq k - i$ for all $4 \leq i \leq k - 2$. So by Lemma 10 we know that G has an equitable L -coloring. ■

We are now finished with the proof of Theorem 18. To complete the proof of Theorem 7 we must show that if $G = \Theta(l_1, l_2, \dots, l_m)$, then $T(G)$ is equitably k -choosable when $m \geq 3$ and $k = m + 2$. Recall for $m \geq 3$ that if $G = \Theta(l_1, l_2, \dots, l_m)$, then $T(G)$ is a copy of $[\Theta(2l_1, 2l_2, \dots, 2l_m)]^2$ where $2l_2 \geq 4$. So, we begin working on the case of $k = m + 2$ by dealing with $[\Theta(2, 4, \dots, 4)]^2$ and $[\Theta(4, \dots, 4)]^2$. We will then finish the case of $k = m + 2$ by considering $[\Theta(l_1, l_2, \dots, l_m)]^2$ with $l_1 \geq 2$, $l_2 \geq 4$, and $l_m \geq 6$.

We begin with two specific cases to which our general arguments do not apply.

Lemma 21. *Suppose $H = \Theta(2, 4, 4)$ and $G = H^2$. Then, G is equitably 5-choosable.*

Proof. Suppose L is an arbitrary 5-assignment for G . We will show that G has an equitable L -coloring in two cases: (1) $L(v)$ is the same list for each $v \in V(G)$ and (2) there exist $x, y \in V(G)$ such that $L(x) \neq L(y)$. In case (1) we may suppose $L(v) = \{1, 2, 3, 4, 5\}$ for each $v \in V(G)$. We let f be the proper L -coloring of G defined as follows: $f(u) = 1$, $f(v_{1,1}) = 2$, $f(v_{2,1}) = 3$, $f(v_{3,1}) = 4$, $f(v_{2,2}) = 2$, $f(v_{3,2}) = 3$, $f(v_{2,3}) = 4$, $f(v_{3,3}) = 1$, and $f(w) = 5$. Clearly, f is an equitable L -coloring of G .

We now turn our attention to case (2). Let $S_1 = \{v_{2,1}, v_{2,2}, v_{2,3}\}$ and $S_2 = \{v_{3,1}, v_{3,2}, v_{3,3}\}$. Note that $(V(G) - S_1) \cup (V(G) - S_2) = V(G)$ and $(V(G) - S_1) \cap (V(G) - S_2) \neq \emptyset$. So, it must be the case that there exists at least two lists in either $\{L(v) : v \in V(G) - S_1\}$ or $\{L(v) : v \in V(G) - S_2\}$. We assume without loss of generality that there are at least two lists in $\{L(v) : v \in V(G) - S_1\}$. It is then possible to find a proper L -coloring, f , of $G - S_1$ that uses six distinct colors. Let $L'(v_{2,i}) = L(v_{2,i}) - \{f(v) : v \in N_G(v_{2,i}) - S_1\}$ for each $i \in [3]$. Note that $|L'(v_{2,1})| \geq 2$, $|L'(v_{2,2})| \geq 3$, and $|L'(v_{2,3})| \geq 2$. So, there is a proper L' -coloring of $G[S_1]$. Such a coloring completes an equitable L -coloring of G . ■

Lemma 22. *Suppose $H = \Theta(2, 4, 4, 4)$ and $G = H^2$. Then, G is equitably 6-choosable.*

Proof. Suppose L is an arbitrary 6-assignment for G . Note an equitable L -coloring of G uses no color more than twice. We will show that G has an equitable L -coloring in two cases: (1) $L(v)$ is the same list for each $v \in V(G) - \{u, w\}$ and (2) there exist $x, y \in V(G) - \{u, w\}$ such that $L(x) \neq L(y)$. In case (1) we may suppose $L(v) = \{1, 2, 3, 4, 5, 6\}$ for each $v \in V(G) - \{u, w\}$. Let f be the proper L -coloring for $G - \{u, w\}$ given by: $f(v_{1,1}) = 2$, $f(v_{2,1}) = 3$, $f(v_{3,1}) = 4$, $f(v_{4,1}) = 5$, $f(v_{2,2}) = 2$, $f(v_{3,2}) = 6$, $f(v_{4,2}) = 6$, $f(v_{2,3}) = 1$, $f(v_{3,3}) = 5$, and $f(v_{4,3}) = 4$. Then, let $L'(u) = L(u) - \{2, 3, 4, 5, 6\}$ and $L'(w) = L(w) - \{1, 2, 4, 5, 6\}$. As long as $L'(u)$ and $L'(w)$ are not the same set of size 1, we can find a proper L' -coloring of $G[\{u, w\}]$ which along with f completes an equitable L -coloring of G .

So, we assume that $L'(w) = L'(u) = \{7\}$. This means $L(u) = \{2, 3, 4, 5, 6, 7\}$ and $L(w) = \{1, 2, 4, 5, 6, 7\}$. An equitable L coloring, g , of G can then be constructed as follows: $g(u) = 7$, $g(v_{1,1}) = 2$, $g(v_{2,1}) = 3$, $g(v_{3,1}) = 4$, $g(v_{4,1}) = 5$, $g(v_{2,2}) = 2$, $g(v_{3,2}) = 3$, $g(v_{4,2}) = 6$, $g(v_{2,3}) = 6$, $g(v_{3,3}) = 5$, $g(v_{4,3}) = 4$, and $g(w) = 1$.

For case (2), let $S_1 = \{u, w, v_{4,1}, v_{4,2}, v_{3,2}\}$, $S_2 = \{u, w, v_{3,1}, v_{3,2}, v_{2,2}\}$, and $S_3 = \{u, w, v_{2,1}, v_{2,2}, v_{4,2}\}$. Note that $(V(G) - S_1) \cup (V(G) - S_2) \cup (V(G) - S_3) = V(G) - \{u, w\}$ and $\bigcap_{i=1}^3 (V(G) - S_i) \neq \emptyset$. Thus, there must exist at least two lists in $\{L(v) : v \in V(G) - S_1\}$, $\{L(v) : v \in V(G) - S_2\}$, or $\{L(v) : v \in V(G) - S_3\}$. Assume without loss of generality that $\{L(v) : v \in V(G) - S_1\}$ contains at least two lists. It is possible to find a proper L -coloring, h , of $G - S_1$ that uses seven distinct colors. Let $L'(v) = L(v) - \{h(x) : x \in N_G(v) - S_1\}$ for each $v \in S_1$. Note that $|L'(u)| \geq 2$, $|L'(w)| \geq 1$, $|L'(v_{4,1})| \geq 2$, $|L'(v_{3,2})| \geq 4$, $|L'(v_{4,2})| \geq 5$. Now, we greedily construct a proper L' -coloring of $G[S_1]$, h' , that uses at least 4 distinct colors by coloring the vertices of S_1 in the order: $w, u, v_{4,1}, v_{3,2}, v_{4,2}$ (this is possible since $v_{4,1}$ is not adjacent to w in G). In the case $|h'(S_1)| = 5$, h together with h' forms an equitable L -coloring of G . So, we suppose that $|h'(S_1)| = 4$. By the way h' is constructed, it must be that $h'(w) = h'(v_{4,1}) = c$.

Now, for the sake of contradiction, we suppose that h together with h' is not an equitable L -coloring of G . This means that c was used by h . Note that $V(G) - S_1 \subseteq N_G(w) \cup N_G(v_{4,1})$. So, by the definition of L' , $c \notin L'(w)$ or $c \notin L'(v_{4,1})$ which is a contradiction. Thus h along with h' forms an equitable L -coloring of G . \blacksquare

We are now ready to prove two Lemmas that will complete the case of $k = m + 2$ for $[\Theta(2, 4, \dots, 4)]^2$ and $[\Theta(4, \dots, 4)]^2$.

Lemma 23. *Suppose $m \geq 5$. Suppose $H = \Theta(l_1, l_2, \dots, l_m)$ where $l_1 = 2$, and $l_2 = l_m = 4$. If $G = H^2$, then G is equitably $(m + 2)$ -choosable.*

Proof. Suppose that L is an arbitrary $(m + 2)$ -assignment for G . We will construct an equitable L -coloring of G . Since $m \geq 5$, in an equitable L -coloring of G , no color can be used more than $\lceil 3m/(m + 2) \rceil = 3$ times. Let $S_1 = \{u\} \cup \{v_{i,1} : i \in [m]\}$, $S_2 = \{w\} \cup \{v_{i,3} : 2 \leq i \leq m\}$, and $S_3 = \{v_{i,2} : 2 \leq i \leq m\}$. Note that $|S_1| = m + 1$, $|S_2| = m$, and $|S_3| = m - 1$. We begin by coloring the vertices in S_1 with $m + 1$ distinct colors. For each $v \in S_2$ suppose that $L'(v)$ is obtained from $L(v)$ by deleting the colors used on the neighbors of v in the coloring of the vertices in S_1 . Note that $|L'(v)| \geq m$ for each $v \in S_2$. So, we can color each $v \in S_2$ with a color $c \in L'(v)$ such that the vertices in S_2 are colored with m distinct colors.

Now, for each $v \in S_1 \cup S_2$ let $f(v)$ be the color used to color v . Note that f uses at most m colors twice, and f uses no color more than twice. For each $v \in S_3$ let $L''(v) = L(v) - \{f(x) : x \in (S_1 \cup S_2) \cap N_G(v)\}$. Since each vertex in S_3 has degree 4 in G , $|L''(v)| \geq m - 2$ for each $v \in S_3$.

If it is not the case that: $|L''(v)| = m - 2$ for all $v \in S_3$ and $L''(v_{2,2}) = L''(v_{3,2}) = \cdots = L''(v_{m,2})$, then it is clear that there exists a proper L'' -coloring of $G[S_3]$ that uses $m - 1$ distinct colors which completes an equitable L -coloring of G . So, we assume that $|L''(v)| = m - 2$ for all $v \in S_3$ and $L''(v_{2,2}) = L''(v_{3,2}) = \cdots = L''(v_{m,2})$. Let $A = L''(v_{2,2})$. For the sake of contradiction, we assume that all colors in A are used twice by f . Note that $f^{-1}(A) \subseteq S_1 \cup S_2$ and $|f^{-1}(A)| = 2|A| = 2m - 4$. Since $|S_1 \cup S_2| = 2m + 1$, there are at most 5 vertices in $(S_1 \cup S_2) - f^{-1}(A)$. Note that

$$\left| \bigcup_{i=2}^m N_G(v_{i,2}) \right| = 2m \geq 10 > 5.$$

So, there must be a $z \in S_1 \cup S_2$ that is in $f^{-1}(A) \cap \bigcup_{i=2}^m N_G(v_{i,2})$. This however implies that for some $2 \leq i \leq m$, $f(z)$ was deleted from $L(v_{i,2})$ when forming $L''(v_{i,2})$ which implies $f(z) \notin L''(v_{i,2})$ which is a contradiction.

Thus, there must exist an element $a \in A$ that was not used twice by f . So, we can complete an equitable L -coloring of G by coloring $v_{2,2}$ and $v_{3,2}$ with a and coloring the remaining vertices in S_3 with the $m - 3$ distinct colors in $A - \{a\}$. ■

Lemma 24. *Suppose $m \geq 3$. Suppose $H = \Theta(l_1, l_2, \dots, l_m)$ where $l_1 = l_m = 4$. If $G = H^2$, then G is equitably $(m + 2)$ -choosable.*

Proof. Suppose that L is an arbitrary $(m + 2)$ -assignment for G . We will construct an equitable L -coloring of G . Since $m \geq 3$, in an equitable L -coloring of G , no color can be used more than $\lceil (3m + 2)/(m + 2) \rceil = 3$ times. Let $S_1 = \{u, v_{1,2}\} \cup \{v_{i,1} : i \in [m]\}$, $S_2 = \{w\} \cup \{v_{i,3} : i \in [m]\}$, and $S_3 = \{v_{i,2} : 2 \leq i \leq m\}$. Note that $|S_1| = m + 2$, $|S_2| = m + 1$, and $|S_3| = m - 1$. We begin by coloring the vertices in S_1 with $m + 2$ distinct colors. For each $v \in S_2$ suppose that $L'(v)$ is obtained from $L(v)$ by deleting the colors used on the neighbors of v in the coloring of the vertices in S_1 . Note that $|L'(w)| \geq m + 1$ and $|L'(v_{i,3})| \geq m$ for all $i \in [m]$. So, we can color each $v \in S_2$ with a color $c \in L'(v)$ such that the vertices in S_2 are colored with $m + 1$ distinct colors.

Now, for each $v \in S_1 \cup S_2$ let $f(v)$ be the color used to color v . Note that f uses at most $m + 1$ colors twice, and f uses no color more than twice. For each $v \in S_3$ suppose that $L''(v) = L(v) - \{f(x) : x \in (S_1 \cup S_2) \cap N_G(v)\}$. Since each vertex in S_3 has degree 4 in G , $|L''(v)| \geq m - 2$ for each $v \in S_3$.

If it is not the case that: $|L''(v)| = m - 2$ for all $v \in S_3$ and $L''(v_{2,2}) = L''(v_{3,2}) = \cdots = L''(v_{m,2})$, then it is clear that there exists a proper L'' -coloring of $G[S_3]$ that uses $m - 1$ distinct colors which completes an equitable L -coloring of G . So, we assume that $|L''(v)| = m - 2$ for all $v \in S_3$ and $L''(v_{2,2}) = L''(v_{3,2}) = \cdots = L''(v_{m,2})$. Let $A = L''(v_{2,2})$. For the sake of contradiction, we assume that all colors in A are used twice by f . Note that $f^{-1}(A) \subseteq S_1 \cup S_2$ and

$|f^{-1}(A)| = 2|A| = 2m - 4$. Since $|S_1 \cup S_2| = 2m + 3$, there are at most 7 vertices in $(S_1 \cup S_2) - f^{-1}(A)$. Note that

$$\left| \bigcup_{i=2}^m N_G(v_{i,2}) \right| = 2m.$$

So, when $m \geq 4$, there must be a $z \in S_1 \cup S_2$ that is in $f^{-1}(A) \cap \bigcup_{i=2}^m N_G(v_{i,2})$, and we reach a contradiction as we did in the proof of Lemma 23. When $m = 3$ note that since $\{v_{1,1}, v_{1,2}, v_{1,3}\}$ is a clique in G , the single element in A must be in $f(\bigcup_{i=2}^3 N_G(v_{i,2}))$. So, when $m = 3$ there must also be a $z \in S_1 \cup S_2$ that is in $f^{-1}(A) \cap \bigcup_{i=2}^m N_G(v_{i,2})$, and we reach a contradiction as we did in the proof of Lemma 23.

Finally, we can complete an equitable L -coloring of G as we did in the proof of Lemma 23. \blacksquare

We now complete the proof of Theorem 7 with two lemmas. The next lemma will be important for proving the final lemma which will address all remaining cases needed for Theorem 7.

Lemma 25. *Suppose $m \geq 3$. Suppose $H = B(l_1, l_2, \dots, l_m)$, where $l_m \geq 3$. Suppose $G' = H^2$. Let G be the graph obtained from G' by adding an extra edge between the vertices v_{a,l_a} and v_{b,l_b} where a and b are chosen so that $1 \leq a < b \leq m$. Then, G is equitably $(m + 2)$ -choosable.*

Note that this lemma can be extended to $l_m = 2$, but this is not necessary to complete the proof of Theorem 7.

Proof. Let L be an arbitrary $(m + 2)$ -assignment for G . We will show that G has an equitable L -coloring in the following cases: (1) $l_m = 3$ and (2) $l_m \geq 4$.

In the case $l_m = 3$, let $S = \{u\} \cup \{v_{i,1} : 2 \leq i \leq m\} \cup \{v_{m,2}, v_{m,3}\}$. Then, we name the vertices of S : x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m+2} = v_{m,2}$, $x_{m+1} = v_{m,3}$, and for each $2 \leq i \leq m$, $x_i = v_{i,1}$. Note $\Delta(G - S) \leq 3$. So, by Theorem 4, $G - S$ is equitably L -colorable. It is easy to see that $|N_G(x_1) - S| = m \leq m + 2 - 1$, $|N_G(x_{m+2}) - S| = 0$, $|N_G(x_{m+1}) - S| \leq 1$, $|N_G(x_m) - S| = 1$, and for each $2 \leq j \leq m - 1$, $|N_G(x_j) - S| \leq 3 \leq m + 2 - j$. Thus G is equitably L -colorable by Lemma 10.

For case (2), we will show that G has an equitable L -coloring in the following subcases: (a) $m \geq 4$ and (b) $m = 3$. When $m \geq 4$, choose a t such that $t \in [m]$, $t \neq a$, $t \neq b$, and $t \neq m$. Let

$$S = \{u, v_{m,2}, v_{m,3}, v_{m,4}\} \cup (\{v_{i,1} : i \in [m]\} - \{v_{a,1}, v_{t,1}\}).$$

Then we name the vertices of S : x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m+2} = v_{m,2}$, $x_{m+1} = v_{m,3}$, $x_m = v_{m,4}$, and $x_{m-1} = v_{m,1}$. Finally, we arbitrarily name the

remaining vertices in S : x_2, \dots, x_{m-2} in an injective fashion. Note $\Delta(G-S) \leq 4$, and so by Theorem 4, $G-S$ is equitably L -colorable. It is easy to see that $|N_G(x_1) - S| = m+1 \leq m+2-1$, $|N_G(x_{m+2}) - S| = 0$, $|N_G(x_{m+1}) - S| \leq 1$, $|N_G(x_m) - S| \leq 2$, and $|N_G(x_{m-1}) - S| = 2$. Finally, for each $2 \leq j \leq m-2$, $|N_G(x_j) - S| \leq 4 \leq m+2-j$. Thus, G is equitably L -colorable by Lemma 10.

For subcase (b), $m = 3$, and we let $S = \{u, v_{3,1}, v_{3,2}, v_{3,3}, v_{3,4}\}$. Note that $\Delta(G-S) \leq 4$. So, by Theorem 4 we know that $G-S$ has an equitable L -coloring. We name the vertices of S as follows: $x_1 = u$, $x_2 = v_{3,1}$, $x_3 = v_{3,4}$, $x_4 = v_{3,3}$, and $x_5 = v_{3,2}$. Note $|N_G(x_1) - S| \leq 4$, $|N_G(x_2) - S| = 2$, $|N_G(x_3) - S| \leq 2$, $|N_G(x_4) - S| \leq 1$, and $|N_G(x_5) - S| = 0$. So, by Lemma 10 we know that G has an equitable L -coloring. ■

Lemma 26. *Suppose $m \geq 3$. Suppose $H = \Theta(l_1, l_2, \dots, l_m)$ where $l_1 \geq 2$, $l_2 \geq 4$, and $l_m \geq 6$. If $G = H^2$, then G is equitably $(m+2)$ -choosable.*

Proof. Let L be an arbitrary $(m+2)$ -assignment for G . We will show that G has an equitable L -coloring. We let

$$S = \{u, v_{m,2}, v_{m,3}, v_{m,4}\} \cup \{v_{i,1} : 3 \leq i \leq m\}.$$

There exist natural numbers a_1, \dots, a_m satisfying $a_1 \leq \dots \leq a_m$ and $a_m \geq 3$ such that the following holds. If $H = [B(a_1, \dots, a_m)]^2$, then there are $i, j \in [m]$ satisfying $1 \leq i < j \leq m$ such that $G-S$ is H plus an edge between the vertices v_{i,a_i} and v_{j,a_j} . Note that we know such an i and j exist since $v_{1,1}$ and $v_{2,1}$ are adjacent in $G-S$. Also note that we know $a_m \geq 3$ since $v_{2,1}$, $v_{2,2}$, and $v_{2,3}$ are in $G-S$. By Lemma 25 we know that $G-S$ has an equitable L -coloring. We name the vertices of S : x_1, x_2, \dots, x_{m+2} where $x_1 = u$, $x_{m-1} = v_{m,4}$, $x_m = v_{m,1}$, $x_{m+1} = v_{m,3}$ and $x_{m+2} = v_{m,2}$. Finally, if $m \geq 4$, we arbitrarily name the remaining vertices in S : x_2, \dots, x_{m-2} in an injective fashion. Note that $|N_G(x_1) - S| = m+1$, $|N_G(x_{m-1}) - S| = 2$, $|N_G(x_m) - S| = 1$, $|N_G(x_{m+1}) - S| = 1$, $|N_G(x_{m+2}) - S| = 0$, and $|N_G(x_i) - S| \leq 4 \leq m+2-i$ for all $2 \leq i \leq m-2$. Thus, G has an equitable L -coloring by Lemma 10. ■

Acknowledgment

The authors would like to thank Hemanshu Kaul and Michael Pelsmajer for their helpful comments on this paper. The authors would also like to thank the two anonymous referees whose comments greatly improved the readability of this paper.

REFERENCES

- [1] M. Behzad, Graphs and their Chromatic Numbers, Ph.D. Thesis (Michigan State University, 1965).

- [2] J. Beier, J. Fierson, R. Haas, H.M. Russel and K. Shavo, *Classifying coloring graphs*, Discrete Math. **339** (2016) 2100–2112.
doi:10.1016/j.disc.2016.03.003
- [3] J.I. Brown, C.A. Hickman, A.D. Sokal and D.G. Wagner, *On the chromatic roots of generalized theta graphs*, J. Combin. Theory Ser. B **83** (2001) 272–297.
doi:10.1006/jctb.2001.2057
- [4] J.M. Carraher, T. Mahoney, G.J. Puelo and D.B. West, *Sum-paintability of generalized theta-graphs*, Graphs Combin. **31** (2015) 1325–1334.
doi:10.1007/s00373-014-1441-1
- [5] B.-L. Chen, K.-W. Lih and P.-L. Wu, *Equitable coloring and the maximum degree*, European J. Combin. **15** (1994) 443–447.
doi:10.1006/eujc.1994.1047
- [6] T. Chunling, L. Xiaohui, Y. Yuansheng and L. Zhihe, *Equitable total coloring of $C_m \square C_n$* , Discrete Appl. Math. **157** (2009) 596–601.
doi:10.1016/j.dam.2008.08.030
- [7] A. Dong and J. Wu, *Equitable coloring and equitable choosability of planar graphs without chordal 4- and 6-cycles*, (2018), preprint.
arXiv: 1806.01064
- [8] A. Dong and X. Zhang, *Equitable coloring and equitable choosability of graphs with small maximum average degree*, Discuss. Math. Graph Theory **38** (2018) 829–839.
doi:10.7151/dmgt.2049
- [9] P. Erdős, *Problem 9*, in: Theory of Graphs and Its Applications, M. Fiedler (Ed(s)), (Proc. Sympos., Smolenice, 1963, Publ. House Czechoslovak Acad. Sci. Prague, 1964) 159.
- [10] P. Erdős, A.L. Rubin and H. Taylor, *Choosability in graphs*, Congr. Numer. **26** (1979) 125–127.
- [11] H.-L. Fu, *Some results on equalized total coloring*, Congr. Numer. **102** (1994) 111–119.
- [12] H. Furmańczyk and R. Zuazua, *Equitable total coloring of corona of cubic graphs*, Discuss. Math. Graph Theory, in press.
doi:10.7151/dmgt.2245
- [13] M.A. Gang and M.A. Ming, *The equitable chromatic number of some join graphs*, Open J. Appl. Sci. (2012) 96–99.
doi:10.4236/ojapps.2012.24B023
- [14] K. Gong, Z.-F. Zhang and J.-F. Wang, *Equitable total coloring of $F_n \vee W_n$* , Acta Math. Appl. Sin. Engl. Ser. **25** (2009) 83–86.
doi:10.1007/s10255-006-6031-4
- [15] A. Hajnal and E. Szemerédi, *Proof of a conjecture of Erdős*, in: Combinatorial Theory and Its Applications, A. Rényi and V.T. Sós (Ed(s)), (Vol. II, North-Holland, Amsterdam, 1970) 601–623.

- [16] S. Janson and A. Ruciński, *The infamous upper tail*, Random Structures Algorithms **20** (2002) 317–342.
doi:10.1002/rsa.10031
- [17] H. Kaul and S.H. Jacobson, *New global optima results for the Kauffman NK model: handling dependency*, Math. Program. **108** (2006) 475–494.
doi:10.1007/s10107-006-0719-3
- [18] H. Kaul, J.A. Mudrock and M.J. Pelsmajer, *Total equitable list coloring*, Graphs Combin. **34** (2018) 1637–1649.
doi:10.1007/s00373-018-1965-x
- [19] H.A. Kierstead and A.V. Kostochka, *Equitable list coloring of graphs with bounded degree*, J. Graph Theory **74** (2013) 309–334.
doi:10.1002/jgt.21710
- [20] A.V. Kostochka, M.J. Pelsmajer and D.B. West, *A list analogue of equitable coloring*, J. Graph Theory **44** (2003) 166–177.
doi:10.1002/jgt.10137
- [21] D. Laiche, I. Bouchemakh and É. Sopena, *On the packing coloring of undirected and oriented generalized theta graphs*, Australas. J. Combin. **66** (2016) 310–329.
- [22] M.E. Leidner, *A Study of the Total Colorings of Graphs*, Ph.D. Thesis (University of Louisville, 2012).
- [23] Q. Li and Y. Bu, *Equitable list coloring of planar graphs without 4- and 6-cycles*, Discrete Math. **309** (2009) 280–287.
doi:10.1016/j.disc.2007.12.070
- [24] R. Li, H. Broersma and S. Zhang, *Properly edge-colored theta graphs in edge-colored complete graphs*, Graphs Combin. **35** (2019) 261–286.
doi:10.1007/s00373-018-1989-2
- [25] K.-W. Lih, *The equitable coloring of graphs*, in: Handbook of Combinatorial Optimization, D.-Z. Du and P. Pardalos (Ed(s)), (Vol. III, Kluwer, Dordrecht, 1998) 543–566.
- [26] K.-W. Lih and P.-L. Wu, *On equitable coloring of bipartite graphs*, Discrete Math. **151** (1996) 155–160.
doi:10.1016/0012-365X(94)00092-W
- [27] W. Meyer, *Equitable coloring*, Amer. Math. Monthly **80** (1973) 920–922.
doi:10.1080/00029890.1973.11993408
- [28] J. Mudrock, *On the List Coloring Problem and Its Equitable Variants*, Ph.D. Thesis (Illinois Institute of Technology, 2018).
- [29] J. Mudrock, M. Chase, I. Kadera and T. Wagstrom, *A note on the equitable choosability of complete bipartite graphs*, Discuss. Math. Graph Theory, in press.
doi:10.7151/dmgt.2232
- [30] K. Nakprasit, personal communication, 2002.

- [31] S.V. Pemmaraju, *Equitable colorings extend Chernoff-Hoeffding bounds*, Proceedings of the 5th International Workshop on Randomization and Approximation Techniques in Computer Science (2001) 285–296.
- [32] G. Sathiamoorthy and T.N. Janakiraman, *Graceful labeling of generalized theta graphs*, Nat. Acad. Sci. Lett. **41** (2018) 121–122.
doi:10.1007/s40009-018-0625-2
- [33] A. Tucker, *Perfect graphs and an application to optimizing municipal services*, SIAM Rev. **15** (1973) 585–590.
doi:10.1137/1015072
- [34] V.G. Vizing, *Some unsolved problems in graph theory*, Ups. Mat. Nauk. **23** (1968) 117–134, in Russian, English translation in Russian Math. Surveys **23** (1968) 125–141.
doi:10.1070/RM1968v023n06ABEH001252
- [35] V.G. Vizing, *Coloring the vertices of a graph in prescribed colors*, Diskret. Analiz. **29**, Metody Diskret. Anal. v Teorii Kodovi Skhem **101** (1976) 3–10.
- [36] W.-F. Wang, *Equitable total coloring of graphs with maximum degree 3*, Graphs Combin. **18** (2002) 677–685.
doi:10.1007/s003730200051
- [37] D.B. West, *Introduction to Graph Theory* (Prentice Hall, NJ, 2001).
- [38] H.P. Yap and Y. Zhang, *The equitable Δ -coloring conjecture holds for outerplanar graphs*, Bull. Inst. Math. Acad. Sin. **25** (1997) 143–149.
- [39] Z. Zhang, W. Wang, S. Bau and J. Li, *On the equitable total colorings of some join graphs*, J. Inf. Comput. Sci. **2** (2005) 829–834.
- [40] J. Zhu and Y. Bu, *Equitable list coloring of planar graphs without short cycles*, Theoret. Comput. Sci. **407** (2008) 21–28.
doi:10.1016/j.tcs.2008.04.018
- [41] X. Zhang and J.-L. Wu, *On equitable and equitable list colorings of series-parallel graphs*, Discrete Math. **311** (2011) 800–803.
doi:10.1016/j.disc.2011.02.001
- [42] J. Zhu and Y. Bu, *Equitable and equitable list colorings of graphs*, Theoret. Comput. Sci. **411** (2010) 3873–3876.
doi:10.1016/j.tcs.2010.06.027
- [43] J. Zhu, Y. Bu and X. Min, *Equitable list-coloring for C_5 -free plane graphs without adjacent triangles*, Graphs Combin. **31** (2015) 795–804.
doi:10.1007/s00373-013-1396-7

Received 22 August 2019
 Revised 7 December 2019
 Accepted 10 December 2019